

1



ICHES AUSTRALIA 2011

3	AUTHOR INDEX					
3						
3	lsikwue, Bernadeth C	739	Kim, JunYoung	75		
1	lwabuchi, Yuhko	726	Kim, Sung-Soo	67		
	Iwakuma, Maki	532 708 756	Kim, Sungwhan	65		
3	Jaain, Monicca	612	Kim, Sunme	31		
7	Jackson, Shya	528	Kim, Taeyoun	65		
-	Jang, Jaewoong	650	Kim, Wao Jeong	64		
1	Jayasuriya, Niranjali	441	Kim, Woo-Jeong	64		
-	Jeon, Moon-Jin	615	Kimura, Ryo	532 690 75		
3	Ji, Na Young	671	King, Rudith S	34		
1	Johnson, Laura	614	Kioni, Paul N	66		
	Jollands, Margaret	310	Kissho, Mizue Y	69		
1	Jones, Lorie	513	Kline, Michael	35		
	Jung, Myung Hee	613	Knox, Sally A	35		
7	Kaberere, Keren K	426	Koba, Alexandra	61		
3	Kadiri, Medina O	645	***************************************			
	Kagira, John M	656	Kongoro, Jedida A	60		
1	Kahangi, Esther M	659	Kosoko, Adebowale O	45		
7	Kaib, Manfred	667	Koth, Barbara	41		
- 44	Kakota, Tasokwa VM	689	Kovalcuk, Tatiana V	62		
7	Kakuda, Fedeko	708	Kubo, Nami	532 690 75		
=	Kallis, Deon	408	Kucharska, Danuta J	73		
7	Kalungu, Jokastah Wanzuu	646 731	Kumar, Ravinder	64		
7 '	Kang, Sang Hun	653	Kusumayudha, Sari B	53		
	Kang, Taejin	658	Kwemesi, Anthony A	45		
3	Kang, Yun Hee	671	Kwon, Dong-Young	61		
2	Kanui, Mary Ng'endo	732	Kwon, Jang Hyuk	41		
	Kanyama-Phiri, George Y	657	Lake, Amanda			
3	Kao, Nhol	668	Lamont, Miranda J	69		
	Karanja, Faith Njoki	330 416	Lane, Augusta	35		
7	Karanja, Simon M	656	Lang'at Thoruwa, Caroline C	69		
7	Kata, Lauren A	327	Laryea, Wilhelmina k	66		
-14	Kato, Maki	428	Layne, Peggy	35		
T	Kentish, Sandra E	516	Leclercq, Séverine	73		
	Keraita, James	660	Lee, Ah-Reum	64		
R	Khare, Manorama M	347 709	Lee, Chung II	67		
7	Khrutskaya, Hanna	710	Lee, Geumseon	65		
	Kim, Day-Young	615	Lee, Hee Gu	67		
T	Kim, Deog-Keun	452	Lee, Hosun	65		
	Kim, Eunhee	722	Lee, Jin-Suk	45		
T	Kim, Gyu-Sun	615	Lee, Jonghyun	31		
T	Kim, Heeyeon	647	Lee, Na-Young	61		
	Kim, HyeYoung	753	Lee, Sookyeon	65		
	Kim, Hyong-Ha	648 649	Lee, Wonkyung	31		
7	Kim, Jae Wha	671	Lee, Yang Ji	65		
4	Kim, JiHye	753	Leggon, Cheryl B	52		
3	Kim, Jung Sun	333 700	Leibnitz, Gretalyn M	34		



#### LIST OF PAPERS

#### **ENERGY, RESOURCES & ENVIRONMENTAL SCIENCE & TECHNOLOGY**

COMPARISON OF SOLAR SORPTION COOLING SYSTEMS USING TRNSYS SOFTWARE (Marie-Esther Guillerm, Rebecca Selwyn, Hind Saidani-Scott)

KINGSGROVE TO REVESBY RAIL QUADRUPLICATION – INNOVATIONS IN DESIGN AND CONSTRUCTION OF THE OVERBRIDGES (Gillian Sisk, Sally Cox, Naomi Stone)

TOWARDS A MORE SUSTAINABLE, RESILIENT INFRASTRUCTURE SYSTEM: REGIONAL RISK ASSESSMENT OF COASTAL BRIDGES DURING HURRICANE EVENTS (Candase D. Arnold, Jamie E. Padgett)

CHALLENGING DESIGN: FOUNDATIONS FOR TALL BUILDINGS (Helen Chow, Frances Badelow)

ACCELERATED DRAINAGE IN SAND PILES BY SUCTION THROUGH CONVOLUTED PERFORATED PIPE (Eliza Hashemi, Muhammed A. Bhuiyan, Niranjali Jayasuriya)

RULES FOR THE PROVISION OF EARTHQUAKE RESISTANCE IN SMALL BUILDINGS IN GHANA (C.D. BOU-Chedid)

THE PRODUCTION OF BIODIESEL FROM OILS AND FATS (Ji-Yeon Park, Jin-Suk Lee, Deog-Keun Kim, You-Kwan Oh)

ENGINEERING SUSTAINABILITY – A SYSTEMS APPROACH (Lorie Jones FIEAust, CPEng, NPER, Janine Barrow, Katie DeJong)

DEMONSTRATING CARBON DIOXIDE CAPTURE THROUGH PILOT PLANT OPERATION (Clare J Anderson, Kathryn A Mumford, Kathryn H Smith, Sandra E Kentish, Abdul Qader and Geoff W Stevens)

STRUCTURE, HYDROGEOLOGY, AND THE GEOTHERMAL SYSTEM OF MOUNT UNGARAN AREA, CENTRAL JAVA, INDONESIA (Sari B. Kusumayudha, Andriyanto P. Markam, Basuki Rahmat)

BARELY SURVIVING OR TRAILBLAZING? PROFESSIONAL LIFE OF WOMEN ENGINEERS IN SHIPBUILDING, MECHANICAL AND CONSTRUCTION INDUSTRIES IN TAIWAN (Wen-Ling Hong)

# STRUCTURE, HYDROGEOLOGY, AND THE GEOTHERMAL SYSTEM OF MOUNT UNGARAN AREA, CENTRAL JAVA, INDONESIA

Sari B. Kusumayudha<sup>1</sup>, Andriyanto P. Markam<sup>2</sup>, Basuki Rahmat<sup>3</sup> UPN "Veteran", JI SWK 104 Yogyakarta, Indoesia. Email: saribk@gmail.com

## ABSTRACT

Indonesia is well known as a country of volcanoes. There are hundreds active and dying volcanoes, one of which is namely Mount Ungaran. It is a Pleistocene age volcano, located in the Semarang region, Central Java Province. In the present time the volcano is identified being in its post volcanism stage. Some observable fact related to the stage of the volcano is there has been no more volcanic eruption record for a thousand year. In contrary, there exist fumaroles and sulfuric hot steam through fissure on the flank of the volcano. The geologic structures obtained at the surrounding area are reverse faults, strike slip faults, and normal faults. On the other hand, the hydrogeologic system is compiled of fractured complex aquifer system, containing volcanic breccias, sandstones, and lava deposits of andesitic to basaltic composition. Due to the high density of fractures, cracks, and faults, the aquifers are able to transmit surface water and groundwater down to contact with such hot plutonic bodies in the volcanic environment to perform geothermal system. This geothermal system is expressed by the occurrence of side and fissure sulfuric fumaroles eruption. Therefore it can be stated that structures and hydrogeologic condition play important role to the occurrence of the geothermal system in the study area.

# Introduction

Mount Ungaran is located at Central Java province, belonging to the Semarang region, Indonesia. The volcano is Pleistocene (about 1 to 1.8 million years) age, and in post volcanism phase, indicated by the existence of fumaroles on its side vent. Ancient Mount Ungaran volcanism was the source of Notopuro Formation, comprises of andesitic to basaltic breccias, sandstones and agglomerates. Geothermal system is generally influenced by geologic structure, and hydrogeologic condition. Therefore, based on the rationales mentioned above, this paper is intended to assess the geologic structures and hydrogeologic condition those control the geothermal system of Mount Ungaran environment.

The area being studied is situated on the southeastern slope of Ungaran volcano, administratively belonging to the Semarang region, Central Java Province. It is astronomically bordered by  $7^{\circ}10'00"$  to  $7^{\circ}15'39"$  latitudes, and  $110^{\circ}22'30"$  to  $110^{\circ}28'32"$  longitudes, able to be reached from Semarang, the capital city of Central Java province, of about 27 km to the south, and from Yogyakarta about 70 km to the Northwest (Figure 1).

Method applied in order to complete the objectives of this study is geologic and hydrogeologic mapping, geologic structural and hydrochemistry analyses. Some rock and groundwater samples were taken for either in situ or laboratories testing.




Figure 1. The location map of the study area

### Geology

Van Bemmelen (1949) divided the physiography of the Central Java into several zone, respectively from the north to the south are alluvial plain of Northern Java, North Serayu Anticlynorium, South Serayu Mountains, Central Depression and Quaternary Volcanoes Zone, and Southern Mountains. The study area that is situated on the southeastern slope of Mount Ungaran, physiographically includes in the North Serayu Mountains (Figure 2). The Ungaran volcano stands on the eastern most edge of uplifted mountains, and separates North Serayu Mountains in the west from Kendeng Ridge in the east (East Java). Both the North Serayu and Kendeng Mountains are folded ridges, built of predominantly plastic sedimentary rocks of Tertiary Neocene ages (Van Bemmelen, 1949). Because of occupying such foot slopes of a volcano, landforms in the study area are generally expressing hills and wavy planes.

The oldest lithology outcropped in the study area is Penyatan formation, comprises of andesitic breccias with lava interbedding, calcareous sandstones, and marl, Middle Miocene to Late Miocene ages (Van Bemmelen, 1949, Genevraye & Samuel 1972, Markam 1998). Penyatan Formation is unconformably overlain by Notopuro Formation. Notopuro formation consists of andesitic to basaltic breccias, lavas, volcanic sandstones, and tuff layers. The breccias were deposited by both laharic and pyroclastic mechanism. This volcanic formation at some locations was intruded by igneous rocks of either andesites or basalts. The age of Notopuro formation is Late Pliocene to Pleistocene (Van Bemmelen,

1949). The youngest lithology of the study area is alluvial deposits, consists of unconsolidated sediments of clay, granule, pebble, cobble, to boulder grain sizes. The alluvial have been produced from fluvial sedimentation of some rivers, such as River Panjang, River Garang, and River Santritulung.




Figure 2. The Physiographic map of Central Java (Van Bemmelen, 1949)

# Structural Geology and Hydrogeology

The study area shows a complex geologic structure of folds and faults. Formation of Tertiary period is strongly folded, while litologic unit of Quaternary period is in general dissected by faults. In spite of being controlled by regional tectonic regime of Java island during Tertiary to Quaternary period, the geologic structures in the southeastern slope of Mount Ungaran is also influenced by local gravitational tectonic processes of Pleistocene to Holocene (Van Bemmelen, 1949). The Quaternary gravitational phenomenon created the collapse of Ungaran volcano, and reactivated some faults those already sliced the plastic formations underlain by the massive volcanic deposits of Notopuro Formation (Van Bemmelen, 1949).

Three great faults crosscutting the study area are the Ungaran reverse fault, Jimbaran normal fault, and Getas left strike slip fault. Beside the three faults, there are normal faults as the impact of gravity force with relatively north – south strikes. On the other hand the reverse fault is west – east strikes, and the left strike slip faults is northeast – southwest orientation. Fault type determination was done by using field data and joints analysis, referring to Rickard classification (Billing, 1979). Field data used to determine the classification of the faults are written in the Table 1, while the geologic condition of the study area can be seen in the Figure 3.

Fault Name	Classification	Field Supporting Data	
Getas Fault	Left strike slip fault	<ul> <li>Scarps along the Santritulung river</li> <li>Fault breccias at river Santritulung, Begajah, and river Klampok</li> <li>Lithologic offset, and shear &amp; gash fractures</li> </ul>	
Jimbaran Fault	Normal fault	<ul> <li>Fault plane with slicken sides</li> <li>Fault breccias, shear and gash fractures</li> <li>Lithologic offset</li> </ul>	
Ungaran Fault Reverse fault		<ul> <li>Fault plane with slicken sides</li> <li>Scarps, water fall, and fault springs</li> <li>Brecciated zones,</li> <li>Lithologic offset,</li> <li>Minor faults, and drag folds</li> </ul>	

Table 1. Field supporting data to fault classification of the study area




Figure 3. Geologic block diagram of the southeastern slope of Mount Ungaran (Markam, 1998 revised by Kusumayudha et.al, 2011)

There are some normal ring faults surrounding the cone of the volcano (Figure 4.). By the control of these faults, the summit part of Mount Ungaran was subjected to gravitational

subsidence (Van Bemmelen, 1949, Zen, 1969, vide Zen, et.al, 1983). Impact of this collapse, magma extruded out to the surface forming some lava domes around the summit of the volcano (Figure 5a.). Despite of the existence of ring faults, there are radial faults which play as the vent of side or flank eruptions. Up to now, fumaroles, sulfuric hot steam and geyser extrude from the flank vent as geothermal energy potency (Figure 5b.).




Figure 4. Schematic block diagram of Ungaran volcano-tectonic (Van Bemmelen, 1949)



Figure 5. a. Lava dome, b. Fomarole, at Gedongsongo, southern flank of Mount Ungaran

The hydrogeologic system of Ungaran area is built by aquifers comprises of volcanic deposits especially andesitic breccias, lava, and sandstone, predominantly forming fissure type of water bearing formation. In this system, Penyatan formation plays as the basement. Based on their stratigraphic position, Ungaran aquifers can be classified into unconfined and semi confined aquifers. Groundwater dynamics in the Ungaran aquifer is very complex, because it is conducted by crack and intergranular porosity systems combination. Under the hydraulic gradient, groundwater moves in general relatively from west to east direction by complex diffuse and cracked flow type.

Recharge zone occupies almost all parts, while the discharge zone lay on the eastern parts of the study area. There are many springs, and some of them are containing sulfur of moderately high concentration, and relatively high temperature. The existences of some hot and sulfuric springs show that the volcanism of Ungaran in the present time is progressing for post volcanism stage. Figure 6 displays the hydrogeologic conceptual model that is controlling the geothermal system of the southeastern slope of Mount Ungaran.




Figure 6. Hydrogeologic conceptual model of Mount Ungaran southeastern slope area (Markam ,1998 revised by Kusumayudha et.al, 2011)

The properties and groundwater chemistry of the aquifers are shown in the Table 2 and Table 3. The data were obtained from field measurements, field sampling, and laboratory testing. Result from hydrochemistry analysis using Aquachem software show that groundwater of Ungaran aquifer belongs to various water types. Based on cation content, it can be classified into calcium dominant water, on the other hand according to anion content the water includes in bicarbonate water, although some samples indicating chloride water (See Table 3 and Figure 7, the Piper plot of groundwater chemistry).

Hydraulic Conductivity	Transmissivity	Effective Porosity	Storage
(m/sec)	(m2/sec)		Coefficient
$0.9 \cdot 10^{-5}$ to $1.9.10^{-5}$	$1.3.10^{-4}$ to $3.3.10^{-2}$	10% – 25%,	$1.10^3$ to $1.7.10^{-1}$

Sample	Location	Hydroche	lydrochemistry Concentration (Mg/l)			Watan Tuna		
ID	(Village)	Ca	Mg	Na	HCO3	SO4	Cl	Water Type
U1a	Jimbaran 1	56.12	14.59	27.59	256.25	19.21	14.18	Ca-Mg-Na-HCO3
U1b	Jimbaran 2	56.14	14.58	27.58	256.25	19.20	14.19	Ca-Mg-Na-HCO3
U2a	Karangjoho1	24.05	14.60	62.07	189.14	9.61	56.72	Na-Mg-Ca-HCO3-Cl
U2b	Karangjoho 2	26.05	13.38	48.28	192.19	9.61	35.45	Na-Ca-Mg-HCO3-Cl
U3	Pakopen	24.05	19.45	50.57	231.84	19.22	24.81	Na-Mg-Ca-HCO3
U4	Kalikembar	36.07	17.02	34.48	247.11	4.80	14.18	Ca-Na-Mg-HCO3
U5	Sruwen	76.15	13.37	29.89	317.26	19.20	10.64	Ca-Na-HCO3
U6	Coblong	26.05	13.38	20.69	167.78	9.60	7.09	Ca-Mg-Na-HCO3
U7a	Diwak 1	28.06	13.37	29.88	184.26	9.61	17.72	Ca-Na-Mg-Cl
U7b	Diwak 2	36.07	17.02	41.38	222.69	9.60	35.46	Ca-Na-Mg-HCO3-Cl

Table 3. Groundwater chemistry of the Ungaran aquifers






Figure 7. Piper diagram plot of groundwater samples of Ungaran aquifers

#### Geothermal System of Mount Ungaran

The earth anatomy from the outer to the inner is composed of crust, mantle, and core respectively. It owns and yields natural heat. Increase of the temperature layers with depth toward the earth core is called geothermal. The temperature of the geothermal has been resulted from heat residue of the initial earth material, and decay of radioactive compounds. The rate of the temperature change in the earth crust per km deep is known as geothermal gradient. Geothermal gradient causes the melting rocks to form such very hot and melted materials in the earth mantle. When hot and melted materials of the earth mantle rise to intrude the crust, it is called magma. If the magma extrude the surface, it becomes change its name to lava.

When water from the earth surface infiltrate subsurface penetrating soil layers reaching the saturated zones, it will forms groundwater. Due to high permeability of the rock formation and particular favorable condition, the groundwater will gravitate down to enter deeper formations, contact to such a high temperature rock body. The water then will be heated, changing its phase to be steam. When the surrounding rock formations are fissured intensively by joints and cracks, the condition will be able to function as a regulation for the steam out flowing from its traps. The steam mixed with such volcanic gases is potentially flushed up and out of the land surface, known as fumaroles that can be geothermal energy resource. The model of geothermal system is shown in the Figure 8 bellow.




Figure 8. Geothermal system model (http://www.indotoplist.com/info)

On the south flank of Mount Ungaran, i.e. Gedongsongo area, there is fumarole through fissures, composed of water vapor and such gases as  $CO_2$ ,  $H_2S$ ,  $H_2$ , and  $CH_4$  (Wahyudi, 2006), with temperature of about 85°C. Water samples taken from Gedongsongo in general show very low ph < 4.0. This high acidity of the water is caused by geochemical reaction of water with some gasses such as  $CO_2$ , and Sulfur. Source of the erupted steam is groundwater in the hydrogeologic system of Ungaran volcano and surrounding area,

including the southeastern slope parts of the volcano which strongly controlled by cracks, fissures, and faults.

Due to the influence of geologic condition as has been discussed earlier, there are weak zones performed from the existence of reverse Ungaran Fault, strike slip Getas Fault, Jimbaran Fault, and other normal faults with all of their secondary order structures of joints and fractures. The weak zones cause the aquifer more permeable. The existence of such a caprock makes the steam accumulated, causing its volume and pressure increase. Lithology that able to function as a caprock is clay or marl. As it was mentioned above, there are marl layers in Penyatan formation. When there is crack system able to be the regulator, the steam will be sprayed out, known as fumaroles. The geothermal potency of Mount Ungaran is assumed to be sustainable, because it is supported by the relatively high precipitation in the surrounding area, i.e. 2,247 mm/year average (www.esdm.go.id).

# Conclusion

Based on the analysis and assessment written above, the conclusions are as the following:

- 1. Geology of the study area is composed of Tertiary Penyatan Formation, unconformably overlain by Pliopleistocene Notopuro Formation. Penyatan Formation comprises of sandstone, breccias, and marl, Notopuro formation consists of andesitic to basaltic breccias, lavas, volcanic sandstones, and tuffs. The formations are intruded by andesites and basalts.
- 2. Geologic structures in the study area are folds and faults. The folds exist in Penyatan Formation, while the faults cut up Notopuro Formation. The three main faults are Ungaran reverse fault, Getas left strike slip fault, and Jimbaran normal fault.
- 3. The hydrogeologic system of the Ungaran southeastern slope is built of complex fissures aquifer system composed of volcanic rocks of Mount Ungaran. Types of groundwater are variation of Ca-Mg-Na-HCO3, Na-Ca-Mg-HCO3-Cl, Ca-Na-HCO3, and Ca-Na-Mg-Cl. The structural hydrogeologic system let groundwater of the surrounding area to contact with magmatic and plutonic bodies of the volcanic environments to perform geothermal system.
- 4. The geothermal system of Mount Ungaran southeastern slope is expressed as flank and fissure fumaroles eruption. The fumaroles consist of water vapor, CO<sub>2</sub>, H<sub>2</sub>S, H<sub>2</sub>, and CH<sub>4</sub> with 85°C temperature.

### References

Billing, M.P., 1979, Structural Geology, 3rd ed., Prentice Hall of India, New Delhi, 606 p

- Budiardjo, B., Nugroho M, Budihardi, 1997, Resource Characteristics of the Ungaran Field, Central Java Indonesia, *Proc. Seminar of Human Resources of Indonesian Geologists*, UPN "Veteran" Yogyakarta, p9
- Domenico, P. A. & Schwartz, F. W., 1990, *Physical and Chemical Hydrogeology*, John Wiley and Sons
- Freeze, R. A. & Cherry, J. A., 1979, *Groundwater*, Prentice Hall Inc, Englewood Cliffs, New Jersey, 604 p.

- Hem, J. D., 1985, Study and Interpretation of the Chemical Characteristics of Natural Water, USGS Water Supply Paper 2254
- Kusumayudha, S. B., 1993, Groundwater Geochemistry of Mount Merapi's Southern Slope Area, Java, Indonesia, Master of Science Thesis, AIT, Bangkok, unpublished, 239 p.
- Kusumayudha, S.B. & Pratiknyo P., 2001, Detecting the existence of groundwater discharge in the fractured volcanic area. *Majalah Geologi Indonesia*, Vol 16 No 2, p 83 91
- Markam, A. P., 1998, "Geologi dan Studi Aplikasi Sistem Informasi Hidrogeologi pada Endapan Vulkanik Kuarter Sistem Akifer Ungaran daerah Jimbaran – Ungaran dan Sekitarnya, Kecamatan Ungaran, Kabupaten Semarang, Jawa Tengah – Indonesia", Thesis, UPN "Veteran" Yogyakarta, Unpublished, 206 p.
- Matthess, G., 1982, *The Properties of Groundwater*, John Wiley \$ Sons Publishers, New York, 406 p.
- Pertamina, 1976, "Stratigrafi Zona Bogor Serayu Utara Daerah Semarang Baratdaya Jawa Tengah", Divisi Eksplorsi Migas, Jakarta, Indonesia, unpublished, 45 p.
- Samuel, L. & Genevraye, 1972, Geology of the Kendemg Zone (Central and East java), *Proc. 1<sup>st</sup> Annual Convention of IPA*, Jakarta, Indonesia,
- Samuel L., Yanto, R., Sumantri, Fx, Suyanto, Sudiono, A., Syhbudin, A., 1995, *Indonesia Geothermal Reserves and Resources*, Pertamina, Jakarta, 45 p
- Todd, D.K., 1980, *Groundwater Hydrology*, 2<sup>nd</sup> edition, John Wiley & Sons, New York., 535 p.
- Van Bemmelen, R.W., 1949, *The Geology of Indonesia*, Vol IA, Martinus Nijhoff, The Hague, 792 p.
- Wahyudi, 2006, "Kajian Potensi panasbumi dan rekomendasi Pemanfaatannya pada daerah Prospek Gunungapi Ungaran, Jawa Tengah, Berkala MIPA, Vol 16 (1), Januari 2006, p 41 - 48
- Zen, M. T., M.A. Syarif, S.H. Simatupang, G. Yuniarto, 1983, Tektogenesa Gaya Berat dan Daur Magma Sepanjang Deretan Gunungapi Ungaran – Merapi di Jawa Tengah, Proc IAGI Annual Convention, Yogyakarta p 76 - 85

http://www.esdm.go.id

http://www.indotoplist.com/info

http://www.volcano.si.edu/world/volcano