



ISSN 2373-423X

# JOAAT



[www.joaat.com](http://www.joaat.com)



ISSN: 2373-423X (Print)

ISSN: 2301-3737(Online)

**Journal of**  
**Advanced Agricultural Technologies**

Vol. 5, No. 2, June 2018

**Editor-in-Chief: Prof. Byoung Ryong Jeong**

The logo for the Journal of Advanced Agricultural Technologies (dAAT) features a stylized green wave symbol to the left of the lowercase letters 'dAAT'. The 'd' is a simple curve, while 'AAT' are bold, block letters. Below the logo, the full journal title is written in a smaller, sans-serif font.  
**dAAT**  
Journal of Advanced Agricultural Technologies

# Journal of Advanced Agricultural Technologies

## CONTENTS

Volume 5, Number 2, June 2018

### Food Science and Engineering

Effects of Maltodextrin and Silicon Dioxide Added as Anticaking Agents on the Properties of Instant Date Palm (*Phoenix dactylifera* L.) Powder Using Spray Drying 86

*Nutthamon Nortuy, Kanyarat Suthapakti, and Niramom Utama-ang*

Detection of Salt Content in Raw Milk on a Gold Electrode through Cyclic Voltammetry 93

*Chen Xing-guang, Chu Bei-bei, Lai Li-zhi, Peng Hui-Jeng, Deng Dan-wen, and Huang Gan-hui*

Effect of Frying in Different Cooking Oils on the Fatty Acid Profile of Nile Tilapia (*Oreochromis niloticus*) Fillets 98

*Dinesh D. Jayasena, Kalana Fernando, and Thilini Awanthika*

How to Display Social and Environmental Sustainability of Food Products? 103

*Fatiha Fort and Laura Solaroli*

### Animal Science

Performance and MilleAnti-Oxidant Property of Dairy Goats Fed Pomegranate Seed Pulp and Soybean Oil .109

*Jongkolnee Yaowapaksophon*

Antioxidant Effects of Peppermint (*Mentha piperita*) Extract on the Oxidative Balance of Rabbit Spermatozoa ..... 117

*Eva Tvrdá; Natalia Konecnd, Katarina Zbyiiovska, and Norbert Lukad*

Goat Farm Performance in Northern Mindanao, Philippines 123

*Reynaldo L. Intong, Julnar E. Pepito, Jhonny M Callejo, and Emilio M Cruz*

### Botany and Biotechnology

Effect of Compost and Different NPK Levels on Growth and Yield of Three Varieties in Sri Lanka

*HASL Jayasinghe and ANR Weerawansha*

Effectivity of *Trichoderma Asperellum* against Water Availability Level in

*Nirwana and St. Sabahannur*

Bioremediation of Mine Waste Using Sulphate Reducing Bacteria againts

*Saida, Netty, Abdullah, and St. Subaedah*

Yield and Phytochemical Attributes of Ashwagandha (*Withania somnifera*) of Nutrients

*Ewon Kaliyadasa, Lalith Jayasinghe, and Sriyani Peiris*

Growth and Yield of Hybrid Com under Different Fertilizer Applications

*Raden R. R. Brotodjojo and Dyah Arbiwati*



# Growth and Yield of Hybrid Corn under Different Fertilizer Applications

Raden R. R. Brotodjojo and Dyah Arbiwati

Agrotechnology, Universitas Pembangunan Nasional "Veteran" Yogyakarta, Yogyakarta, Indonesia

Email: brotodjojo@upnyk.ac.id, arbi\_rizal@yahoo.com

**Abstract**—Conventional agriculture is heavily depended on the use of inorganic fertilizer that provides sufficient macro nutrients, with some draw back on the environment. On the other hand, organic agriculture that only use organic fertilizers is considered environmentally friendly but sometimes could not provide adequate nutrients for plants. The aim of this research was study the growth and yield of hybrid corn treated with different dose of Granular Organic Fertilizer (GOF) in comparison to inorganic fertilizers. The corns were planted in polybags arranged in Randomized Completely Block Design with four treatments, Le, 10 tons/ha GOF, 20 tons/ha GOF, 30 tons/ha GOF, inorganic fertilizers (630 kg/ha ~ 350 kg/ha Urea, 100 kg/ha SP36, 380 kg/ha Phonska). Each treatment consisted of 20 plants with 6 replications. The results showed that the height of plants treated with 20 tons/ha GOF or 30 tons/ha GOF was not significantly different from plants treated with inorganic fertilizers, but was significantly higher than those treated with 10 tons/ha GOF. The number of leaves did not significantly affected by different fertilizers application. Furthermore, plants treated with inorganic fertilizers significantly had higher chlorophyll content and produced higher yield than those treated with GOF at various doses. This revealed that application of GOF on corn at dose of up to 30 tons/ha was not sufficient to produce high yield. To increased yield, higher dose of GOF is needed or it can be supplemented by application of inorganic fertilizers.

**Index Terms**—granular organic fertilizer, inorganic fertilizer, corn, growth, yield

## I. INTRODUCTION

Com is one of important grains in the worlds. It is cultivated for food, feed and recently for renewable fuels in many countries. The world com production is estimated to increase from 968,06 Million Metric Tons (MMT) in 2015/2016 to 1065,11 MMT in 2016/2017 ~d world consumption increase from 965,1 MMT in 2015/2016 to 1053,59 MMT (est.) in 2016/2017. In addition, the com demand is projected to increase to 1062,3 MMT in 2017/2018, whereas the production is projected only 1033,66 MMT [1]. The United States of America is the highest producer of com in the world with its production is estimated 384,778 MMT in 2016/2017. Indonesia is in the 12th position produced 10,200 MMT (est.) in 2016/2017 [2]. Com is usually planted as a second crop after rice in Indonesia. Com consumption in

Indonesia as food and feed in 2016/2017 is estimated 3,8 MIT and 8,4 MTI, respectively [3]. The government of Indonesia has restricted the import of corn that implicated to boost the country production.

In general com is cultivated conventionally with the intensive application of inorganic fertilizers. In Indonesia the recommended fertilizer dosage for com are N: 65-95 kg/ha/season, P: 30-50 kg/ha/season, K: 10-30 kg/ha/season [4]. However, some time, farmers use fertilizer under or beyond recommended rates. There is a great concern that over use of inorganic fertilizers, continuously for long period unaccompanied by organic materials has adverse effect to the environment, including degradation of soil quality and pollution. Degraded soil characterized by compacted soil and reduced porosity, decreased soil fertility and content of essential micro nutrients [5]. In addition, soil treated with inorganic fertilizer alone significantly has lower populations of soil microbial biomass, such as bacteria, fungi and actinomycetes, as well as soil enzyme activities [6], [7]. Decreasing soil quality can, cause levelling off crop yield, although farmers use more fertilizer for their crop the yield remain the same or sometimes decrease.

In comparison, organic fertilizer poses many advantages including (1) provides a more balance macro and micro nutrients, (2) improves activities of soil microbes, (3) increases P supply by improves mycorrhizae colonization, (3) improves soil structure thus enhance growth of roots, (4) increases Soil Organic Materials [8]. The positive impact of organic fertilizer seen in the short term and long term, which increase soil fertility and crop yields. In the short term, organic fertilizer application can improve the biological and biochemical properties of the-soil, so as to produce the similar yield of sweet corn as that treated with inorganic fertilizer [9]. A 32-year study in Sweden showed that the use of organic fertilizers would yield wheat and potatoes approximately the same as those using inorganic fertilizers. Increased yields on land given organic fertilizers were even higher than those given inorganic fertilizers. The quality of wheat and potatoes treated with organic fertilizers is better than those given inorganic fertilizers, in addition to the decrease in weight after storage is lower in wheat and potatoes given organic fertilizers than those given inorganic fertilizers [10]. Moreover, organic crops contained significantly more vitamin C, iron, magnesium, and phosphorus and significantly less nitrates than conventional crops [11].

Manuscript received September 13, 2017; revised January 23, 2018.

The nutrients content of organic fertilizer depend on the materials that are used as main substrate. Organic fertilizer made from cattle manure and broiler litter contains 1.5% N, 1.5% P<sub>2</sub>O<sub>5</sub>, 1.2% K<sub>2</sub>O, 1.1% Ca, 0.3% Mg and 3% N, 3% P<sub>2</sub>O<sub>5</sub>, 2% K<sub>2</sub>O, 1.8% Ca, 0.4% Mg, respectively [12]. Boiler ash as the residue of bagasse combustion produced in sugar production process can be added as organic materials, it contains 71% SiO<sub>2</sub>; 3% P<sub>2</sub>O<sub>5</sub>; 8.2% K<sub>2</sub>O; 3.4% CaO and some micronutrients (0.2% MnO; 1.9% Al<sub>2</sub>O<sub>3</sub>, 7.8% and 0.3% Fe<sub>2</sub>O<sub>3</sub> MgO) [13].

As source of organic materials, boiler ash contains high percentage of silicate that improve plant resistance against biotic and abiotic stresses. Plants from the Gramineae family, including corn, are Silicate accumulators (Si). The absorbed silicate acid is stored in the epidermal tissue of the plant thereby strengthening the stems and leaves [14]. Silicate can increase the efficiency of plant photosynthesis. The application of boiler ash with dose of 50 ton / ha can increase soil porosity, P, S and K, Fe, Mn, Zn and Cu availability as well as grain yield on calcareous soil [15].

However, constraints perceived by farmers with the use of organic fertilizers are it is to be given in large quantities because of low nutrient content, need more cost to transport and not practical. Farmers are less responsive in accepting the transformation of the idea of the benefits of solid organic fertilizers, not even believing that organic fertilizer is a fundamental component of soil fertility improvement as well as plant growth and yield. Research on the application of organic fertilizers to the growth and yield of various plants has been done. Nevertheless, research on the use of granular organic fertilizers enriched with boiler ash and neem leaves powder to increase yield on corn has never been done. Therefore, the aim of this research was to compare the growth and yield of hybrid corn treated with various dosage of granular organic fertilizers with inorganic fertilizer.

## II. MATERIALS AND METHODS

This experiment was conducted in The Experimental Field Station of Faculty of Agriculture, Universitas Pembangunan Nasional "Veteran" Yogyakarta, Indonesia. The experiment design was a Randomized Completely Block Design (RCBD) of four treatments with six replications, each treatment consisted of 20 polybags. The treatments were dose of granular organic fertilizer with control treatment according to conventional cultivation using recommended dose of inorganic fertilizers.

Granular organic fertilizer used for this experiment is specially formulated enriched with boiler ash and neem leaves powder. Granular organic fertilizer was supplied according to the treatments i.e., dose of 30 tons / ha (D1); dose of 20 ton / ha (D2); dose of 10 tons / ha (D3). For control, the plant was provided inorganic fertilizers as follows: at 15 dap (days after planting): 280 kg/ha ZA, 100 kg/ha SP36; at 35 dap: 200 kg/ ha Phonska; at 50 dap: 350 kg/ha urea, 180 kg/ha Phonska. Hybrid corns (Pioneer variety) were planted in polybags (50 cm diameter; filled with 16 kg soil). Soil used for this experiment was

obtained from location that has not been used for agricultural farming. The soil was sieved to remove the gravel. Granular organic fertilizer was placed into soil at a depth of about 10 cm two weeks before planting. Soil humidity was kept to field capacity by watering plants every day. Corn seeds were planted 2 pieces per polybag. One plant that grew unequally with others was removed from polybag after the plants were 2 wap (weeks after planting), leaving one plant per polybag.

Plant height, number of leaves and plant damage were measured every two weeks until eight weeks after planting (wap). Handheld Chlorophyll Meter (Konica Minolta SPAD-502Plus) was used to measure Chlorophyll content at 13 wap. The optical density of chlorophyll within two wavelength, blue (400-500 nm) and red (600-700nm) regions was recorded by the chlorophyll meter. The measurements showed the relative density of chlorophyll in the leaf [11]. Three leaves per plant were taken as sample to measure chlorophyll content. Yield parameters were cob fresh weight/polibag, economical weight (cob weight without sheath)/polibag that was measured at 13 wap.

The SPSS for Windows version 15 was used to analyze the data. Percentage data were transformed into  $\text{Arcsin} \sqrt{x+1}$  before being analyzed to achieve normality. All data were analyzed using a one way ANOVA. All multiple comparison tests for means were performed using Duncan Multiple Range Tests.

## III. RESULTS AND DISCUSSION

Plant height and number of leaves can reflect the relative growth of plant. Nutrient status may affect the plant growth. This experiment showed that corn plant height at 2-6 wap was not significantly affected by the application of different dose of granular organic fertilizer (GOF) or inorganic fertilizer. Hybrid corns treated with inorganic fertilizer significantly have higher plant height than those treated with 10 ton/ha granular organic fertilizer, although the plant height was not significantly different from those treated with 20 ton/ha or 30 ton/ha granular organic fertilizer (Table I). This results showed that 10 ton/ha granular organic fertilizer did not provide sufficient nutrient for plant growth. Visual observation showed that the plant vigor was low and exhibited stunted growth.

Number of corn leaves at 2-8 wap did not affected significantly by application of different dose of granular organic fertilizer or inorganic fertilizer. However, chlorophyll content in plants treated with granular organic fertilizer significantly lower than those treated with inorganic fertilizer (Table II). Leaves are important parts of plant that facilitate photosynthetic. There is a positive correlation between chlorophyll content with nitrogen content [16], [17]. Nitrogen is macro nutrient that essential for proteins synthesis and is needed to develop chloroplasts and chlorophyll [18], [19]. The colour of the leaf reflect the chlorophyll content in the leaf. In this study, plants treated with inorganic fertilizer that supplied high nitrogen, their leaves were dark green, whereas plant treated with granular organic fertilizer,



their leaves were light green indicating that they lack of nitrogen.

Fresh cob weight with or without sheath represent yield of corn. The results showed that plant treated with inorganic fertilizer produced the highest fresh cob weight with or without sheath and plant treated with 10 tons/ha granular organic fertilizer produced the lowest fresh cob weight (Table III). This research indicate that the application of granular organic fertilizer on hybrid corn with dose up to 30 ton/ha was not sufficient to achieve high yield as those with recommended dose of inorganic fertilizer. Although organic fertilizer contains macro and micro nutrient, the amount did not provide adequate nutrient needed for plant growth and yield.

Inorganic provide plants with sufficient nitrogen, phosphate and kalium. Nitrogen is needed for protein

synthesis and support the plant growth in general [19]. Phosphate is needed to generate energy, nucleic acid synthesis, carbohydrate metabolism and nitrogen fixation [20]. Kalium is needed to support the biochemical and physiological processes that influence plant growth and plant metabolism. In addition, adequate kalium will increase plant resistance against drought, salinity and pests attack [21]. Furthermore, kalium activate enzyme and support nutrient and water absorption from soil and distribute assimilates from leaf to other part of plant [22]. The photosynthetic rate is largely affected by the chlorophyll content. Plant with high content of chlorophyll as those treated with inorganic fertilizer will produce more assimilates, which subsequently will be accumulated to acquire yield. Therefore plant treated with inorganic fertilizer produced the highest yield.

TABLE I. CORN PLANT HEIGHT (CM) AT 2- 8 WAP (WEEKS AFTER PLANTING) UNDER DIFFERENT FERTILIZER APPLICATION

Fertilizer dose	2wap	4wap	6wap	Swap
30 ton/ha	29,9±1,9 a	55,9±3,6a	84,1±5,9 a	108,4±5,3 ab
20 ton/ha	27,3±1,5 a	61,2±5,8 a	83,3±5,2 a	109,33±5,3 ab
10 ton/ha	30,6±4,2 a	59,3:1:4,4a	76,0±3,8 a	95,1±10,1 b
Inorganic fertilizer	27,4±1,2 a	51,1:1:3,1 a	81,7±1,5 a	123,1:1:4,9 a

Means in the same column followed by the same letters are not significantly different (P<0.05), according to Duncan's test (a: 5%).

TABLE II. NUMBER OF CORN LEAVES AT 2-8 WAP AND CHLOROPHYLL CONTENT AT 13 MST WAP (WEEKS AFTER PLANTING) UNDER DIFFERENT FERTILIZER APPLICATION

Fertilizer dose	2wap	4wap	6wap	Swap	Chlorophyll (SPAD value)
30 ton/ha	2,4±0,1 a	4,7±0,2 a	7,3±0,2 a	9,1±0,1 a	22,1±1,9b
20 ton/ha	2,9±0,3 a	5,1±0,2 a	7,7±0,2 a	9,2:1:0,4a	24,1:1:2,4 b
10 ton/ha	3,0:1:0,2 a	4,9±0,4a	6,7±0,0 b	9,1:1:0,6 a	24,2:1:3,5 b
Inorganic fertilizer	2,9±0,1 a	4,8±0,3 a	7,3±0,2 a	9,8±0,4 a	44,5±2,7 a

Means in the same column followed by the same letters are not significantly different (P<0.05), according to Duncan's test (a: 5%). SPAD value: Index shown by Konica Minolta SPAD-502Plus that indicating the content of chlorophyll in the leaf

TABLE fii. CORN YIELD (GY POLYBAG UNDER DIFFERENT FERTILIZER APPLICATION

Fertilizer dose	Fresh Cob with sheath	Fresh Cob without sheath
30 ton/ha	44,8±9,8 b	26,9±3,5 c
20 ton/ha	51,7:1:8,4b	32,3:1:1,2 b
10 ton/ha	33,5:1:9,2 c	20,0:1:1,8 d
Inorganic fertilizer	124,3±7,7 a	48,8±4,1 a

Means in the same column followed by the same letters are not significantly different (P<0.05), according to Duncan's test (a: 5%)

ACKNOWLEDGMENT

The authors wish to thank the Directorate General Research and Development Solidification, Ministry of Research, Technology and Higher Education for providing research funding through competitive grants scheme, Letter of Assignment Number: 075/SP2H/LT /DPRM/II/2016.

REFERENCES

[1] USDA. (2017). World Agricultural Supply and Demand Estimates. [Online]. Available:

<http://usdamannlib.coQ1ell.edu/usda/current/wasde/wasde-050-2017.pdf>

[2] Statista. (2017). Com production worldwide 2016/2017, by country. [Online]. Available: <https://www.statista.com/statistics/25492/global-com-production-by-country/>

[3] GAIN. (2017). Indonesia Grain and Feed Annual Report 2017 (Report No. ID1707). [Online]. Available: [https://gain.fas.usdagov/Recent/o20GAIN%20Publications/Grain%20and/o20Feed%20Annual\\_Jakarta\\_Indonesia\\_3-30-2017.pdf](https://gain.fas.usdagov/Recent/o20GAIN%20Publications/Grain%20and/o20Feed%20Annual_Jakarta_Indonesia_3-30-2017.pdf)

[4] FAO, "Fertilizer use by crop in Indonesia," Food and Agriculture Organization of the United Nations, Rome, p. 78, 2005.

[5] H. Zhang, B. Wang, and M. Xu, "Effects of inorganic fertilizer inputs on grain yields and soil properties in a long-term wheat-corn cropping system in South China," *Communications in Soil Science and Plant Analysis*, vol. 39, pp. 1583-1599, May 2008.

- [6] J. A. Adediran, L.B. Taiwo, M. O. Akande, R. A. Sobulo, and O. J. Idowu, "Application of organic and inorganic fertilizer for sustainable maize and cowpea yields in Nigeria," *Journal of Plant Nutrition*, vol. 27, pp. 1163-1181, Feb. 2007.
- [7] E. H. Chang, R. S. Chung, and Y. H. Tsai, "Effect of different application rates of organic fertilizer on soil enzyme activity and microbial population," *Soil Science and Plant Nutrition*, vol. 53, pp. 132-140, Feb. 2007.
- [8] J. H. Chen, "The combined use of chemical and organic fertilizers and/or biofertilizer for crop growth and soil fertility," in *Proc. International Workshop on Soil Stained Management of the Soil-Rhizosphere System for Efficient Crop Production and Fertilizer Use*, Thailand, 16--20 October 2006.
- [9] C. Lazcano, M. Gomez-Brandon, P. Revilla, and J. Dominguez, "Short-term effects of organic and inorganic fertilizers on soil microbial community structure and function - A field study with sweet corn," *Biology and Fertility of Soils*, vol. 49, pp. 723-733, Aug. 2013.
- [10] A. Granstedt and L. Kjellenberg, "Long-term field experiment in sweden: effects of organic and inorganic fertilizers on soil fertility and crop quality," in *Proc. International Conference in Boston, Tufts University, Agricultural Production and Nutrition*, Massachusetts, March 19-21, 1997.
- [11] V. Worthington, "Nutritional quality of organic versus conventional fruits, vegetables, and grains," *The Journal of Alternative and Complementary Medicine*, vol. 7, pp. 161-173, 2004.
- [12] C. C. Mitchell. (2008). Nutrient Content of Fertilizer Materials. Alabama Cooperative Extension System. [Online]. Available: <http://www.aces.edu/pubs/docs/N/ANR-0174/ANR-0174.pdf>
- [13] Disbunjatim. (2011). Potensi Limbah Bahan organik. [Online]. Available: [http://disbunjatim.co.cc/bahanorganik/potensi\\_limbah\\_sbg\\_bo.html](http://disbunjatim.co.cc/bahanorganik/potensi_limbah_sbg_bo.html)
- [14] N. K. Savant, G. H. Korndorfer, L. E. Datnoff, and G. H. Snyder, "Silicon nutrition and sugarcane production: A review," *Journal of Plant Nutrition*, vol. 22, no. 12, pp. 1853-1903, Dec. 1999.
- [15] M. J. Khan and M. Qasim, "Integrated use of boiler ash as organic fertilizer and soil conditioner with NPK in calcareous soil," *Songklanakarinn Journal of Science and Technology*, vol. 30, no. 3, pp. 281-289, May-Jun. 2008.
- [16] J. R. Evans, "Nitrogen and photosynthesis in the flag leaf of wheat (*Triticum aestivum* L.)," *Plant Physiology*, vol. 72, pp. 297-302, Jun. 1983.
- [17] B. Bojovic and A. Markovic, "Correlation between nitrogen and chlorophyll content in wheat (*Triticum aestivum* L.)," *Kragujevac Journal of Science*, vol. 31, pp. 69-74, 2009.
- [18] C. S. T. Daughtry, C. L. Walthall, M. S. Kim, E. B. D. Colstoun, and J. E. McMurtrey, "Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance," *Remote Sensing of Environment*, vol. 74, pp. 229-239, 2000.
- [19] D. W. Lawlor, "Carbon and nitrogen assimilation in yield: mechanisms are the key to understanding systems," *Journal of Experimental Botany*, vol. 53, Apr. 2002.
- [20] C. P. Vance, C. Uhde-Stone, and D. L. Allan, "acquisition and use: Critical adaptations by plants for nonrenewable resource," *New Phytologist*, vol. 157, Mar. 2003.
- [21] M. Wang, Q. Zheng, Q. Shen, and S. Guo, "The critical potassium in plant stress response," *International Molecular Sciences*, vol. 14, pp. 7370-7390, Apr. 2013.
- [22] P. Marschner, *Marschner's Mineral Nutrition of Higher Plants*, 3rd ed., London, UK: Academic Press, 2012, pp. 178-1



Raden R. R. Brotodjojo graduated from Faculty of Agriculture, Gadjah Mada University, Indonesia with Bachelor degree in 1989. She majored in Plant Protection and obtained her Master degree in Agricultural Entomology from The University of Sydney, Australia in 1998 and obtained her PhD in the field of Plant Protection from The University of Queensland in 2007.

She works as a lecturer in the Faculty of Agriculture, Universitas Pembangunan Nasional "Veteran" (UPNVY), Indonesia since 1990. She is a member of the Review Team at the Faculty of Agriculture, UPNVY. She is the Head of Office of International Affairs UPNVY since 2010. She teaches Introductory to Agriculture, Fundamentals of Plant Protection, Plant Pathology and Pest Management. Her research interest is in the area of plant protection, especially related to biological control and sustainable agriculture. Dr. R.R. Rukmowati Brotodjojo is a member of The Indonesian Societies of Sugar Cane Technologist, and The Entomological Society of Indonesia. She got Endeavour Award as a research fellow from the Department of Education, Employment and Workplace Relations Australia in 2009. She awarded as An Outstanding Lecturer UPNVY in 2010.

Dyab Arbiwati graduated from Faculty of Agriculture, Universitas Pembangunan Nasional "Veteran" Yogyakarta (UPNVY) with Bachelor degree in 1988. She majored in Soil Sciences, and obtained her Master degree in Soil Sciences from Gadjah Mada University in 2000. She works as a lecturer in the Faculty of Agriculture, Universitas Pembangunan Nasional "Veteran" Yogyakarta (UPNVY) since 1989. She teaches Fertilizer Technology, Soil Fertility and Introductory to Soil Sciences, Soil and Water Conservation. Her research interest is on management and improvement of soil fertility. Mrs. Dyah Arbiwati is a member of The Indonesian Society of Soil Science.