



# **Proceedings**

## **19<sup>th</sup> World Congress of Soil Science**

### **Soil solutions for a changing world**

**BRISBANE AUSTRALIA 1-6 August 2010**

**Robert J Gilkes and Nattaporn Prakongkep**

**Editors**


## Information for readers of the 19<sup>th</sup> WCSS Proceedings

The papers in the 19<sup>th</sup> WCSS Proceedings have been peer reviewed and edited prior to publication. They constitute brief (maximum 4 pages) descriptions of research work or are reviews of soil science topics. These papers should not be submitted for publication elsewhere in their present form. It is anticipated that some authors will develop and modify the content of their papers for publication as full-length papers in journals.


Opinions, statements and information within papers belong to the authors and not to IUSS or the organisers of the 19<sup>th</sup> WCSS. The 19<sup>th</sup> WCSS Proceedings are available on a DVD (ISBN 978-0-646-53783-2) and also from the website (<http://www.iuss.org>).

**Citation of the papers in 19<sup>th</sup> WCSS Proceedings should use the following formats:**

*Example of a Citation for a 19<sup>th</sup> World Congress of Soil Science Symposium*



*Example of a Citation for a 19<sup>th</sup> World Congress of Soil Science Paper*



Some papers presented at the Congress are not included in the Proceedings because *inter alia* they were incorrectly formatted, authors did not provide a corrected version, speakers enrolled for the Congress after the Proceedings were compiled or due to data base deficiencies.

In the Proceedings papers are located in Symposia current on 21 June 2010; a few papers have been moved to other Symposia subsequent to this date. They can be located by searching for title or first author.

Author of papers inadvertently omitted from this DVD can apply to the editor ([bob.gilkes@uwa.edu.au](mailto:bob.gilkes@uwa.edu.au) and [natta@cyllene.uwa.edu.au](mailto:natta@cyllene.uwa.edu.au)) before 1 October 2010 for the paper to be included in the online Proceedings (<http://www.iuss.org>).

The editors gratefully acknowledge the contributions of many reviewers and editorial staff. We are particularly indebted to Kate Vella, Worachart Wisawapipat and Tingtong Darunsontaya.

Professor Robert Gilkes  
Dr. Nattaporn Prakongkep  
Editors Proceedings 19<sup>th</sup> WCSS  
University of Western Australia  
Crawley, WA, Australia  
1<sup>st</sup> July 2010

# Development of Halloysite/Smectite Mixed Layer Mineral in Paleudult of Java Island

Mohammad Nurcholis

Department of Soil Science, Univ. Pembangunan Nasional "Veteran" Yogyakarta  
Jl. Lingkar Utara Condongcatur Yogyakarta Indonesia, 55283  
Personal communication: M Nurcholis [nurch2003@yahoo.com](mailto:nurch2003@yahoo.com)

## Abstract

This study was initiated to investigate the development of unique kaolin/smectite mixed layer minerals in the Paleudult of Java Island. The present study was aimed to identify the kaolin minerals and to understand the development of the kaolin/smectite mixed layer. Clay fraction samples were collected from the Paleudult profile. Selective dissolutions using dithionite citrate bicarbonate, NaOH, and Na-citrate treatments were conducted on the clay fraction. The result showed that free iron oxide content increased with the depth. Halloysite which showed a 7.6 Å peak on the X-ray diffraction (XRD) was confirmed by the hot NaOH extraction. SiO<sub>2</sub>/Al<sub>2</sub>O<sub>3</sub> molar ratio from the hot-citrate extracts provided the halloysitic property only in the upper horizons. However, the peaks of 10-14 Å on the XRD collapse at 300°C following the hot citrate treatment suggesting that crystallinity of the halloysite mineral is weak. This halloysite mineral developed well in the upper horizons. The development of halloysite/smectite mixed layer is proposed in the four stages.

## Key Words

Halloysite, development of halloysite/smectite mixed layer.

## Introduction

Smectite minerals are found in several acidic soils such as Red-yellow Podzolic Soils in Java Island. The formation of smectite minerals in this area might be resulted from ejected materials from activities of several volcanoes in the past. These materials were then deposited in the sea and further weathering processes resulting smectite minerals. Geological processes might lead an uplift of the sea floor to be terrestrial and then developed soils. McDaniel (1995) reported the presence of beidellite in the E horizon in Spodosols that developed on volcanic ash in Northern Idaho.

Hydroxyl-Al interlayered vermiculite mineral was found in some Ultisols in Indonesia (Goenadi and Tan 1988; Supriyo *et al.* 1992). They described that this minerals showed 14.02 Å peak on the XRD on the Mg<sup>2+</sup>-saturated clay and glycerol solvation, and the peak was gradually collapsed for K<sup>+</sup>-saturated clay and following subsequent heating treatments. Rich (1960) mentioned the interlayering process of aluminum in the interlayer space of expansive 2:1 minerals. Transition of smectite to chlorite that resulted from geothermal alteration was reported by Robinson and de Zamora (1999). Kaolin/smectite mixed layer was identified in the clay fraction of the Paleudult developed on dacitic-andesitic tuffs from Jasinga Subdistrict Bogor Regency West Java Province Indonesia (Nurcholis and Tokashiki 1998).

The objectives of the study were: (i) to identify the kaolin minerals in the kaolin/smectite mixed layer minerals, and (ii) to understand the development of this kaolin/smectite mixed layer minerals in the soil profile of the Paleudult.

## Materials and Methods

The studied soils were the Paleudult, developed from volcanic materials, at Jasinga subdistrict, Bogor regency, West Java province. Five soil samples were collected from each horizon that developed in the soil profile. Soil texture was analyzed using mechanical analysis after oxidizing the organic matter using H<sub>2</sub>O<sub>2</sub>. The coarse and fine sand, silt and clay fractions were separated and collected for mineralogical studies. The fine sand fraction was then analyzed the primary mineral composition using XRD. The clay fraction was then pretreated using dithionite citrate bicarbonate (DCB) using Mehra and Jackson (1960) method. Formamide treatment was conducted to identify the kinds of kaolin minerals. Successive dissolutions were done to identify the properties and kinds of mineral that compose the clay specimens. Hot NaOH treatment was done on the DCB pretreated clay to extract kaolin mineral (Tokashiki *et al.* 1986). The extractable Si and Al elements were then determined using an inductively couple plasma spectrophotometer (ICPS). Sodium citrate treatment (Tamura 1958) was conducted on the residual clay after NaOH-DCB treatment. In

the same way, the extractable Si and Al were then determined using ICPS. The residual clays were then analysed mineralogical properties using XRD.

## Results

### *Lithologic discontinuity*

Table 1 shows the primary minerals identified using XRD for fine sand and silt fractions. Integrated intensity of feldspar and amphibole increased with the depth, except feldspar decreased in B3 horizon, on the contrary quartz decreased with the depth. Based on the Goldich stability series, amphibole is more stable than feldspar. Feldspar was only detected in the A horizon. Feldspar was still detected in the 2B horizon. Anatase was detected in the silt fraction in all horizons and it showed increase with the depth. Free iron oxides slightly increased from the surface horizon to B3 horizon, and slightly increased in 2B horizon. According to the results it can be stated that the soil profile exhibited lithologic discontinuities, as there was a different in the mineralogical composition in the fine sand fraction.


**Table 1. The primary minerals identified using XRD of the fine sand and silt fractions**

Horizon	Fine sand			Silt		
	Feldspar	Amphibole	Quartz	Feldspar	Quartz	Antase
	Integrated intensity fraction*					
A	0.07		0.93	0.07	0.89	0.10
B1	0.12	0.05	0.83		0.89	0.11
B2	0.23	0.05	0.72		0.86	0.14
B3	0.07	0.19	0.72		0.86	0.14
2B			1.00	0.12	0.71	0.17

\*relative value among the detectable minerals

### *Halloysite/smectite mixed layers*

Figure 1 shows the XRD pattern of the clay fraction of the soil A horizon as a representative result for five clay samples. The Mg saturated clay showed 15.9 Å and 7.6 Å peaks. NaOH treatment resulted in a collapse of the 7.6 Å peak. Glycerol solvation resulted in swelling the peak of 15.9 Å to 19.64 Å, whereas the 7.6 Å peak changed from a relatively sharp peak to a broad peak. K saturated clay showed a change in the 15.9 Å peak of the Mg saturated clay to a broadening of the 12.1 Å peak accompanied with a continuous form between 7.6 Å and 12.1 Å. Formamide solvation resulted in a shift of the 7.6 Å peak to 9.72 Å. Accordingly, it is suggested that the 7.6 Å peak is halloysite.



**Figure 1. The XRD pattern of the clay fraction of the soil at A horizon**

### *Development of the halloysite/smectite mixed layers*

Table 2 shows the results of the successive dissolution using hot NaOH and hot citrate extraction. The hot-NaOH extraction resulted in the SiO<sub>2</sub>/Al<sub>2</sub>O<sub>3</sub> molar ratio of 2.29 for the A horizon and increased from A to B2 horizon and then decreased for the following B3 and 2B horizons. According to Tan and Troth (1982), the extracted material is kaolin minerals, and for the present study is halloysite. Prolonged hot-citrate extraction

was successfully used to identify the interlayer materials in hydroxyl-Al interlayered vermiculite (Nurcholis 2003). The hot citrate extractable  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$  resulted in the  $\text{SiO}_2/\text{Al}_2\text{O}_3$  molar ratio of 2.83 for the A horizon, and the highest value was for the B2 horizon. The value of the  $\text{SiO}_2/\text{Al}_2\text{O}_3$  molar ratio resulting from the hot citrate extraction is too high for halloysite. It is probably that prolonged hot citrate extraction also caused the destruction of the crystals of whole minerals. The lithologic discontinuity in the studied soil profile might also contribute to the variation in materials.

**Table 2. Dissolution of silica and alumina and silica/alumina molar ratio by the NaOH treatment for the DCB pretreated clays**

Horizon	NaOH extraction on the DCB pretreated clay			Hot-citrate extraction on the NaOH-DCB pretreated clay		
	$\text{Al}_2\text{O}_3$ mg/g	$\text{SiO}_2$	$\text{SiO}_2/\text{Al}_2\text{O}_3$ molar ratio	$\text{Al}_2\text{O}_3$ mg/g	$\text{SiO}_2$	$\text{SiO}_2/\text{Al}_2\text{O}_3$ molar ratio
A	58.75	79.37	2.29	1.93	3.22	2.83
B1	67.71	90.91	2.28	1.49	2.08	2.36
B2	62.62	96.09	2.60	1.81	3.70	3.46
B3	67.17	96.09	2.43	1.54	2.95	3.25
2B	61.06	72.71	2.01	1.62	2.98	3.31

## Conclusion

According to the results, it is possible to propose a mechanism of the development of halloysite/smectite mixed layer in four stages, as follows: the first stage is designated with the deposition of quaternary volcanic ash under the sea environment and it was diagenetically altered to smectite minerals. The second stage is gradual transgression of the sea, weathering and soil genesis processes. The third stage is Al-OH growth in the interlayer space of smectite mineral as hydroxyl-interlayered smectite. And the fourth stage is an inversion of a portion of the Si tetrahedral sheets of smectite mineral and halloysite which are then separated as free forms, and there is an arrangement of halloysite-smectite layers.

## References

- Goenadi DH, Tan KH (1988) Differences in clay mineralogy and oxidic ratio of selected LAC soils in temperate and tropical regions. *Soil science* **146**, 151-159.
- McDaniel PA, Falen AL, Tice KR (1995) Beridelite in E horizon of Northern Idaho Spodosols formed in volcanic ash. *Clays and Clay Minerals* **43**, 525-532.
- Mehra OP, Jackson ML (1960) Iron oxide removal from soils and clays by a dithionite citrate system buffered with sodium bicarbonate. *Clays and clay Minerals* **7**, 317-327.
- Nurcholis M, Tokashiki Y (1998) Characterization of kaolin/smectite mixed layer mineral in Paleudult of Java Island. *Clay Science* **10**, 291-302.
- Nurcholis M (2003) Identification of interlayer materials in the Hydroxy Interlayered Vermiculite. In 'Proceedings International conference on Mineral and energy Resources Management'. (Eds Sutarto) pp. 127—134. (UPNVY Press)
- Rich CI (1960) Aluminum in interlayer of vermiculite. *Soil Science Society of America Proceeding* **24**, 26-32.
- Robinson D, de Zamora AS (1999) The smectite to chlorite transition in the Chipilapa geothermal system, El Salvador. *American Mineralogist* **84**, 607-619.
- Supriyo H, Matsue N, Yoshinaga N (1992) Chemistry and mineralogy of some soils from Indonesia. *Soil Science and Plant Nutrition* **38**, 217-225.
- Tamura T (1958) Identification of clay minerals from acid soils. *Journal of Soil Science* **9**, 141-147.
- Tan KH, Troth PS (1982) Silica-sesquioxides ratios as aids in characterization of some temperate and tropical region. *Soil Science society of America Journal* **46**, 1109-1114.
- Tokashiki Y, Dixon JB, Golden DC (1986) Manganese oxide analysis in soils by combined X-ray diffraction and selective dissolution methods. *Soil Science society of America Journal* **50**, 1079-1084.