

Analysis on the Capacity..

by Sari B Kusumayudha

Submission date: 23-Jan-2018 09:39AM (UTC+0700)

Submission ID: 905604343

File name: 4._JEEG_Capacity_Building_Fullpaper.pdf (1.32M)

Word count: 4211

Character count: 23919

2

International Journal of Economic and Environment Geology

Search:

[Home](#)[Current Issue](#)[Past Issue](#)[About Journal](#)[Call For Paper](#)[Contact Us](#)

2

Title Page of the Journal Illustrated



Current Issue
Volume-3(2) July-Dec, 2012

[CALL FOR PAPER](#)

[CALL FOR PAPER](#)

- [About Journal](#)
- [Publication Information](#)
- [Instructions to Author](#)
- [Manuscript Submission](#)
- [Editorial Board](#)
- [Contact Us](#)



Viqar Husain, Department of Geology, University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Copyright © 2002 - 2009 www.econ-environ-geol.org

Designed and Developed by [Aspire Solutions](#)

2

International Journal of Economic and Environment Geology

Search:

[Home](#)[Current Issue](#)[Past Issue](#)[About Journal](#)[Call For Paper](#)[Contact Us](#)

Current Issue Volume-3(1) July-Dec, 2012

Table of Content

[Estimation of Material loss due to Chemical Weathering of Some Rocks in Siri Lanka.](#)

U.de S. JAYAWARDENA

[Analysis on the Capacity Building Efforts for Mitigating Volcanic Risks during 2010 Eruption of Mount Merapi, Central Java, Indonesia.](#)

SARI BAHAGIARTI KUSUMAYUDHA

[Assessing the Effect of Cement Dust Emission on the Physicochemical Nature of Soil around Messebo area, Tigray, North Ethiopia.](#)

SAMUEL ESTIFANOS and AYNALEM DEGEFA

[Alteration and Litho geochemistry of Alteration Rocks at Well Kmj-49 Kamojang Geothermal Field West Java Indonesia.](#)

D.F.YUDIANTORO, EMMY SUPARKA, ISAO TAKASHIMA, DAIZO ISHIYAMA YUSTIN KAMA

[Coal Mine Accidents in Bangladesh: Its Causes and Remedial Measures.](#)

MD. MINHAJ UDDIN MONIR AND H. M. ZAKIR HOSSAIN



2

Viqar Husain, Department of Geology, University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Copyright © 2002 - 2009 www.econ-environ-geol.org

Designed and Developed by [Aspire Solutions](#)

2

International Journal of Economic and Environment Geology

Search:

[Home](#)[Current Issue](#)[Past Issue](#)[About Journal](#)[Call For Paper](#)[Contact Us](#)

About Journal

International Journal of Economic and Environment Geology (IJEEG) is an online, open access and peer reviewed international journal dedicated to the latest advancements in the fields of Economic and Environmental Geology. It is half yearly journal, published by Society of Economic Geologists and Mineral Technologists (SEGMITE) and supported by Pakistan Science Foundation (PSF) since 2010. This journal is abstracted in Pakistan Scientific and Technological Information Center (PASTIC) which is the premier organization in the field of scientific and technological information dissemination worldwide.

International Journal of Economic and Environment Geology (ISSN: 2223-957X) is widely circulated. The aim of this journal is to provide a platform for scientists, academicians and technologists all over the world to publish their research work related to the scope of IJEEG. This journal publishes original and review articles (full length paper/short communication). Topics include but not restricted to: Exploration of metallic and non metallic mineral deposits including gemstones, coal and petroleum; Mining geology, mineral processing and industrial applications; Surface and groundwater resources, water pollution, impact on health, agriculture, environment and remedial measures; Geohazards and mitigation options.

Manuscript Submission: The manuscript (s) prepared as per instructions to Author (www.econ-environ-geol.org), should be submitted to the Editor-in-Chief on prof.viqarhusain@yahoo.com. Manuscript submitted for publication will immediately be sent for editorial scrutiny by the journal's international Editorial Board.



2

Viqar Husain, Department of Geology, University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Copyright © 2002 - 2009 www.econ-environ-geol.org

Designed and Developed by [Aspire Solutions](#)

2

International Journal of Economic and Environment Geology

Search:

[Home](#)[Current Issue](#)[Past Issue](#)[About Journal](#)[Call For Paper](#)[Contact Us](#)

2
International Journal of Economic and Environment Geology
(Int.j.econ.env.geol) ISSN: 2223-957X
www.econ-environ-geol.org

Editor-in-Chief

Viqar Husain, Department of Geology (Convener SEGMITE & AGID Vice President, Asia)
University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com , segmite@gmail.com)

2
Associate Editors

Associate Editors Hossain Md Anwar, School of Earth and Environment, University of Western
Australia, Crawley WA 6009, Australia.

Prosun Bhattacharya, Royal Institute of Technology (KTH), Stockholm, Sweden.

Alan E. Fryar, Department of Earth and Environmental Sciences, University of Kentucky, USA.

Jochen Bundschuh, University of Applied Sciences, Institute of Applied Research, Karlsruhe,
Germany & Royal Institute of Technology (KTH), Stockholm, Sweden.

Jochen Bundschuh, University of Applied Sciences, Institute of Applied Research, Karlsruhe,
Germany & Royal Institute of Technology (KTH), Stockholm, Sweden

Managing Editor

Adnan Khan, Department of Geology, University of Karachi, Pakistan.

Editorial Board

George P.C. (USA)	David K. Kremer (USA)	S. Sarfaraz Solangi (Pakistan)
A. J. Reedman (UK)	2 del M. Aly Hassan (Egypt)	Ali Al-Mishwat (Kuwait)
Rehanul Haq Siddiqi (Pakistan)	Akhtar A. Saleemi (Pakistan)	Pramendra Dev (India)
Shahida Begum (Pakistan)	S. D. Limaye (India)	Tahseenullah Khan (Pakistan)
M. Tahir Shah (Pakistan)	C. P. Rajendran (India)	M. Nehal uddin (Bangladesh)
Michael Katz (Australia)	Corrad, D. Lindholm (Norway)	M. A. Obaid (Egypt)
Hongbin Zhan (USA)	2 Tayel El-Hasan (Jordan)	Werner Balderer (Sweden)
A. R. Tabrez (Pakistan)	Ashok Kumar Singhvi (India)	Ghulam Murtaza Arain (Pakistan)
Madhumita Das (India)	Ashish Sarkar (India)	Rubina Bilqees (Pakistan)
U. De S. Jayawardena (Sri Lanka)	P. K. Srivastava (India)	Abdul Salam (Pakistan)
Mihir Deb (India)	O. Okunlola (Nigeria)	2 Am Ali Khawaja (Pakistan)
S. Weerawarnakula (Sri Lanka)	Vaclav Nemeč (Czech Republic)	James T. Teller (Canada)
M. Mansoor Khan (Pakistan)	Silvia Peppoloni (Italy)	Abida Farooqi (Pakistan)
F. M. Zafar Kaifi (Pakistan)	Fares Howari (UAE)	Sari B. Kusumayudha (Indonesia)

Asif Inam (Pakistan)

Sadaf Naseem (Pakistan)

M.Y. Khuhawar (Pakistan)

- [About Journal](#)
- [Publication Information](#)
- [Instructions to Author](#)
- [Manuscript Submission](#)
- [Editorial Board](#)
- [Contact Us](#)



Viqar Husain, Department of Geology, University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Copyright © 2002 - 2009 www.econ-environ-geol.org

Designed and Developed by [Aspire Solutions](#)

International Journal of Economic and Environment Geology

Search:

[Home](#)[Current Issue](#)[Past Issue](#)[About Journal](#)[Call For Paper](#)[Contact Us](#)

Chief Editor

Viqar Husain, Department of Geology (Convener SEGMITE & AGID Vice President, Asia)
University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Associate Editors

Associate Editors Hossain Md Anwar, School of Earth and Environment, University of Western Australia, Crawley WA 6009, Australia.

Prosun Bhattacharya, Royal Institute of Technology (KTH), Stockholm, Sweden.

Alan E. Fryar, Department of Earth and Environmental Sciences, University of Kentucky, USA.

Jochen Bundschuh, University of Applied Sciences, Institute of Applied Research, Karlsruhe, Germany & Royal Institute of Technology (KTH), Stockholm, Sweden.

Managing Editor

Adnan Khan, Department of Geology, University of Karachi, Pakistan.



2

Viqar Husain, Department of Geology, University of Karachi, Pakistan
(prof.viqarhusain@yahoo.com, segmite1989@yahoo.com)

Copyright © 2002 - 2009 www.econ-environ-geol.org

Designed and Developed by [Aspire Solutions](#)

1

Analysis on the Capacity Building Efforts for Mitigating Volcanic Risks during 2010 Eruption of Mount Merapi, Central Java, Indonesia

SARI BAHAGIARTI KUSUMAYUDHA*

* UPN "Veteran", Jl SWK 104, Condongcatur, Yogyakarta 55283, Indonesia

**STTNas, Jl Babarsari, Yogyakarta 55281, Indonesia

E-mail: saribk@gmail.com

Abstract

Mount Merapi that is one of the most active volcanoes on the world erupted again during October to November 2010. Its climax activities happened on 5th November at 00.10 pm, with different type of eruption from Mount Merapi of last 50 years. Ordinary, Mount Merapi activity starts from lava dome development, followed by dome collapse to create pyroclastic flow. This specific character of eruptions called Merapi Type. This type of eruption did not happen in the year 2010, but Vulcanian and Pelean types. The pyroclastic flows at that time killed 341 people and buried many villages on the southeastern slope, while the secondary hazard of lahar destroyed many other human settlements and infrastructures on the western slope of the volcano.

Actually, capacity building program in the areas of around Mount Merapi has been done and established since more than 15 years ago. In most villages, there are community associations that well trained on volcanic hazard mitigation and early warning system. The association name is Association of Mountains Belt of Merapi. Map of Mount Merapi hazards was also already set by the Center of Volcanology and Geologic Disaster Mitigation. Unfortunately, human are not able to order the nature. The character of Mount Merapi eruption in the year 2010 was inconsistent. There was much higher gas pressure, much longer distant of pyroclastic flow, and much greater volume of volcanic material poured from the crater. This made people and stake holders very astonished in handling the evacuation. However, a socio-cultural factor in this respect is that the local people and agriculturists view Mount Merapi as a God which gives them fertile soil and water for agriculture and are reluctant to move away even under an impending threat of a volcanic hazard. This mind-set of people is a challenge in capacity building as the people prefer in-situ protective measures rather than moving away.

Introduction

The study was done in the area of around Mount Merapi, while the volcano is located within densely populated territory of Central Java province and Yogyakarta Special Region (Figure 1). As it is known, Mount Merapi belongs to the most active volcano of Indonesia, and one of the most attractive volcanoes for volcanologists on the world. Once in every about 4 years its activity increases, threatening surrounding environment with its phenomenal pyroclastic flows, glowing clouds, and lahar. In the year 2006, Mount Merapi showed its force, and erupted on 14 June, burying a tourist object at Kaliadem of Cangkringan District, Sleman Regency. In October to November 2010 Mount Merapi was active again with different characteristic of eruption, killing many people and wiping out numerous villages of Cangkringan District, Kemalang District, and Magelang Regency. Although the volcano is some time dangerous, but it brings blessings to the surrounding area by supplying sands, stones, fertile land, and beautiful scenery.

This paper reports and analyzes the characteristics of Mount Merapi eruption 2010, its impact to surrounding, and the capacity building of local communities in term of volcanic hazards mitigation. Methods applied in this study were field surveying, mapping, and primary or secondary data analysis.




Fig. 1. Map showing location of the study area

Mount Merapi the Active Volcano

Mount Merapi is classified into a very active and very young volcano. Camus et al (2000) infer the earliest growth of the volcano began at least 40,000 years B.P. (Newhall et al, 2000). Historical records verify that the character of

1 Mount Merapi eruption is dynamic and changed by time. Variation of effusive and explosive types have occurred, shown by diverse volcanic products such as thick lavas, widespread volcanic sand, ash and dust deposits, pyroclastic breccias, and auto breccias (Kusumayudha 2001, 2008).

Mount Merapi stands on the intersection of two volcanic lineaments, i.e. Ungaran - Telomoyo - Merbabu - Merapi and Lawu - Merapi - Sumbing - Sundoro - Slamet (Figure 2). Mount Merapi also lays on the meeting point of Semarang fault and Solo fault (Figure 3). The volcano exists due to the subduction of Indo-Australia oceanic plate to Eurasia continental plate. Partial melting beneath 75 to 125 km depth brings about calc-alkaline magma, as the source of Merapi products. Based on geophysical investigation, it is identified that the magma chamber of Merapi is relatively shallow, i.e. about 3 km below the ground surface (Sri Brotopuspito, 2006), the magma is very viscous, and its gas pressure is relatively low. This condition makes Merapi activity just displaying weak explosion, lava dome development, and *muee ardentes d'avalanche* (Kusumadinata, 1979).



1 Fig 2. Volcanic lineaments of Ungaran-Telomoyo-Merbabu- Marapi and Lawu-Merapi-Sumbing-Sundoro-Slamet on the physiographic map of Central -East Java (Van Bemmelen, 1949)





Fig 3. Tectonic sketch map of Java (After Martodjojo, 1994 vide Riswandi & Purwanto, 2008)

3

According to Camus et al (2000), there are four periods of Merapi volcanic activities, comprising Ancient Merapi period, Middle Merapi period, Recent Merapi period, and Modern Merapi period. Ancient Merapi period is characterized by thick olivine andesite lava. The Middle Merapi period produced thick andesitic lava flows and *mee ardentees* deposits. The products of Recent Merapi period are thin lava flows, pyroclastics and epiclastic deposits. Meanwhile Modern Merapi period is specified by the Merapi type of eruptions, i.e. a continuous growth of summit dome, followed by collapses and phases of quiescence (Camus et al, 2000). The figure 4 below is the geologic map of Mount Merapi after 2010 eruption (after Paripurno, 2006)



1 Fig. 4. The geologic map of Mount Merapi (Revised after Paripurno, 2006)

The 2010 Eruption

Mount Merapi eruption of 2010 was not similar to that happened of last 50 years. The activity which initially starting from lava dome development on the summit was not occurred this period. This time movement of the magma was too fast. According to the Institute of Research and Development of Volcanic Technology (IRDVT), the energy of Mount Merapi eruption of 2010 was many times higher than that of 2006 eruption. In the year 2010, since the activity level was pointed to be updated from normal active level to be first alert level, second alert level, and then third alert level, there was no glowing lava flows or dome detected on the summit. But suddenly Mount Merapi erupted and exploded on 26 October 2010.

1 When Mount Merapi erupted in 2010, everyone was not well prepared yet. The eruption was more explosive, higher gas pressure, and tremendous *mee ardentees* (Figure 5). The climax eruption occurred on 5 November early morning. According to the National Board of Disaster Prevention BNPB (2010) (<http://www.bnpb.go.id/irw/berita>).

asp?id=247), and Kompas (2012), 341 people died, 366 injured, 50,272 evacuated, and 3,307 buildings destroyed (Table 1). The cost of material lost was estimated approximately 4,230 billion rupiahs, including agricultures.

Mount Merapi eruption of 2010 can be classified into Vulcanian type with more than 5 km high rise dark smokes. Besides upward explosion, the eruptions were also combined with *nuee ardentes d'avalanche* as ever happened to Mount St Helens, known as Pelean type. The greatest pyroclastic flow on 5 November reached 15 km distant from the eruption center to southeast direction through River Gendol valley (Figure 6). This eruption swept and buried many villages, in Cangkringan district, Sleman regency, and Kemalang district of Klaten regency. Even Temple Borobudur, the biggest Buddhist holy place that situated about 35 km from Mount Merapi was covered by volcanic ash (Figure 7). Figure 8 show the situation of post Merapi eruption at some villages.

After the eruption was over, the secondary threats of lahar then appeared. There were 100 million m³ volcanic materials deposited on the upper slopes of the volcano (Kompas, 2012). In the valley of River Gendol, the thickness of pyroclastic deposit reaches 10 m. When the deposits become denser by rain water and flow down lead by gravity, lahar will occur. Since Mount Merapi activity started, some villages along River Code in Yogyakarta, River Putih, and River Pabelan in Magelang Regency were struck by lahar (Figure 9). Impacts of Merapi activity in 2010 to the environment are written in Table 1.

Table 1. Impacts of Merapi Eruption in 2010 (<http://www.bnpb.go.id/irw/berita.asp?id=247>)

Name of Regency	Province	Number of Victims			Volcanic Process as the Destroyer
		Died	Injured	Evacuated	
Sleman	Yogyakarta	243	203	26,774	Pyroclastic flows
Klaten	Central Java	36	30	4,321	Pyroclastic flows
Magelang	Central Java	52	96	18,505	Lahar
Boyolali	Central Java	10	37	672	Volcanic ash
Total		341	336	50,272	Volcanic ash



Fig. 5. Mount Merapi eruption, 10 November 2010



Fig. 6. Pyroclastic flow deposits of Merapi eruption on 5 November 2010, at River Gendol.



Fig. 7. Borobudur temple that located about 35 km western of Merapi was coated by ash



1

Fig. 8. Situation at Kalitengah village, and ruins at Kinahrejo village, after 2010 eruption



Fig. 9. Lahar of River Putih that has broken Yogyakarta – Magelang transportation line

1

Community Based of Disaster Management

1

Disaster management in general has been well understood and run by most Indonesian people (Government, NGO, academician, and community), especially who live in the vulnerable area of volcanic hazards. Theoretically, disaster management can be operated by steps of activities, i.e. pre disaster, during disaster, and post disaster.

In Yogyakarta Special Region, especially for areas which are frequently impacted by Mount Merapi eruption, the government set procedures of disaster management activity steps as the following (Paripurno et al, 2011):

1. **Research:** It is done to study the trends and characteristics of Mount Merapi activities, especially in the area frequently affected by Merapi eruption. Culture of the local people including local wisdoms is important to be studied and understood.
2. **Vulnerability Analysis and Risk Assessment:** There are several threats of Merapi activity, i.e. *ne ardentis* (glowing cloud), volcanic ash and dust, and lahar. Matrix of these threat variants and their risks should be assessed. How will be the risk for one threat variant, combination of two variant or several variant.
3. **Socialization and Community Preparedness:** Local people or community member are taught on natural phenomenon and actions for anticipation. This program is done by local and province Government in collaboration with Non Government Organizations (NGO) and universities in Yogyakarta.
4. **Mitigation:** It is the preparedness to facing disaster or alert situation. Preparation to facing Mount Merapi hazards includes providing evacuation lines, disaster centers, evacuation barracks, and logistic. On the other hand, monitoring of Merapi activities will be done by IRDVT, and the results will always be communicated and informed to the local Government, local community leaders, and the NGO.
5. **Warning System:** When Mount Merapi situation is in the second alert level, the possibility of big eruption should be socialized as soon as possible, not only by persuasive way, but also powered effort. Early warning can be disseminated by lectures in schools or meetings, sermon at churches or mosques, siren, or Short Message System (SMS).

6. **Rescue:** When glowing cloud occurs, the safest method to avoid the risk is escape from the vulnerable area to places or evacuation barracks with suitable logistic.
7. **Communication:** Communication is important in order to detect the threats as early as possible from Yogyakarta as the capital city of the province, and Jakarta as the capital city of the country. It can be done by using satellite telephone system.
8. **Emergency Handling:** When there is someone injured, need to be medically treated, or even missing, the preparedness of the SAR (Save and Rescue) team must be well organized and coordinated.
9. **Sustainability Management:** If Mount Merapi activity does not subside in a short time, the mitigation will need to be proceeded continuously. Well coordination and collaboration must be set up involving local Government, province Government, central Government, and all stakeholders.
10. **Restoration:** This activity belongs to post disaster step, the process may consume such a long time. Renovation planning need to be done carefully because the cost may be high. It can be supported by either local, regional, National, even international communities.
11. **Training and Education:** To achieve best results of disaster management, there should be some skilled and trained officers in every vulnerable area. Then they will be able to train other officers, community members, and NGO members.
12. **Simulation:** After the volunteers are prepared, every vulnerable area should hold simulation on disaster management process in order all the community members and their family able to anticipate and save their selves from disaster threats.

Mitigation and Capacity building

Mitigation action as a part of disaster management was actually already done when Mount Merapi active in 2010. In that time, based on the monitoring result of IRDVT Yogyakarta, the activity status had been updated to be third alert on 24 October morning, at 06.00 am local time. By this status, all people were reminded that Mount Merapi possibly erupts any time. Some works had been done including seismicity analysis, geochemical testing on gas and rock samples, and ground deformation observation using EDM (*Electronic Distance Measurement*). The monitoring was accomplished with visual observation to identify lava dome development, lava dome collapse, and solfatar emission. This showed that such comprehensive approaches had been well done in order to affirm the activity status of the volcano.

On behalf of the change of the volcanic activity status on 24 October 2010, people were suggested with self awareness to leave their places quickly. In that time, old people, women, and children were the first priority to be evacuated. But unfortunately, next day, some people that had been evacuated went back to their village for daily activities.

They were back to their places, because after they stay in the barrack, in fact Mount Merapi did not erupt. On the other hand, they think that they already had the experiences on facing Merapi eruption. They believed that Mount Merapi would take a long enough time for preparing its eruption. There were no natural signals yet, such as flying eagles around the sky, running wild animals (tiger, monkey) down the mountain, and thundering sound from

Merapi “stomach”. Furthermore, at that time the volcano keeper, Mr Maridjan stated that he did not yet get any signs and feelings that Merapi will erupt.

Capacity building actions has actually been already set up in various levels. In the government level, it was done by planning disaster risk reduction (DRR) program, the program is then advocated to the lower level, in order to be operating. The community members establish their DRR action plan for example by forming a team of Preparedness and other activities in their villages. Government and local communities are linked through joint activities such as contingency planning, drills, monitoring and evaluation. The program has been running since the last two years, showing positive growth condition (Paripurno et al, 2011).

Merapi eruption 2010 period was a very valuable lesson for all stakeholders. Disaster management have already run, but still there were hundreds people killed. In the future well coordination, well preparation, and well communication among all stakeholders and especially local people must be well done. In order to minimize the number of victims and risk as minimal as possible, local people should be submissive to Government and official instruction when they have to be evacuated. This is

In principle, capacity building is intended to reduce the vulnerabilities in disaster preparedness and mitigation. In contrary, communities will be weak to face the threats with lack of capacity to reduce the risks. Generally, the capacity of disaster risk reduction of civil society members will be able to well develop, through frequent communication, discussion, workshop, trainings, education, and socialization, without discarding the local wisdoms.

Conclusion

1. Mount Merapi erupted during October-November 2010. Its climax eruption happened on 5th November at 00.10 am. The eruption type in this period was Vulcanian and Pelean type, with much higher gas pressure, much longer distant of pyroclastic flow, and much greater volume of material poured from the crater.
2. The pyroclastic flows of 2010 reached 15 km distant from the center of eruption, killing 341 people, injuring 366 and destroying 3,307 buildings. Value of the lost was estimated approximately 4,230 billion rupiahs, including agricultures.
3. Capacity building program in Mount Merapi surrounding areas has been done and established since more than 15 years ago. In most villages, there are community associations, such as *Association of Mountains Belt of Merapi* that well trained on volcanic hazard mitigation and early warning system. Map of Merapi hazards based on the last data was also already set by the Center of Volcanology and Geologic Disaster Mitigation.
4. During 2010 eruption of Mount Merapi, the mitigation was constrained by a socio-cultural factor, due to people were reluctant to be evacuated. They only temporary leave when Merapi actives, and comeback home when the situation is normal, because for them Merapi is “a God” who gives them fertile soil, water, and everything for life.

References

Andreastuti SD, Alloway BV, Smith IEM (2000) A Detailed Tephrostratigraphic Framework at Merapi Volcano, Central Java, Indonesia: Implications for Eruptive Predictions and Hazard Assessment. *Journal of Volcanology and Geothermal Research*, Vol 100, Elsevier: 51 - 67

- Camus G, Gaurgaud A., Moussand-Berthommier PC, Vincent PM (2000) Merapi (Central Java Indonesia): An Outline the Structural and Magmatological Evolution, with a Special Emphasis to the Major Pyroclastic Events. *Journal of Volcanology and Geothermal Research*, Vol 100, Elsevier: 139 – 163.
- Kompas Daily Newspaper (2012) *Ring of Fire Expedition: Merapi*, Special publication, 18 February 2012
- Kusumadinata K. (1979) *Data Dasar Gunungapi Indonesia* (Basic data on volcanoes of Indonesia). Departemen Pertambangan dan Energi R.I.
- Kusumayudha SB (2001) Hydrogeologic system of Merapi southern slope: The role of volcanic evolution. *Majalah Geologi Indonesia*, Vol 16 No 1, April 2001: 16 - 21
- Kusumayudha SB (2008) Mount Merapi Activity 2006: Its Impact on Groundwater Environment. *Proc International Conference of Women Engineers and Scientists*, Lille, France, 15 – 18 Agustus 2008
- Newhall CG, Bronto S, Alloway B, Banks NG, Bahar I, del Marmol MA, Hadisantoso RD, Holcomb RT, McGeen J, Miksic JN, Rubin M, Sayudi SD, Sukhyar R, Andreastuti S, Tilling RI, Torley R, Trimble D, Wirakusumah AD (2006) 10,000 years of explosive eruptions of Merapi volcano, Central Java: archeological and modern implications. *Journal of Volcanology and Geothermal Research*, Vol 100, Elsevier: 9 - 51
- Paripurno ET (2006) *Geologic Map of Merapi Volcano*. Unpublished
- Paripurno ET, Ronny So, Nakmofa Y, Tanty S, Thamrin, Ferdiwijaya D (2011) *Progress Toward Community Resilience To Disaster Risk*. <http://psmbupn.org/article>
- Riswandi, H. & Purwanto H.S. (2008) Interpretation on the Structure Zone of Alteration Based on Geophysics, Nirmala Area, Bogor, West Java, *Jurnal Ilmiah* Volume 1 No.2 (ISSN 1979-0090), Magister Teknik Geologi, UPN “Veteran” Yogyakarta.
- Sayudi DS, Muzani M, Djalal J (2008) Volcanic activities of Mount Merapi (Evaluation on activity of Januari – April 2008). *Merapi Journal*, Vol 05/01/Edisi April 2008: 1 – 5
- Sri Brotopuspito, K (2006) Merapi Volcano Inspires Scientific Curiosity, *Proc of Volcano International Gathering 2006*, UPN “Veteran” Yogyakarta, 119-126
- Van Bemmelen RW (1949) *The Geology of Indonesia*. Vol I A, Martinus Nijhoff, The Hague, 732 pp
- <http://www.bnppb.go.id/irw/berita.asp?id=247>; www.crisp.nus.edu.sg

Analysis on the Capacity..

ORIGINALITY REPORT

92%

SIMILARITY INDEX

92%

INTERNET SOURCES

9%

PUBLICATIONS

14%

STUDENT PAPERS

PRIMARY SOURCES

1

www.econ-environ-geol.org

Internet Source

68%

2

econ-environ-geol.org

Internet Source

20%

3

ir.lib.hiroshima-u.ac.jp

Internet Source

2%

4

Submitted to Asian Institute of Technology

Student Paper

2%

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On

Analysis on the Capacity..

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

RUBRIC: REVIEW PAPER		0 / 6
PENDAHULUAN (15%)		0 / 6
SANGAT BAGUS (6)	Merujuk pada penelitian sebelumnya	
BAGUS (4)	Tidak ada rujukan tapi esuatu yang baru	
KURANG (1)	Pengulangan dari penelitian yang pernah ada	
PEMBAHASAN (50%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		
KESIMPULAN (20%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		
SARAN (15%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		