

Geomorphologic Model...

by Sari B Kusumayudha

Submission date: 23-Jan-2018 09:39AM (UTC+0700)

Submission ID: 905604355

File name: 7._JGRE_geomorphologic_model.pdf (1.14M)

Word count: 3207

Character count: 19361

From Knowledge to Wisdom

Journal of Geological Resource and Engineering

Volume 3, Number 1, Jan.-Feb. 2015



David Publishing Company
www.davidpublisher.com

ISSN 2328-2193
DOI:10.17265/2328-2193

1

Journal of Geological Resource and Engineering

Volume 3, Number 1, Jan. -Feb. 2015 (Serial Number 6)

David Publishing Company
www.davidpublisher.com

PUBLISHING

Publication Information:

Journal of Geological Resource and Engineering is published bimonthly in hard copy (ISSN 2328-2193) by David Publishing Company located at 240 Nagle Avenue #15C, New York, NY 10034, USA.

Aims and Scope:

Journal of Geological Resource and Engineering, a bimonthly professional academic journal. Its aim is to achieve the advancement and dissemination of information in the fields of geology, geotechnical, geological engineering and so on. And articles interpreting successful policies, programs or cases are also welcome.

Editorial Board Members:

Prof. Barbara, Maria Namysłowska-Wilczyńska (Poland), Dr. Morovvat Faridazad (Iran), Prof. Bahadur Singh Kotlia (Nainital), Dr. Behshad Jodeiri Shokri (Iran), Prof. Mustafa Aytakin (Kingdom of Bahrain), Prof. Vlado Gicev (Macedonia), Prof. Abdelaziz Lafi Khlaifat (Abu Dhabi), Dr. Hossein Masoumi (Australia), Prof. Sylejman Hyseni (Republic of Kosovo), Prof. Mohamed EL WARTITI (Morocco), Prof. José Fernando Thomé Jucá (Brazil)

Manuscripts and correspondence are invited for publication. You can submit your papers via web submission, or E-mail to geology@davidpublishing.com or geology_davidpublishing@hotmail.com. Submission guidelines and web submission system are available at <http://www.davidpublisher.com>.

Editorial Office:

240 Nagle Avenue #15C, New York, NY 10034, USA

Tel: 1-323-984-7526, 323-410-1082, 607-4416220; Fax: 1-323-984-7374, 323-908-0457

E-mail: geology@davidpublishing.com, geology_davidpublishing@hotmail.com

Copyright©2015 by David Publishing Company and individual contributors. All rights reserved. David Publishing Company holds the exclusive copyright of all the contents of this journal. In accordance with the international convention, no part of this journal may be reproduced or transmitted by any media or publishing organs (including various websites) without the written permission of the copyright holder. Otherwise, any conduct would be considered as the violation of the copyright. The contents of this journal are available for any citation. However, all the citations should be clearly indicated with the title of this journal, serial number and the name of the author.

Abstracted / Indexed in:

Chinese Database of CEPS, CNKI, China

Chinese Scientific Journals Database, VIP Corporation, Chongqing, China

ProQuest

CiteFactor (USA), Ulrich's International Periodicals Directory, USA

Google Scholar

Summon Serials Solutions

Pubicon Science Index

SJournal Index

Scientific Indexing Services

Subscription Information:

Price (per year):

Print \$520; Online \$360; Print and Online \$680

David Publishing Company

240 Nagle Avenue #15C, New York, NY 10034, USA

Tel: 1-323-984-7526, 323-410-1082, 607-4416220; Fax: 1-323-984-7374, 323-908-0457

E-mail: order@davidpublishing.com

David Publishing Company

www.davidpublisher.com

PUBLISHING

Journal of Geological Resource and Engineering

Volume 3, Number 1, Jan.-Feb. 2015 (Serial Number 6)

Contents

Theoretical Research

1 **Geomorphologic Model of Gunungsewu Karst, Gunung Kidul Regency, Yogyakarta Special Territory, Indonesia: The Role of Lithologic Variation and Geologic Structure**

Sari Bahagiarti Kusumayudha, Jatmiko Setiawan, Ayu N. Ciptahening and Prabawa Dwi Septianta

8 **Lineament Analysis Using ASTER Satellite Images in the Southern Part of the Korean Peninsula**

Weon-hack Choi, Sung-il Cho, Daiei Inoue, Makoto Yanagida, Dong-hee Park and Chun-joong Chang

17 **The Influence of Soil Characteristics in Seismic Response of Embedded Structures**

Adrian Popovici, Cornel Ilinca and Răzvan Vărvorea

Techniques and Methods

28 **Punch Multi-slice Longwall Mining System for Thick Coal Seam under Weak Geological Conditions**

Takashi Sasaoka, Akihiro Hamanaka, Hideki Shimada, Kikuo Matsui, Nay Zar Lin and Budi Sulistianto

37 **WHR (Waste Heat Technology) Method in Tri-generation Model**

Imrich Discantiny

42 **Gas Ratio Analysis in Hovsan Oil Field**

Samir Hashimov

49 **Calibration of Hydrological Streamflow Modeling Using MODIS**

Manithaphone Mahaxay, Wanchai Arunpraparut, Yongyut Trisurat and Nipon Tangtham

Geomorphologic Model of Gunungsewu Karst, Gunung Kidul Regency, Yogyakarta Special Territory, Indonesia: The Role of Lithologic Variation and Geologic Structure

Sari Bahagiarti Kusumayudha¹, Jatmiko Setiawan¹, Ayu N. Ciptahening¹, Prabawa Dwi Septianta¹

1. *Geology Study Program, Faculty of Mineral Technology, UPN "Veteran" Yogyakarta
Jl. SWK 104 Condongcatur, Yogyakarta. E-mail: saribk@gmail.com; sarik@upnyk.ac.id*

Abstract: Gunungsewu is a karst terrain that shows unique geomorphologic phenomena. The area is mainly composed of limestone of Wonosari Formation. Interaction of tectonic, erosion, denudation, and deposition occurred since Miocene period brings about diversity of landforms that supporting the development of this area as a natural laboratory, specific interest, scientific and educational tourisms. In general the geomorphology of Gunungsewu can be classified into two groups of landforms, the first is positive relief and the second is negative relief. The positive relief includes hills of conical, dome, convex-cone, convex, and ridge. The negative relief involves cave, shaft, doline, uvala, locva, and polje. This diversity is influenced by variation of physical properties of rock composing the landforms, such as hardness, internal friction angle, and geologic structures including thickness, position of bedding plane, and joint pattern.

Key words: geomorphology, karst, rock properties, geologic structures

1. Introduction

Gunungsewu is very famous in Indonesia, especially in Java Island, as a karst terrain that is geologically, geomorphologically, and hydrogeologically shows unique phenomena. Geology of Gunungsewu expresses tracks of tectonic, erosion, denudation, and deposition occurred since Miocene epoch. These processes brought about morphologic diversities that support for the development of this area as natural laboratory, special interest, scientific, and education tourisms. Therefore Gunungsewu is very potential to be promoted as an international level of geopark.

Gunungsewu displays a specific Tropical landform, characterized by the existence of hills, closed depressions, and caves. Although morphology of Gunungsewu area is as an ordinary karst landforms, but there is a diversity. The diversity seems to be reflecting the occurrence of various geologic factors influencing the landform process.

It is interpreted that the morphology is influenced by variation of physical properties of the lithology and the existing geologic structures. This hypothetical view has to be verified. In relation to that, such an assessment is needed to be done. Therefore, this study is aimed to identify geologic factors that role in the formation of karst morphology of the Gunungsewu, and develop a geomorphologic model of this area.

The study area is located in Gunungsewu karst area of Yogyakarta Special Territory, Indonesia, geographically lies of $7^{\circ}57' - 8^{\circ}14'$ south latitude, and $110^{\circ}19' - 110^{\circ}50'$ east longitude, about 25 km southeastern of Yogyakarta city. Case study was especially taken in the Ponjong and Karangmojo districts, Gunung Kidul Regency, on the coordinate of 468750mE - 473750mE and $9117000\text{mN} - 9122000\text{mN}$. Fig. 1 shows the location of the study area that can be reached easily by cars from Yogyakarta city.

* **Corresponding author:** Sari Bahagiarti Kusumayudha, professor, research fields: geology, hydrogeology. E-mail: saribk@upnyk.ac.id



Fig. 1. Map showing the location of the study area

2. Methods of Study

Approach method applied in the study was field surveying, surface geologic and geomorphologic mapping, topographic interpretation, remote sensing, petrologic and geologic structure assessment. Data used for the analyses were derived from field surveys and mapping, and from existing previous studies. Remote sensing method was run by using airphotos of 1 : 35,000 scale and satellite images (Ers and Google Earth).

Survey and mapping were done to identify variation physical characteristics and distribution of the lithology. Geologic structure identification and measurements were operated for strike and dip and also shape of karst morphology such as hill, ridge, and valleys. Topographic and image interpretation were utilized for determining pattern and orientation of fracture system (joint, fractures, fault) also hills and valleys lineaments.

3. Geology

Geologic setting of Gunungsewu belongs to western part of Southern Mountains [18]. The stratigraphy from the oldest to the youngest is composed of Semilir

formation, Nglanggran formation, Sambipitu formation, Oya formation, Wonosari formation, Kepek formation, and locally Terrarosa deposit.

Semilir formation belongs to deep sea sedimentary rocks consists of sandstone, tuff, tuffaceous sandstone, lapilli, tuff breccias, lappilli breccias, agglomerate, andesitic breccias, clay stone, siltstone, and shale of Oligocene – Early Miocene [17]. Semilir formation is overlain by Nglanggeran formation, consists of andesitic breccias, agglomerate, polymixed breccias, lava deposits, sandstone and tuffaceous sandstone. Age of the formation is oligomiocene to Middle Miocene [17]. Nglanggeran formation is conformably overlain by sambipitu formation that comprises marl, claystone, calcareous siltstone, calcareous sandstone, and tuffaceous sandstone. Sambipitu formation is Middle Miocene age [17]. Oya formation conformably overlays Sambipitu formation. This rock unit consists of calcarenite, bedded limestone, calcareous sandstone, and tuffaceous-calcareous sandstone. Age of the formation is Middle Miocene to Mio-Pliocene [10]. Oya formation is overlain by Wonosari formation that consists of reef, massive and bedded limestones of Middle Miocene to Pliocene age [10]. Kepek formation consists of marl and calcarenite that lays over the Wonosari formation of Pliocene – Pleistocene epoch [10]. The geologic map of the study area is shown in Fig. 2.

Gunungsewu has been uplifted and a little bit folded since middle miocene or about 6 million years ago ([17]). There is a syncline expresses as the Wonosari plateau with axial orientation of N75°E - N255°E, dipping no more than 10°, while the hills display homoclinal dipping 5° to 15° relatively southward [9], [10]. In the Ponjong and Karangmojo areas, dip of the bedding planes reaches 18°.

Patterns of fault, joints, and fissures that is mapped by using ERS, generally shows northwest - southeast direction [8], [9]. On the other hand, the general dip direction is to southwest, southeast, and south [10], [16], while the joint patterns shows northwest - southeast and northeast – southwest.




Fig 2. Geologic map of Gunungsewu area [10]

4. Karst Geomorphology

According to White (1988), there are some requirements to form such a perfect karst, this include the existence of limestone of >20 m thickness, massive but jointed, and bedded. There must be valleys that the elevation is lower than the position of the limestone. Karstification also needs warm climate, high enough rainfall (>2000 mm/year); and relatively slow tectonic activity (uplifting) involving all of the area.

Beside the requirements mentioned above, morphologic diversity of karst is influenced by physical characteristics of the limestone, including *strength* and *hardness*. Joints, cracks, faults, bedding planes are factors that contribute in karstification process (Bogli, 1980), into which dissolution occurs.

Area of the Gunungsewu area spreading is about 1500 km². Based on air photo interpretation, there are more or less 45.000 hills [10], with various shapes, such as cone, dome, convex, and ridge (Fig. 3 – Fig. 6). The negative reliefs include doline, uvala, locva, polje, shaft, and cave.



Fig 3. Cone karst



Fig 6. Ridge karst



Fig 4. Convex cone karst



Fig 5. Dome karst

5. Influence of Lithologic Variation and Geologic Structure

Limestone of Wonosari Formation in the Gunungsewu Area is composed of some lithofacieses, such as bedded limestone consist of *packstone* and *wackestone*, and reef limestone or *boundstone* [8],[9],[10]. Beside *wackestone*, *packstone* and *boundstone*, there is also *grainstone*. The limestone physically shows karstification or hard and cavernous, and calichification (*chalky limestone*), weak or brittle. If such limestones is uplifted and exposed in the area with wet tropical climate, karstification will happen. But when the limestone is exposed in an arid with a few rainfall condition, calichification will occur. The sequence of caliche consists of *hardpan*, *platty caliche*, *nodular caliche*, and *chalky caliche*. [15].






In this study, result of geomorphologic and geologic mapping on the distribution and variation of lithology and geologic structure verifies that different physical characteristic of lithology forms different shape of morphology, either positive or negative reliefs. In this case, internal friction angle (ϕ) of limestone has an important role in producing the slope of morphology of the karst. Table 1 contains various values of internal friction angle of limestone from some sources.

The orientation of geologic structures especially joint, fault, and fissures control the lineaments of hills, valleys, that are Northwest – Southeast and Northeast – Southwest. The dip of bedding plane influences morphologic dimension. Table 2 displays orientations of joint patterns and hills or valleys lineaments.

Table 1 Values of internal friction angle (ϕ) of karstic limestone and caliche, quoted from some sources

ϕ Limestone ($^{\circ}$)	ϕ Dolomite, Chalky limestone ($^{\circ}$)	Source
33 - 40	27 - 31	Hoek & Bray (1981)
35 - 50	22	Jumikis (1983)
34,8 - 42	31,5 - 35,5	Goodman (1989)
32 - 49	30 - 41	Franklin & Dussault (1989)

Table 2 Orientations of joint pattern, hills and valleys lineaments

Measured from:	Orientation of the pattern (N...°E)	
	Joint	Valleys lineament
	35 and 315	
Field (175 sample)		
Satellite image	45 and 335	50 and 325
		
Topographic map	45 and 325	45 and 325
		

6. Karst Geomorphologic Model of Gunungsewu

Karst geomorphology modeling of the Gunungsewu area refers to Verstappen (1983), comprises some aspects including morphographic, morphometric, and morphogenesis. Morphogenesis involves passive morphostructure, physical properties of the lithology, active morphostructure including tectonic, and morphodynamic. It is also related to exogenic processes such as erosion, dissolution, and mass wasting.

Geomorphologic diversity of Gunungsewu area involves positive and negative reliefs. Positive relief comprises hills of 10 m to 90 m height. The shape varies from conical (angular top cone), dome, rounded top cone (convex-cone), convex, and ridge. While negative reliefs includes doline, uvala, polje, and shaft, or cave.

Attractive morphology is not only found on the surface, but also in subsurface or in caves. There are various ornaments such as stalactite, stalagmite, pillar, sinter, and flowstones. The formation of endokarst in the caves are also influenced by the existence of fracture structures and bedding planes. The stalactites are always found in longitudinal row following strike of fractures that is crosscutting the wall or roof of the cave.

Table 3 and Table 4 describe the model of karst morphology of Gunungsewu area. As already discussed above, it is controlled by variation of physical properties of the lithology and the existence of geologic structures.

Conclusions

Results of this study and analyses on the geomorphology of Gunungsewu Area conclude that:

1. Karst morphology of Gunungsewu area can be grouped into morphology with positive relief including hills of cone, convex-cone, dome, convex, and ridge shapes, and morphology with negative relief such as cave, shaft, doline, uvala, locva, and polje
2. The variation of karst morphology of Gunungsewu area is influenced by the variation of physical properties of the lithology such as hardness and internal friction angle, and geologic structures such as joint orientation, and existence, position, thickness of bedding plane.
3. Karstic limestone either massive or bedded forms hills of cone, convex-cone, and dome shaped, while caliche limestone displays convex hills. Internal friction angle (ϕ) of the lithology determines the angle of slope inclination of the hills that are formed. The higher the value of friction angle (ϕ), the steeper the slope of hills which are formed.
4. Joint, crack, and fault patterns are correlated to the lineaments of hills and valleys of the Gunungsewu area. Position and thickness of bedding plane also controls the shape of the hills: thin incline bedding plane will result cone shape, thick incline bedding plane will result convex-cone, while gentle to horizontal bedding plane will result dome, convex, and ridge shapes.
5. Cave, shaft, doline, and uvala are only formed in karstic limestone, while polje can be found either in karstic or caliche limestone.

Acknowledgment

Gratefulness is dedicated to the Institute of Research and Community Service, UPN "Veteran" Yogyakarta for its support to this successful study.

References

- [1] Bogli, A., 1980, *Karst Hydrology and Physical Speleology*, Springer-Verlag, 284 p.
- [2] Dunham, R. J., 1962, Classification of Carbonate Rocks According to Depositional Texture. In: W.E. Ham (Ed), *Classification of Carbonate Rocks*. AAPG Mem. 1, p 108 – 121.
- [3] Esteban, M., 1996, *Karst System from Prospect to Reservoir*, Carbonate International Ltd, 168 p.
- [4] Franklin, J.A. & Dussault M.B., 1989, *Rock Engineering*, McGraw Hill Publishing Company, 600 p.
- [5] Goodman, R.E., 1989, *Introduction to Rock Mechanics*, 2nd ed, John Wiley & Sons, 56 p.
- [6] Hoek E & Bray J.W., 1981, *Rock Slope Engineering*, Revised 3rd ed, The Institution of Mining and Metallurgy London, 358 p.
- [7] Jumikis, A.R, 1983, *Rock Mechanics*, nd ed, Gulf Publishing Company, 613 p.
- [8] Kusumayudha, S.B., Zen, M.T., Notosiswoyo S., Gautama R.S., 1997, Study on the distribution of carbonate rocks based on fractal characteristics of the valley patterns and secondary porosity, case: Paliyan and surrounding area, Gunung Kidul, *DIY, Jurnal Teknologi Mineral*, Vol IV, No. 2, 1997, p 71-86.
- [9] Kusumayudha, S.B., M.T. Zen, Notosiswoyo S., Gautama R.S., 1999, hydrogeologic System of Gunungsewu Area, *Proc. IAGI The 28th Annual Convention*, Vol IV, p 73-84
- [10] Kusumayudha, S. B., 2004, *Introduction of Karst Hydrogeology*, Pusat Studi Karst UPN "Veteran" Yogyakarta, 137 p.
- [11] Kusumayudha, S.B., 2005, *Fractal geometry and hydrogeology of Gunungsewu Area*, Adicita Publisher, Yogyakarta, 171 p.
- [12] Kusumayudha S.B., 2008, *Karst: cave, underground river, and spring*, Citra Aji Parama Publisher, Yogyakarta, 44 p.
- [13] Kusumayudha, S. B., 2011, Exploring the Prospective of Gunungsewu Area for a World Geopark, Malaysia – Indonesia Join Geoheritage Convention, 1 – 2 Oktober 2011, Terengganu, Malaysia
- [14] Monroe, W.H., 1976, *The Karst Landforms of Puerto Rico*, United State Government Printing Office, 69 p.
- [15] Moore, C.H., 1989, *Carbonate Diagenesis and Porosity*, Elsevier, 338 p.
- [16] Setiawan, J. & Kusumayudha, S.B., 2011, Groundwater system of Gunung Kendil and Umbul Ponjong as a unique geoheritage, Malaysia – Indonesia Join Geoheritage Convention, 1 – 2 Oktober 2011, Terengganu, Malaysia
- [17] Suyoto, 1994, Stratigraphic sequence of the Gunungsewu carbonates, *Proc. PIT IAGI XXIII*, Vol 1, p 19-32.
- [18] Van Bemmelen, R.W., 1949, *The Geology of Indonesia*, Vol IA, Martinus Nijhoff, The Hague 792 p.
- [19] Verstappen, H. Th., (1983) *Applied Geomorphology*, Elsevier Science Publishers, 437 p.
- [20] White, W.B., 1988, *Geomorphology and Hydrology of Karst Terrains*, Oxford University Press, New York, 464 p.

Table 3 Model of the influence of lithologic physical properties and geologic structures to positive relief karst morphology










LITHOLOGY	MORPHOLOGY			MORPHOGENESIS	
	Morphographic	Morphometric	Active Morphostructure (Geologic Structures)	Passive Morphostructure (Physical Properties)	Morphodynamic (Exogenic Processes)
Dominated by bedded limestone, grainstone, packstone		<ul style="list-style-type: none"> - Steep slope - Slope: 30°-45° - Relatively top sharp - Height: 30 – 90 m 	<ul style="list-style-type: none"> - Joint - Thin bedded: < 1m - Bedding plane inclination: > 5° 	<ul style="list-style-type: none"> hard, cavernous, lapies, $\phi = 32 - 50$ 	Karstification
Dominated by reef limestone, boundstone		<ul style="list-style-type: none"> - Steep slope - Slope: 35°-45° - Top convex - Height: 30 – 70 m 	<ul style="list-style-type: none"> - Joint - Massive, or thick bedded: > 1m - Bedding plane inclination > 5° 	<ul style="list-style-type: none"> hard, cavernous, lapies,, $\phi = 32 - 50$ 	Karstification
Caliche limestone, wackestone		<ul style="list-style-type: none"> - Steep slope - Slope: 30°-45° - Height: 20-50 m 	<ul style="list-style-type: none"> - Joint - Massive or bedded - Relatively horizontal bedding plane: < 5° 	<ul style="list-style-type: none"> hard, cavernous, lapies, $\phi = 32 - 50$ 	Karstification Calichification
Caliche limestone, wackestone		<ul style="list-style-type: none"> - Gentle slope - Slope: 12°-20° - Height: 10 - 30 m 	<ul style="list-style-type: none"> - Joint - With or without bedding plane - Relatively horizontal bedding plane: < 5° 	<ul style="list-style-type: none"> weak, chalky, uncavernous $\phi = 17 - 31$ 	Calichification
Bedded limestone		<ul style="list-style-type: none"> - Moderately inclined to steep slope - Slope: 20°-35° - Top plane - Height: 10 - 30 m 	<ul style="list-style-type: none"> - Joint - Thin bedded: < 1m - Relatively horizontal: < 5° 	<ul style="list-style-type: none"> hard, cavernous, lapies, $\phi = 32 - 50$ Weak, chalky, uncavernous $\phi = 17 - 31$ 	Karstification Calichification

Table 4 Model of the influence of lithologic physical properties and geologic structures to negative relief karst morphology

LITHOLOGY	MORPHOLOGY			MORPHOGENESIS	
	Morphographic	Morphometric	Active Morphostructure (Geologic Structures)	Passive Morphostructure (Physical Properties)	Morphodynamic (Exogenic processes)
Reef limestone, boundstone, Bedded limestone, grainstone, packstone		<ul style="list-style-type: none"> - Various height, reaching 30 m - Various width, reaching 50 m 	<ul style="list-style-type: none"> - Joint - Massive or bedded - Incline to horizontal bedding plane 	hard, cavernous, lapies, $\phi = 32 - 50$	Karstification
Bedded limestone, packstone, wackestone		<ul style="list-style-type: none"> - Diameter of the hole: 10 m - 50 m - Depth: 20 m - 60 m 	<ul style="list-style-type: none"> - Joint - Bedded - Gently incline to horizontal bedded plane : $< 5^\circ$ 	hard, cavernous, lapies, $\phi = 32 - 50$	Karstification
Reef limestone, Bedded limestone, packstone, wackestone		<ul style="list-style-type: none"> - Steep slope - Slope: $30^\circ - 40^\circ$ - Depth: 20 m - 50 m 	<ul style="list-style-type: none"> - Massive of bedded - Incline bedding plane: $> 5^\circ$ 	hard, cavernous, lapies, $\phi = 32 - 50$	Karstification
Bedded limestone Caliche limestone	<p>Dolina (<i>Doline</i>), Uvala</p> 	<ul style="list-style-type: none"> - Moderately inclined to steep slope - Slope: $20^\circ - 30^\circ$ - Base plane - Maximum depth : 10 m - Area: $500 \text{ m}^2 - 1500 \text{ m}^2$ 	<ul style="list-style-type: none"> - Bedded - Relatively horizontal bedding plane: $< 5^\circ$ 	hard, cavernous, lapies, $\phi = 32 - 50$ Weak, chalky, uncavernous $\phi = 17 - 31$	Karstification Calichification
	Polje				

Geomorphologic Model...

ORIGINALITY REPORT

95%

SIMILARITY INDEX

95%

INTERNET SOURCES

2%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

1

eprints.upnyk.ac.id

Internet Source

95%

2

vig2006.recent.or.id

Internet Source

<1%

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On

Geomorphologic Model...

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

RUBRIC: REVIEW PAPER		0 / 6
PENDAHULUAN (15%)		0 / 6
SANGAT BAGUS (6)	Merujuk pada penelitian sebelumnya	
BAGUS (4)	Tidak ada rujukan tapi esuatu yang baru	
KURANG (1)	Pengulangan dari penelitian yang pernah ada	
PEMBAHASAN (50%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		
KESIMPULAN (20%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		
SARAN (15%)		0 / 6
SANGAT BAGUS (6)		
BAGUS (4)		
KURANG (1)		