THE LONGEVITY OF Diaphorina citri ON VARIOUS DOMINANT WEEDS

Siwi Hardiastuti, Mofit Eko Poerwanto*

Agrotechnology Department, University of Pembangunan Nasional "Veteran" Yogyakarta *mofitnuk@yahoo.com, telp/fax: (0274) 487793

ABSTRACT

Citrus vein phloem degeneration (CVPD) is the most impediment disease in citrus production in the world. It mainly vectored by *Diaphorina citri* Kuwayama. Intensive application of insecticides is ineffective and also costly. Various weeds species in citrus orchard are suspected as alternative host of vector. Research on the longevity of *D. citri* on the various weeds is needed for completing the integrated CVPD disease management program. Each of five dominant weeds gained from vegetation analysis were exposed to five days old *D. citri* adults inside the nylon mesh cage, which previously starved for 24 hours. The number of *D. citri* survived was recorded daily (24 h). Analysis of variance and DMRT were done on data. Most of *D. citri* stayed for feeding on *Boerhavia erecta* and survived for 5.91 days. It was significantly different on *Amaranthus spinosus*, and also on *Althermanthera philoxeroides*, *Portulaca oleracea*, *Ludwigia perrenis* which survived for 3.57, 0.20, 0.00, and 0.20 days respectively.

Key words: CVPD, longevity, citrus, Diaphorina citri

INTRODUCTION

Citrus vein phloem degeneration (CVPD) is the most impediment disease in citrus culture in the world and has been attacking citrus in Asia and America (Aubert, 1990; Halbert & Majunath, 2004). Many highly productive plantations where yields were 20 tones per ha decreased sharply into 8.6-15 tones per ha per annum (Irawan *et al.*, 2003). Recently, citrus production in Indonesia has developed rapidly. In 2007, the production increased to 2,625,884 tones. However, CVPD is still a great threat to the citrus production in Indonesia. In 2008, the production decreased again into 2,467,632 tonnes (BPS, 2009).

CVPD mainly vectored by *D. citri* Kuwayama. Fourth and fifth instars as well as adults are able to transmit diseases (Xu et al, 1988; Aubert, 1990). Its role as the insect vector, led to the vector control using insecticides as the principal strategies in CVPD control management. However, significant result of the CVPD disease control has not yet been obtained in the field due to the insufficient understanding of the life of a vector. The existence of various weeds in citrus orchards could be suspected as alternative host of vector (*D. citri*). Some weeds are able as alternative host of pests and pathogens. Kolomento weeds (*Leersia hexandra*) and Sembungan (*Sacciolepis*



239

interrupta) are the host plants of rice stem borer, Cerulang (*Eleusine indica*) is the host plants of *Piricularia oryzae*, and nematodes *Rotylenchus reniformis*. Wuluhan (*Setaria plicata*) and Jajahean or Lempuyangan (*Panicum sp.*) are host plant of rice bug (Rukmana and Saputra, 1999). In order to complete an integrated control program of CVPD, research on *D. citri* longevity in various weeds species should be conducted.

MATERIALS AND METHOD

D. citri mass rearing

Eggs of D. citri was obtained from the field and maintaned on orange jasmine plants. The nymphs wich hatched from eggs reared on orange jasmine shoots, planted in pots (Ø 17 cm, height 12 cm and Ø 25 cm, height 18 cm) and covered with nylon mesh cages (length 100 cm, width 100 cm, height 100 cm). *Weeds identification and vegetation analysis of citrus orchards*

Dominant citrus weeds was determined using analysis vegetation method by taking five ring samples (50 cm x 50 cm) per plots randomly on three citrus orchards. Values of ADe (absolute density), AF (absolute frequency), ADo (absolute dominance), RDe (relative density), RF (relative frequency), and RDo (relative dominance) were calculated and value of SDR (summed dominance ratio) was determined.

D. citri longivity test on weeds

Experiments were carried out by using five dominant weeds planted in pots. Each weed was inserted into nylon mesh cage (\emptyset 10 cm, height 30 cm) and 10 of five days old *D. citri* adults were released in the midle of the cage. They were previously starved for 24 hours before releasing. The cages were covered with sponge (\emptyset 10 cm, 2 cm thick). The number of *D. citri* survived on each weed was observed every 24 hours up to all of them in dead. The treatment was repeated 10 times.

Data analysis

Analysis of variance at 95% significance level was done by using SPSS 10.0.5 (SPSS, 1999) to determine the effect of treatment. Duncan's multiple range test (DMRT) at 95% significance level was carried out to identify significant difference between treatments (Gomez and Gomez, 1983).



RESULTS AND DISCUSSION

Results of vegetation analysis on weeds of citrus orchard found 16 species of weeds which consists of three species of grass, 10 species of broadleaf, and three species of sedges group (Table 1.). Based on these results, five most dominant weed species were taken for testing *D. citri* longevity; there were *Alternathera philoeroides* of grass group, *Oleraceae Portulaca, Amaranthus spinosus, Ludwigia perrenis,* and *Boerhavia erecta* of broadleaf weeds with SDR values of 19.72%, 12.06%, 11.65%, 9.31% and 6.93% respectively. The analysis result also shows that the value of absolute density (ADe) was not necessarily generating high values of SDR, because the value of SDR was determined by three parameters: density (De), dominance (Do), and frequency (F). The highest value of SDR on *A. philoeroides* indicated that this weed was the most dominant weeds in the field.

D, *citri* was able to survive a maximum of eight days with the mean longevity of 5.91 ± 0.251 days on the broadleaf weed *B. erecta*, but was not able to complete one life cycle (Figure 1.). It was significantly different to other weeds, *A. spinosus*, *A. philoeroides*, *L. perrenis* and *P. oleraceae* which were survived for 3.57, 0.20, 0.20 and 0.00 days respectively (Table 2.).

							- 22	
Weeds species	Group	Ade	AF	ADo	RD	RF	RD	SDR(%)
Alternathera philoeroides*	G	15	3	20,5	14,29	12	32,87	19,72
Digitaria longiflora	G	4	1	2,6	3,81	4	4,17	3,99
Physalis angulata	В	1	1	0,6	0,95	4	0,96	1,97
Cyperus iria	S	4	2	1,3	3,81	8	2,08	4,63
Amaranthus spinosus*	В	14	3	6	13,33	12	9,62	11,65
Ageratum conyzoides	В	5	1.	3,7-	4,76	4	5,93	4,90
Ludwigia perrenis*	В	12	3	2,8	11,43	12	4,49	9,31
Eclipta prostrata	В	1	1	0,6	0,95	4	0,96	1,97
Borreria leavis	В	2	1	1,2	1,91	4	1,92	3,27
Eleusine indica	S	5	1	6,1	4,76	4	9,78	6,18
Phylanthus debilis	В	3	2	0,7	2,86	8	1,12	3,99
Boerhavia erecta*	В	6	1	-6,9	5,71	4	11,06	6,93
Portulaca oleraceae*	В	21	2	5,1	20,0	8	8,18	12,06
Euphorbia hirta	В	2	1	0,2	1,91	4	0,32	2,08
Pentella repens	G	8	1	3,9	7,62	4	6,25	5,96
Hedrotis corymbasa	S	2	1	0,1	1,91	4	0,16	2,02
Total		105	25	62,4	99,99	100	99,89	100,6

Table 1. Vegetation analysis on weeds of citrus orchards

Notice : ADe (absolute density), AF (absolute frequency), ADo (absolute dominance), Rde (relative density), RF (relative frequency), and RDo (relative dominance), SDR (summed dominance ratio), * (five most dominance weeds), G (grass), B (broad leaf), S (sedge)



The ability to survive in non host plants indicated the existence of nutrients content in the weed that resembles its host plant nutrient content or the existence of certain substances that could stimulate *D. citri* to stay for feeding. The substances could be a various compounds of alcohol and aldehyde from the leaves of which were specific and volatile used by insect to find its host plant (Visser, 1986). However, the type and number of nutrient content was not as complete as in the host plant for surviving and completing a life cycle. Similar results were also obtained by Sudiono and Purnomo (2008) in Gemini virus isect vectors (*Bemecia tabaci*), and Hardiastono (2001) on Peanut Stripe Virus (PStV). *B. tabaci* were able to live on broadleaf weed *Ageratum conyzoides*. The weeds also serve as a source of inoculum since they were able to be infected by Gemini virus. While *Amaranthus spinosus*, *Bidens pilosa*, *Crotalaria incana*, *Glycine max* and *Physalis angulata* as alternative hosts and source of inoculums for PStV infection.





Table 2. The mean of longevity (days) of *D. citri* on various dominant weeds

Weeds species	Mean ± SE	
Boerhavia erecta Amaranthus spinosus Alternathera philoeroides Portulaca oleraceae	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	a b c c c
Ludwigia perrerlis		

Numbers followed by the same letter are not significantly different between treatments (P < 0,001)



Some of *D. citri* population (31%), obviously did not prefer to stay for feeding on any weeds treatments and just stayed on the cage. Presumably they did not recognize these five kinds of weeds as a host that can be used as a food source. While the absence of *D. citri* population on *P. Oleracea* likely could be caused by high water content in this weeds, which were structurally quite different from their host plants (citrus). The high water content also causes the low content of volatile compounds (specific odors) which were emitted, and were unable to attract *D. citri* stayed for feeding.

The implications of those survivorships of *D. citri* in some weed species is the presence of alternative host for *D. citri* when citrus plant is not available as a food source, either because there are no plants or when plants are treated with pesticides. The existence of alternative host will cause the population of *D. citri* is available throughout the season and serve as initial population for the next generation population. Role of *D. citri* as a propagative vector of CVPD pathogen will also make the disease inoculums always available throughout the season in field. This will further complicate the control measures against CVPD on citrus plant, since the disease transmission is highly dependent on the availability of disease inoculums and insect vector population in the field (Chen, 1998).

CONCLUSION

- 1. Weeds were able as an alternative host for D. citri.
- 2. The mean of longevity of *D. citri* on broad leaf weeds *Bhoeravia erecta* was 5.91 ± 0.251 days.

REFERENCES

- Aubert B. 1990. Integrated activities for the control of *Huanglongbing*-greening and its vector *Diaphorina citri* Kuwayama in Asia. *In:* Aubert B, Tontyaporn S, Buangsuwon D (eds). Proceedings of the Fourth International Asia Pacific Conference on Citrus Rehabilitation, Chiang Mai, Thailand, 4-10 February 1990. Rome: FAO UNDP. pp133-144.
- BPS. 2009. Produksi buah-buahan di Indonesia. <u>http://www.bps.go.id/tab_sub/</u> view.php? tabel=1&daftar=1&id_subyek= 55& notab=2
- Chen CN. 1998. Ecology of the insect vectors of citrus systemic diseases and their control in Taiwan. Food & fertilizer technology center. www.agnet.org/library/data/eb/eb459a/eb459a.pdf

Gomez KA, Gomez AA. 1983. Statistical procedures for agricultural research. Second edition. John Wiley and Sons. New York. 680 pp.



243

- Halbert SE, Manjunath KL. 2004. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. Florida Entomologist 87: 330-353.
- Hardiastono, T. 2001. Potensi gulma berdaun lebar pada pertanian kacang tanah (Arachis hypogaea L.) sebagai inang alternative penyakit bilur peanut stripe virus (PStV). Habitat. 12 (3): 139-146
- Irawan IGP, Sulistyowati L, Wijaya IN. 2003. Penyakit CVPD Pada Tanaman Jeruk (Analisis Baru Berbasis Bioteknologi). Dirjen Perlindungan Hortikultura.

Rukmana dan Saputra. 1999. Gulma dan Teknik Pengendalian. Kanisius. 88 pp.

SPSS Inc. 1999. SPSS[®] for Windows[™] Version 10.0.5. Chicago: SPSS Inc.

- Sudiono dan Purnomo. 2008. Studi kisaran inang kutu kebul (*Bemisia tabaci* Genn.) di sentra sayuran dataran tinggi tanggamus. Jurnal Penelitian Terapan. 8 (3): 103-108.
- Visser, JH. 1986. Host odor perception in herbivores insects. Annual Reviews of Entomology. 31, 121-144
- Xu CF, Xia YH, Li KB, Ke C. 1988. Further study of the Transmission of citrus *Huanglongbing* by a psyllid *Diaphorina citri* kuwayana. Proceeding of 10th Conference of IOCV. Riverside. pp.243-248



ISBN: 978-602-8915-93-9

BOOK 1



International Seminar on Natural Resources, Climate Change and Food Security in Developing Countries

Proceeding









TABLE OF CONTENTS

	Foreword	i – iv
	Kovnoto Speech	v – xi
	The Ministry of Agriculture Republic of Indonesia	
	Table of Contents	xii - xviii
Dian	any Speakers	
Pien	ary Speakers	4 40
1.	Food Insecurity and Climate Change: The Roles of Private Sector in Overcoming The Changes	1 – 13
2.	Genetic Diversity, Climate Uncertainty and Food Security Hugo A. Volkaert	14 – 23
3.	Exploring The Roles of Virtual Classrooms in Biodiversity Conservation Discourses: an Experiential Learning Perspective <i>Ricardo T. Bagarinao</i>	24 – 33
	The second state and Conservation	
Sub	Presentation	
0/4/		34 - 40
4.	Fungus Diversity on Weeds And In Soil after Soybean Harvested	54 - 40
5	Conservation Status of Indigenous Trees in Argao River Watershed	41 – 53
	Reserve, Cebu Philippines: a local initiative	
6	Archiebald Baltazar B. Malaki and Inocencio E. Buot, Jr. Structures and Shape Variations Of The Scales Of Quoy Parrotfish, Scarus	54 – 69
0.	quoyi (Valenciennes 1840)	
_	Arri Q. Adajar, Mark Anthony J. Torres and Cesar G. Demayo	70 – 82
7.	Scales of Viridescent Parrotfish, Calotomus viridescents (Rüppell, 1835)	
	Carlo Stephen Omar Moneva, Mark Anthony J. Torres, Sharon Rose M.	
	Tabugo and Cesar G. Demayo	83 – 97
8.	Colormorphs of The Rice Bug, Leptocorisa oratorius Fabricius, 1794	
	(Hemiptera: Alydidae)	
~	Clifford P. Bendoy, Mark Anthony J. Torres and Cesar G. Demayo	98 – 105
9.	in Flaveria bidentis (C4 Photosinthetic Type)	
	Didik Pudji Restanto	106 - 115
10.	Biodiversity of Java Tea (Orthosiphon Stamineus Benth) Collected From	100 - 115
	Dwi Susanto and Teng D. C. L.	
11.	Biodiversity of Salak Plant (Salacca zalacca (Gaertner) Voss)	116 – 126
	Hari Bowo and Sukartiningrum	127 - 133
12.	Study on Saprophytic Phyloplane Antagonist Against Purple Biolon Disease	121 - 100
	Herv Nirwanto	
13.	Assestment of the Utilization of Ex-Mining Area in	134 – 144
	East Java Province	
4.4	Kemai Wijaya Carotenoids Content of Commercial Seaweed In Bali	145 – 151
14.	K Sri Marhaeni Julvasih	

xi Natural Resources Climate Change and Food Security in Developing Countries Surabaya, Indonesia, June 27-28 2011

xii

15.	Modularity, Covariation and Integration in the Hind Wings of Grasshopper in Family Acrididae	152 – 162
	Kimverly Hazel I. Coronel, Mark Anthony J. Torres and Cesar G.	
16.	Demayo, Arthropods and Weeds Species Diversities on Post Soybean Harvested Kurnia Paramita Sari and Alfi Inayati	163 - 168
17.	Evaluation of Green Open Space Availability in Sleman District Lis Noer Aini	169 – 177
18.	Planning of Soil and Water Conservation on Several Land Use in Watershed Petanu, Gianyar <i>Made Sri Sumarniasih</i>	178 182
19.	Image Analysis of Intra-Population Variability In A Non-Outbreak Population Of The Rice Black Bug Scotinophara SP. From Tacurong, Sultan Kudarat, Philippines	183 - 193
	Mark Anthony J. Torres, Geoffrey Reuel Quinaquin Yañez and Cesar G. Demayo	
20.	The Diversity of Epiphytic Orchid and its Host Tree Along Cemoro Sewu Hiking Pathway, Lawu Mountain, District of Magetan, East Java <i>Nina Dwi Yulia and Sugeng Budiharta</i>	194 – 202
21.	Species Diversity of Seagrasses in Camotes Islands, Central Philippines Serapion N. Tanduyan, Berenice T. Andriano and Ricardo B. Gonzaga	203 - 217
22.	Diversity and Habitat Profile of The Shallow Water Holothurians in Camotes Islands, Central Philippines	218 – 229
	Serapion N. Tanduyan, Berenice T. Andriano, Ricardo B. Gonzaga,	
23.	Effect of Land-Use Change on Diversity And Ecosystem Functions of Insects in Central Sulawesi and Its Conservation Effort	230 – 238
	Shahabuddin	
24.	The Longevity of Diaphorina Citri on Various Dominant Weeds Siwi Hardiastuti and Mofit Eko Poerwanto	239 244
25.	Morphological Characters of Kopyor Coconut Grown in Sumenep, Madura, Indonesia	245 - 254
26.	Sukendah, Zainal Abidin, Sri Wiyatiningsih and Bakti Wisnu Wijayanti Occurence of Bacterial Soft Rot of Phalaenopsis Orchids in Yogyakarta and West Java, Indonesia	255 265
	Tri Joko, Dwi Kiswanti, Hanudin and Siti Subandiyah	
27.	Macro Algae Diversity in Madura Island Wisanti, Evie Ratnasari, Novita Indah Kartika	266 – 275
28.	Land Snail Diversity of MT. Polis, Central Cordillera Range, Luzon Island, Philippines	276 - 293
	Zenaida G. Baoanan and Steve P. Obanan	
Sub Post	Theme : <u>Biodiversity and Conservation</u> er Presentation	
29.	The Development of Nests Collocalia esculenta in Jombang, East Java Novita Kartika Indah and Sunu Kuntjoro	294 – 298
30	Lake Dense, Son Econologie, Control, Philippings, its Status and	000 045

 30.
 Lake Danao, San Francisco, Central, Philippines: its Status and
 299 – 315

 Government Initiatives for its Conservation and Sustainability
 Serapion N. Tanduyan*, Berenice T. Andriano and Ricardo B. Gonzaga

Sub Theme : <u>Adaptation and Production</u> Oral Presentation

Ului	//cscination	
31.	Study of Mango Plant Reproductive Phase's (Mangifera Indica.I) Arumanis 143	316 – 322
32.	Bambang Priyanto and W. Guntoro Land Characteristics To Increase Sugar Content Of Melon Fruit (Cucumis melo)	323 – 339
33.	Bakti Wisnu W, Purwadi and Siswanto Soil Tillage Effect on Growth and Development of Root System of Ten Mango Seedling Varieties	340 - 352
34.	Djarwatiningsih PS and Agus Sulistyono Changes in Leaves and Stem Diameter of Cocoa Tree (Theobroma Cacao I.) in Drought Stress	353 - 361
35.	Salinity Tolerance of Baluran Variety of Soybean (Glycine max. L) by in Vitro culture	362 368
36.	Etty Handayani, Agus Nugroho Setiawan, Slamet Wahyudiyanto Effect of Mulching by Corn Biomass on Growth and Production of Corn in Drought Condition	369 – 376
37.	Denna Eriani Munandar Effect of Fertilizer Mixed Macro Ammonium Sulfate Nitrate on The Growth And Production Shallot In Malang	377 – 386
38.	Effect of Plant Strengthening Agents on Pak Choi Grown Organically Farida	387 – 392
39.	Molecular Analysis of Genetic Stability in Micropropagated Citrus Regenerated Through Somatic Embryogenesis Using Inter-simple Sequence Repeat (ISSR) Markers	393 – 400
40.	Farida Yulianti, Nirmala F. Devy and Hardiyanto Effect of Planting System to IP Rice 400 for Development of Pest Diseases Handoko and Evy Latifah	401 – 406
41.	Agronomic Aspects of Intercropping Between Energy Crop of Castor (Ricinus communis I.) and Pulses	407 – 415
42.	Study of Preference of Diaphorina Citri Kuw.(homoptera : psyllidae) on Several Types of Orange Crops (citrus spp.)	416 – 422
43.	Indriya Radiyanto and Ketut Sri Marhaeni J. Study of Manure Addition (Biogreen and Fine Compost) on Za Fertilizer to Growth and Yield of Green Mustard	423 – 428
44.	Developing High Yield and Tolerant to Low Nitrogen Tomato Genotypes Makhziah and Ida Retno Moeljani	429 – 437
45.	Nitrogen Use Efficiency and Yielding Ability of Four Maize Varieties Makhziah and Yonny Koentjoro	438 – 443
46.	Hybrid Vigor of Crossing Among Japanese and Indonesian Soybean <i>M. Setyo Poerwoko</i>	444 452
47.	Adaptation of Food Crops to Climate Change Moch. Sodiq and Wiludjeng Widayati	453 - 460
48.	Mass Propagation of Five Citrus Cultivars Through Somatic Embryogenesis Technique <i>Nirmala Friyanti Devy, Farida Yulianti and Hardiyant</i> o	461 - 471

xiv

ł

49.	Tolerance Levels of Ornamental Shrubs to Urban Air Pollution in Surabaya City Based on Air Pollution Tolerance Index (APTI)	472 – 479
	Pangesti Nugrahani, Endang Tri Wanyu, Sugijanto and	
50.	Hery Purnobasuki Application of Organic Mulches on Cultivation of Red Pepper (Capsicum annuum, L.) in different seasons	480 – 488
51.	Puji Harsono and Dja'far Shiddieq Qualitative and Quantitative Analysis of Scale Shape Variation Between Different Body Regions of Redbreast Wrasse (Cheilinus fasciatus)	489 – 502
52.	Queenilyn B. Albutra Influence of Planting Distance and Pruning on Growth and Development of Sengon (Albizzia Falcataria L.)	503 – 515
53.	Ramdan Hidayat, B.W. Wijayani and Sukendah Agronomic and Physiological Responses of Maize Cultivarsto Low Nutrient	516 – 526
	Supply	
54.	Determination of Efficiency in Extraction Anthocyanin of Teleng Flower (Clitoria ternatea L)	527 – 535
55.	Retno Endrasari, Pudji Hastuti and Sri Anggrahini Rate and Time Nitrogen Application on The Growth and Yield of Two Varieties Oilseed Rape (Brassica napus L.)	536 - 544
56.	Sigit Soeparjono Resistance of 6 Shallot Cultivars to Moler Disease in Field Sri Wivatiningsih, Arif Wibowo, and Endang Triwahyu P.	545 - 552
57.	Declining of Apple Production at Three Production Centers of East Java Suhariyono	553 - 565
58.	Chicken Manure Application to Improve The Growth of Paddy Grown on Critical Land of Ex-Brick Industry	566 – 572
59.	Sundahri Characterization of EGCG Compound Use ¹ H NMR Spectrum on Camellia Sinansis (L) Callus	573 – 578
	Sutini Tetik W Sutiman B, and R. Verpoorte	
60.	Impact The Lapindo Mudflow toward Biosafety of Fisheries Product at East Coastal of Sidoarjo District	579 – 589
	Tarzan Purnomo	590 - 594
61.	Bactrocera (Diptera: Tephritidae)	
62.	Characteristic of Growth and Production of Oyster Mushrooms at Medium Land at Litter Media	595 – 603
63.	Widiwurjani and Guniarti Growth Pattern of Coffee Seeds on Various Water Potential Pressure Yonny Koentioro and Makhziah	604 - 613
64.	Diversity Technology Strawberry Cultivation in Different Regional Production	614 - 626
	Zainuri Hanif and Emi Budiyati	
Sut	Theme : Adaptation and Production	

Poster Presentation

65. Effect of Foliar Fertilizer Use on The Growth and Production Of Chinese 627 – 640 Cabbage/ Caisim (Brassica juncea) AI. Gamal Pratomo and Luki Rosmahani

XV

ISBN

66.	Test The Effectiveness of Organic Fertilizer Super "Zoo-Mass" on the Growth and Yield of Mustard (Brassica Juncea L.)	641 – 654
67.	Amik Krismawati and Dini Hardini Application Integrated Crop Management Approach (PTT) New Variety of Rice in Rice Field Integration with Beef in Madiun Residen	655 – 667
68.	Amik Krismawati and Setiasin Improvement Feeding Management of Pre Weaning Twinning Calf in East Java	668 – 673
69.	Dini Hardini, Hendri Arianto and Irian HD Potency of Twin Calving of Some Beef Cattle Local Breed In East Java D. Hardini and Irfan HD	674 – 678
70.	Fruit Fly Responses to Plant Type and Distillates Basil Plant on Mango Eli Korlina, Diding Rachmawati, Luki Rosmahani and Sri Yunlastuti	679 - 684
71.	Productivity and Quality of 6 Varieties of Superior Grapevine Plante at High Land of Tlekung (950 m asl)	685 — 693
72.	Emi Budiyati and Anis Andrini Determinants Factors in Technology Development with Internal Input Raw Materials Dirt Cow	694 - 702
73.	Hadi Suhardjono and Yonny Koentjoro Productivity of Three Dwarf Kopyor Coconut Varieties From Pati, Central	703 – 711
74.	Ismail Maskromo, Hengky Novarianto, Sukendah, and Sudarsono Simultaneous Infection of Mandarin (citrus reticulata blanko.) with Tristeza Virus and Liberobacter Asiatus in Indonesia	712 – 724
75.	Mutia Erti Dwiastuti The Assessment of Production And Resistance to Purple Spot Disease (Altenaria Porri) on Lines Shallot (Allium cepa) Potentially High Production	725 – 733
76.	<i>N. Istiqomah, E. Korlina, and Z. Arifin</i> Characteristic Agronomic Components and Organoleptic Test Eleven of Lines Paddy (<i>Oryza Sativa</i>)	734 - 743
77.	N. Istiqomah and S.S. Antarlina Application of Integrated Crop Management (ICM) to Increase Productivity, Revenue and Efficiency of Rice Farming	744 - 753
78.	Purwanto and Suwono Soybean Productivity Improvement through Innovation Technology O. Dedang Ernawanto dan Noeriwan B.S.	754 – 763
79.	Adaptation of the Stem Structure Mangrove on the Est Coast Surabaya Rinie Pratiwi. J. Dioko Budiono, Enny Susiywati and Nur Hidayah	764 – 777
80.	Action Research of Feeding Fermentation Treatment Using Complete Local Raw Materials from Agricultural Waste to Fattening Sheep at Tulupgagung District	778 – 785
81.	Setiasih, A.M. Abdurrahman, D.W. Astuti and S. Purnomo Upland Rice Production Test and Pest Disease Attack Rate Sugiono and Zainal Arifin	786 – 795
82.	The Source of Potassium Fertilizer and its Effect on The Increasing of Rice Yield in The Vertisol	796 – 806
83.	Suwono and Purwanto Evaluation of Picking Tool Application Assessment on Three Arumanis/Gadung Mango Plantations	807 – 814
	wanyuningyawau ang Shi narwang	

xvi

Ŧ

ISBN :		978-602-8915-93-9	
--------	--	-------------------	--

Book 1

84.	The Growth of Leguminoceae in Calcareous Soil Yuliani and Yuni Sri Rahayu	815 - 828
85.	Improvement of Rice and Corn Cropping System in One Cropping Pattern on Rainfed Lowland	829 - 840
86.	Improving Productivity of Jatropha More Than 30 Percent (>10 Ton/Ha) Using Aplication of Kno3 To Flowering Faster (>30%) and Honey Bee Apis Mellifera (Apidae) Pollination To Increasing of Fruit (>30%) on Dry Climate of Dry Land	841 - 846
	F. Deru Dewanti, Budi Hariyono and Agus Sulistyono	
87.	Benthic Meiofaunal Composition in Selected Sandy Substrates Along Pala- O River, Iligan City, Philippines <i>Ifeche, A. A., M. A. D. Delos Angeles, F. I. V. Cabahug, J. A. R. S. Lopez,</i> A. S. Oliveros ¹ And J. A. Godinez	847 – 854
88.	Estimation of Primary Productivity and Periphyton Biomass in The Water Reservior of Agus Iv Hydroelectric Plant In Nangka, Balo-I, Lanao Del Norte, Philippines (A High School Research Project)	855 - 871
99	LIST OF PARTICIPANTS	872 - 885