



**ISNAR-C2FS 2011**

# **International Seminar on Natural Resources, Climate Change and Food Security in Developing Countries**

## **Proceeding**





International Seminar Natural Resources, Climate Change and Food Security  
in Developing Countries - ISNAR C2FS  
June 27 - 28, 2011 Graha Pena Building, Surabaya - Indonesia  
ISBN : 978-602-8915-93-9

---

### **Editorial Board**

*Alferd Hugo Volkaert, PhD. (CAB - University of Kasetsart, Thailand)*

*Zenaida Baoanan, PhD. (University of The Philippines Baguio, Philippines)*

*Katayama Takeshi, Prof. PhD. (University of Kagawa, Japan)*

*Sukendah, Dr. Ir. MSc. (University of Pembangunan nasional "Veteran" East Java, Indonesia)*

*Wani Hadi Utomo, Prof. PhD. (University of Tribuana, Malang, Indonesia)*

*Ricardo Bagarinao, PhD. (University of The Philippines Open University, Philippines)*

*Lin Qing, Prof. PhD. (School of Economic Fujian Normal University, China)*

*Surya Rathore, Prof. PhD. (University of Agriculture and Technology, Pantnagar, India)*

### **Editor :**

*Yonny Koentjoro*

### **Assistant Editor :**

*Wahyu Santoso*

*Pangesti Nugrahani*

*Makhziah*

### **Cover :**

*Magnificent Digital Printing by : Purnomo Edi Sasongko*

### **Proceeding design :**

*Yonny Koentjoro and Wahyu Santoso*

Copyright : December 2011

Published by : Faculty of Agriculture, University of Pembangunan Nasional "Veteran" East Java, Indonesia

### **Held in between :**

- *Faculty of Agriculture, University of Pembangunan Nasional "Veteran" East Java, Indonesia*
- *Philippine Society for The Study of Nature (PSSN), Philippine*
- *CAB - University of Kasetsart, Thailand*

---

**TABLE OF CONTENTS**

Foreword	<i>i – iv</i>
Keynote Speech The Ministry of Agriculture Republic of Indonesia	<i>v – xi</i>
Table of Contents	<i>xii - xviii</i>
<b>Plenary Speakers</b>	
1. Food Insecurity and Climate Change: The Roles of Private Sector in Overcoming The Changes <b>Bustanul Arifin</b>	<i>1 – 13</i>
2. Genetic Diversity, Climate Uncertainty and Food Security <b>Hugo A. Volkaert</b>	<i>14 – 23</i>
3. Exploring The Roles of Virtual Classrooms in Biodiversity Conservation Discourses: an Experiential Learning Perspective <b>Ricardo T. Bagarinao</b>	<i>24 – 33</i>
<b>Sub Theme : <u>Biodiversity and Conservation</u></b>	
<b>Oral Presentation</b>	
4. Fungus Diversity on Weeds And In Soil after Soybean Harvested <b>Alfi Inayati and Kurnia Paramita</b>	<i>34 – 40</i>
5. Conservation Status of Indigenous Trees in Argao River Watershed Reserve, Cebu Philippines: a local initiative <b>Archiebald Baltazar B. Malaki and Inocencio E. Buot, Jr.</b>	<i>41 – 53</i>
6. Structures and Shape Variations Of The Scales Of Quoy Parrotfish, <i>Scarus quoyi</i> (Valenciennes 1840) <b>Arri Q. Adajar, Mark Anthony J. Torres and Cesar G. Demayo</b>	<i>54 – 69</i>
7. Qualitative and Geometric Morphometric Description of The Shapes of Scales of Viridescent Parrotfish, <i>Calotomus viridescens</i> (Rüppell, 1835) <b>Carlo Stephen Omar Moneva, Mark Anthony J. Torres, Sharon Rose M. Tabugo and Cesar G. Demayo</b>	<i>70 – 82</i>
8. Describing Head Shape Variation Between Sexes and Among Two Colormorphs of The Rice Bug, <i>Leptocorisa oratorius</i> Fabricius, 1794 (Hemiptera:Alydidae) <b>Clifford P. Bendoy, Mark Anthony J. Torres and Cesar G. Demayo</b>	<i>83 – 97</i>
9. Genomic Fragment Gene of Carbonic Anhydrase (Ca) in <i>Flaveria bidentis</i> (C4 Photosinthetic Type) <b>Didik Pudji Restanto</b>	<i>98 – 105</i>
10. Biodiversity of Java Tea ( <i>Orthosiphon Stamineus</i> Benth) Collected From Indonesia and Malaysia <b>Dwi Susanto and Teng D. C. L.</b>	<i>106 – 115</i>
11. Biodiversity of Salak Plant ( <i>Salacca zalacca</i> (Gaertner) Voss) <b>Hari Bowo and Sukartiningrum</b>	<i>116 – 126</i>
12. Study on Saprophytic Phyloplane Antagonist Against Purple Blotch Disease on Onion Caused by <i>Alternaria porri</i> IN VITRO <b>Hery Nirwanto</b>	<i>127 – 133</i>
13. Assesment of the Utilization of Ex-Mining Area in East Java Province <b>Kemal Wijaya</b>	<i>134 – 144</i>
14. Carotenoids Content of Commercial Seaweed In Bali <b>K. Sri Marhaeni Julyasih</b>	<i>145 – 151</i>



- |     |   |             |
|-----|---|-------------|
| 15. | Modularity, Covariation and Integration in the Hind Wings of Grasshopper in Family Acrididae<br><b>Kimverly Hazel I. Coronel, Mark Anthony J. Torres and Cesar G. Demayo,</b>   | 152 – 162   |
| 16. | Arthropods and Weeds Species Diversities on Post Soybean Harvested<br><b>Kurnia Paramita Sari and Alfi Inayati</b>  | 163 – 168   |
| 17. | Evaluation of Green Open Space Availability in Sleman District<br><b>Lis Noer Aini</b>  | 169 – 177   |
| 18. | Planning of Soil and Water Conservation on Several Land Use in Watershed Petanu, Gianyar<br><b>Made Sri Sumarniasih</b>   | 178 – 182   |
| 19. | Image Analysis of Intra-Population Variability In A Non-Outbreak Population Of The Rice Black Bug <i>Scotinophara</i> SP. From Tacurong, Sultan Kudarat, Philippines<br><b>Mark Anthony J. Torres, Geoffrey Reuel Quinaquin Yañez and Cesar G. Demayo</b> | 183 – 193   |
| 20. | The Diversity of Epiphytic Orchid and its Host Tree Along Cemoro Sewu Hiking Pathway, Lawu Mountain, District of Magetan, East Java<br><b>Nina Dwi Yulia and Sugeng Budiharta</b>   | 194 – 202   |
| 21. | Species Diversity of Seagrasses in Camotes Islands, Central Philippines<br><b>Serapion N. Tanduyan, Berenice T. Andriano and Ricardo B. Gonzaga</b>   | 203 – 217   |
| 22. | Diversity and Habitat Profile of The Shallow Water Holothurians in Camotes Islands, Central Philippines<br><b>Serapion N. Tanduyan, Berenice T. Andriano, Ricardo B. Gonzaga, Panfilo E. Ciriaco, Wilfredo G. Anos, Lourdes M. Garciano</b>               | 218 – 229   |
| 23. | Effect of Land-Use Change on Diversity And Ecosystem Functions of Insects in Central Sulawesi and Its Conservation Effort<br><b>Shahabuddin</b>   | 230 – 238   |
| 24. | The Longevity of <i>Diaphorina Citri</i> on Various Dominant Weeds<br><b>Siwi Hardiastuti and Mofit Eko Poerwanto</b>   | 239 – 244 ✓ |
| 25. | Morphological Characters of Kopyor Coconut Grown in Sumenep, Madura, Indonesia<br><b>Sukendah, Zainal Abidin, Sri Wiyatiningsih and Bakti Wisnu Wijayanti</b>   | 245 – 254   |
| 26. | Occurrence of Bacterial Soft Rot of <i>Phalaenopsis</i> Orchids in Yogyakarta and West Java, Indonesia<br><b>Tri Joko, Dwi Kiswanti, Hanudin and Siti Subandiyah</b>  | 255 – 265   |
| 27. | Macro Algae Diversity in Madura Island<br><b>Wisanti, Evie Ratnasari, Novita Indah Kartika</b>  | 266 – 275   |
| 28. | Land Snail Diversity of MT. Polis, Central Cordillera Range, Luzon Island, Philippines<br><b>Zenaida G. Baoanan and Steve P. Obanan</b>   | 276 – 293   |

**Sub Theme : Biodiversity and Conservation**  
**Poster Presentation**

- |     |   |           |
|-----|---|-----------|
| 29. | The Development of Nests <i>Collocalia esculenta</i> in Jombang, East Java<br><b>Novita Kartika Indah and Sunu Kuntjoro</b>   | 294 – 298 |
| 30. | Lake Danao, San Francisco, Central, Philippines: its Status and Government Initiatives for its Conservation and Sustainability<br><b>Serapion N. Tanduyan*, Berenice T. Andriano and Ricardo B. Gonzaga</b> | 299 – 315 |



**Sub Theme : Adaptation and Production**  
**Oral Presentation**

- |     |   |           |
|-----|---|-----------|
| 31. | Study of Mango Plant Reproductive Phase's (Mangifera Indica.l)<br>Arumanis 143<br><b>Bambang Priyanto and W. Guntoro</b>  | 316 – 322 |
| 32. | Land Characteristics To Increase Sugar Content Of Melon Fruit (Cucumis melo)<br><b>Bakti Wisnu W, Purwadi and Siswanto</b>  | 323 – 339 |
| 33. | Soil Tillage Effect on Growth and Development of Root System of Ten Mango Seedling Varieties<br><b>Djarwatiningsih PS and Agus Sulistyono</b>   | 340 – 352 |
| 34. | Changes in Leaves and Stem Diameter of Cocoa Tree (Theobroma Cacao l.) in Drought Stress<br><b>Erma Prihastanti</b>   | 353 – 361 |
| 35. | Salinity Tolerance of Baluran Variety of Soybean (Glycine max. L) by in Vitro culture<br><b>Etty Handayani, Agus Nugroho Setiawan, Slamet Wahyudiyanto</b>  | 362 – 368 |
| 36. | Effect of Mulching by Corn Biomass on Growth and Production of Corn in Drought Condition<br><b>Denna Eriani Munandar</b>  | 369 – 376 |
| 37. | Effect of Fertilizer Mixed Macro Ammonium Sulfate Nitrate on The Growth And Production Shallot In Malang<br><b>Dwi Setyorini and Zaenal Arifin</b>  | 377 – 386 |
| 38. | Effect of Plant Strengthening Agents on Pak Choi Grown Organically<br><b>Farida</b>   | 387 – 392 |
| 39. | Molecular Analysis of Genetic Stability in Micropropagated Citrus Regenerated Through Somatic Embryogenesis Using Inter-simple Sequence Repeat (ISSR) Markers<br><b>Farida Yulianti, Nirmala F. Devy and Hardiyanto</b> | 393 – 400 |
| 40. | Effect of Planting System to IP Rice 400 for Development of Pest Diseases<br><b>Handoko and Evy Latifah</b>   | 401 – 406 |
| 41. | Agronomic Aspects of Intercropping Between Energy Crop of Castor (Ricinus communis l.) and Pulses<br><b>I Komang Damar Jaya</b>   | 407 – 415 |
| 42. | Study of Preference of Diaphorina Citri Kuw.(homoptera : psyllidae) on Several Types of Orange Crops (citrus spp.)<br><b>Indriya Radiyahanto and Ketut Sri Marhaeni J.</b>  | 416 – 422 |
| 43. | Study of Manure Addition (Biogreen and Fine Compost) on Za Fertilizer to Growth and Yield of Green Mustard<br><b>Juli Santoso and Yunia Ulfah</b>   | 423 – 428 |
| 44. | Developing High Yield and Tolerant to Low Nitrogen Tomato Genotypes<br><b>Makhziah and Ida Retno Moeljani</b>   | 429 – 437 |
| 45. | Nitrogen Use Efficiency and Yielding Ability of Four Maize Varieties<br><b>Makhziah and Yonny Koentjoro</b>   | 438 – 443 |
| 46. | Hybrid Vigor of Crossing Among Japanese and Indonesian Soybean<br><b>M. Setyo Poerwoko</b>  | 444 – 452 |
| 47. | Adaptation of Food Crops to Climate Change<br><b>Moch. Sodik and Wiludjeng Widayati</b>   | 453 – 460 |
| 48. | Mass Propagation of Five Citrus Cultivars Through Somatic Embryogenesis Technique<br><b>Nirmala Friyanti Devy, Farida Yulianti and Hardiyanto</b>   | 461 – 471 |



49. Tolerance Levels of Ornamental Shrubs to Urban Air Pollution in Surabaya City Based on Air Pollution Tolerance Index (APTI)  
**Pangesti Nugrahani, Endang Tri Wahyu, Sugijanto and Hery Purnobasuki** 472 – 479
50. Application of Organic Mulches on Cultivation of Red Pepper (*Capsicum annuum*, L.) in different seasons  
**Puji Harsono and Dja'far Shiddieq** 480 – 488
51. Qualitative and Quantitative Analysis of Scale Shape Variation Between Different Body Regions of Redbreast Wrasse (*Cheilinus fasciatus*)  
**Queenilyn B. Albutra** 489 – 502
52. Influence of Planting Distance and Pruning on Growth and Development of Sengon (*Albizia Falcataria* L.)  
**Ramdan Hidayat, B.W. Wijayani and Sukendah** 503 – 515
53. Agronomic and Physiological Responses of Maize Cultivarsto Low Nutrient Supply  
**Renih Hayati, Asmawati and Munandar** 516 – 526
54. Determination of Efficiency in Extraction Anthocyanin of Teleng Flower (*Clitoria ternatea* L.)  
**Retno Endrasari, Pudji Hastuti and Sri Anggrahini** 527 – 535
55. Rate and Time Nitrogen Application on The Growth and Yield of Two Varieties Oilseed Rape (*Brassica napus* L.)  
**Sigit Soeparjono** 536 – 544
56. Resistance of 6 Shallot Cultivars to Moler Disease in Field  
**Sri Wiyatiningsih, Arif Wibowo, and Endang Triwahyu P.** 545 – 552
57. Declining of Apple Production at Three Production Centers of East Java  
**Suhariyono** 553 – 565
58. Chicken Manure Application to Improve The Growth of Paddy Grown on Critical Land of Ex-Brick Industry  
**Sundahri** 566 – 572
59. Characterization of EGCG Compound Use <sup>1</sup>H NMR Spectrum on Camellia Sinensis (l.) Callus  
**Sutini, Tatik W, Sutiman B, and R. Verpoorte** 573 – 578
60. Impact The Lapindo Mudflow toward Biosafety of Fisheries Product at East Coastal of Sidoarjo District  
**Tarzan Purnomo** 579 – 589
61. The Effect of The Volume of Methyl Eugenol to Total Catch Fruit Flies *Bactrocera* (Diptera: Tephritidae)  
**Tjipto Haryono** 590 – 594
62. Characteristic of Growth and Production of Oyster Mushrooms at Medium Land at Litter Media  
**Widiwurjani and Guniarti** 595 – 603
63. Growth Pattern of Coffee Seeds on Various Water Potential Pressure  
**Yonny Koentjoro and Makhziah** 604 – 613
64. Diversity Technology Strawberry Cultivation in Different Regional Production Centre  
**Zainuri Hanif and Emi Budiayati** 614 – 626

**Sub Theme : Adaptation and Production**  
**Poster Presentation**

65. Effect of Foliar Fertilizer Use on The Growth and Production Of Chinese Cabbage/ Caisim (*Brassica juncea*)  
**Al. Gamal Pratomo and Luki Rosmahani** 627 – 640



66. Test The Effectiveness of Organic Fertilizer Super "Zoo-Mass" on the Growth and Yield of Mustard (*Brassica Juncea* L.) 641 – 654  
**Amik Krismawati and Dini Hardini**
67. Application Integrated Crop Management Approach (PTT) New Variety of Rice in Rice Field Integration with Beef in Madiun Residen 655 – 667  
**Amik Krismawati and Setiasih**
68. Improvement Feeding Management of Pre Weaning Twinning Calf in East Java 668 – 673  
**Dini Hardini, Hendri Arianto and Irfan HD**
69. Potency of Twin Calving of Some Beef Cattle Local Breed In East Java 674 – 678  
**D. Hardini and Irfan HD**
70. Fruit Fly Responses to Plant Type and Distillates Basil Plant on Mango 679 – 684  
**Eli Korlina, Diding Rachmawati, Luki Rosmahani and Sri Yunlastuti**
71. Productivity and Quality of 6 Varieties of Superior Grapevine Plante at High Land of Tlekung (950 m asl) 685 – 693  
**Emi Budiyati and Anis Andriani**
72. Determinants Factors in Technology Development with Internal Input Raw Materials Dirt Cow 694 – 702  
**Hadi Suhardjono and Yonny Koentjoro**
73. Productivity of Three Dwarf Kopyor Coconut Varieties From Pati, Central Java – Indonesia 703 – 711  
**Ismail Maskromo, Hengky Novarianto, Sukendah, and Sudarsono**
74. Simultaneous Infection of Mandarin (*citrus reticulata* blanko.) with *Tristeza* Virus and *Liberobacter Asiatus* in Indonesia 712 – 724  
**Mutia Erti Dwiastuti**
75. The Assessment of Production And Resistance to Purple Spot Disease (*Altenaria Porri*) on Lines Shallot (*Allium cepa*) Potentially High Production 725 – 733  
**N. Istiqomah, E. Korlina, and Z. Arifin**
76. Characteristic Agronomic Components and Organoleptic Test Eleven of Lines Paddy (*Oryza Sativa*) 734 – 743  
**N. Istiqomah and S.S. Antarlina**
77. Application of Integrated Crop Management (ICM) to Increase Productivity, Revenue and Efficiency of Rice Farming 744 – 753  
**Purwanto and Suwono**
78. Soybean Productivity Improvement through Innovation Technology 754 – 763  
**Q. Dadang Ernawanto dan Noeriwan B.S.**
79. Adaptation of the Stem Structure Mangrove on the Est Coast Surabaya 764 – 777  
**Rinie Pratiwi, J. Djoko Budiono, Enny Susiywati and Nur Hidayah**
80. Action Research of Feeding Fermentation Treatment Using Complete Local Raw Materials from Agricultural Waste to Fattening Sheep at Tulungagung District 778 – 785  
**Setiasih, A.M. Abdurrahman, D.W. Astuti and S. Purnomo**
81. Upland Rice Production Test and Pest Disease Attack Rate 786 – 795  
**Sugiono and Zainal Arifin**
82. The Source of Potassium Fertilizer and its Effect on The Increasing of Rice Yield in The Vertisol 796 – 806  
**Suwono and Purwanto**
83. Evaluation of Picking Tool Application Assessment on Three Arumanis/Gadung Mango Plantations 807 – 814  
**Wahyunindyawati and Sri Harwanti**



- 
- |     |  |           |
|-----|--|-----------|
| 84. | The Growth of Leguminosae in Calcareous Soil<br>Yuliani and Yuni Sri Rahayu  | 815 – 828 |
| 85. | Improvement of Rice and Corn Cropping System<br>in One Cropping Pattern on Rainfed Lowland<br><b>Zainal Arifin, Sugiono and Dwi Setyorini</b>  | 829 – 840 |
| 86. | Improving Productivity of Jatropha More Than 30 Percent (>10 Ton/Ha)<br>Using Application of K <sub>2</sub> O To Flowering Faster (>30%) and Honey Bee Apis<br>Mellifera (Apidae) Pollination To Increasing of Fruit (>30%) on Dry Climate<br>of Dry Land<br><b>F. Deru Dewanti, Budi Hariyono and Agus Sulistyono</b> | 841 - 846 |
| 87. | Benthic Meiofaunal Composition in Selected Sandy Substrates Along Pala-<br>O River, Iligan City, Philippines<br><b>Ifeche, A. A., M. A. D. Delos Angeles, F. I. V. Cabahug, J. A. R. S. Lopez,<br/>A. S. Oliveros<sup>1</sup> And J. A. Godinez</b>  | 847 – 854 |
| 88. | Estimation of Primary Productivity and Periphyton Biomass in The Water<br>Reservoir of Agus Iv Hydroelectric Plant In Nangka, Balo-I, Lanao Del<br>Norte, Philippines (A High School Research Project)<br><b>Rona Kriszelle P. Jauculan and Jesha Faye T. Librea</b>   | 855 - 871 |
| 99. | <b>LIST OF PARTICIPANTS</b>  | 872 - 885 |





## THE LONGEVITY OF *Diaphorina citri* ON VARIOUS DOMINANT WEEDS

Siwi Hardiastuti, Mofit Eko Poerwanto\*

Agrotechnology Department, University of Pembangunan Nasional "Veteran" Yogyakarta  
\*mofitnuk@yahoo.com, telp/fax: (0274) 487793

### ABSTRACT

Citrus vein phloem degeneration (CVPD) is the most impediment disease in citrus production in the world. It mainly vectored by *Diaphorina citri* Kuwayama. Intensive application of insecticides is ineffective and also costly. Various weeds species in citrus orchard are suspected as alternative host of vector. Research on the longevity of *D. citri* on the various weeds is needed for completing the integrated CVPD disease management program. Each of five dominant weeds gained from vegetation analysis were exposed to five days old *D. citri* adults inside the nylon mesh cage, which previously starved for 24 hours. The number of *D. citri* survived was recorded daily (24 h). Analysis of variance and DMRT were done on data. Most of *D. citri* stayed for feeding on *Boerhavia erecta* and survived for 5.91 days. It was significantly different on *Amaranthus spinosus*, and also on *Althernanthera philoxeroides*, *Portulaca oleracea*, *Ludwigia perrenis* which survived for 3.57, 0.20, 0.00, and 0.20 days respectively.

*Key words: CVPD, longevity, citrus, Diaphorina citri*

### INTRODUCTION

Citrus vein phloem degeneration (CVPD) is the most impediment disease in citrus culture in the world and has been attacking citrus in Asia and America (Aubert, 1990; Halbert & Majunath, 2004). Many highly productive plantations where yields were 20 tones per ha decreased sharply into 8.6-15 tones per ha per annum (Irawan *et al.*, 2003). Recently, citrus production in Indonesia has developed rapidly. In 2007, the production increased to 2,625,884 tones. However, CVPD is still a great threat to the citrus production in Indonesia. In 2008, the production decreased again into 2,467,632 tonnes (BPS, 2009).

CVPD mainly vectored by *D. citri* Kuwayama. Fourth and fifth instars as well as adults are able to transmit diseases (Xu *et al.*, 1988; Aubert, 1990). Its role as the insect vector, led to the vector control using insecticides as the principal strategies in CVPD control management. However, significant result of the CVPD disease control has not yet been obtained in the field due to the insufficient understanding of the life of a vector. The existence of various weeds in citrus orchards could be suspected as alternative host of vector (*D. citri*). Some weeds are able as alternative host of pests and pathogens. Kolomento weeds (*Leersia hexandra*) and Sembungan (*Sacciolepis*



*interrupta*) are the host plants of rice stem borer, Cerulang (*Eleusine indica*) is the host plants of *Piricularia oryzae*, and nematodes *Rotylenchus reniformis*. Wuluhan (*Setaria plicata*) and Jajahean or Lempuyangan (*Panicum sp.*) are host plant of rice bug (Rukmana and Saputra, 1999). In order to complete an integrated control program of CVPD, research on *D. citri* longevity in various weeds species should be conducted.

## MATERIALS AND METHOD

### *D. citri* mass rearing

Eggs of *D. citri* was obtained from the field and maintained on orange jasmine plants. The nymphs which hatched from eggs reared on orange jasmine shoots, planted in pots (Ø 17 cm, height 12 cm and Ø 25 cm, height 18 cm) and covered with nylon mesh cages (length 100 cm, width 100 cm, height 100 cm).

### *Weeds identification and vegetation analysis of citrus orchards*

Dominant citrus weeds was determined using analysis vegetation method by taking five ring samples (50 cm x 50 cm) per plots randomly on three citrus orchards. Values of ADe (absolute density), AF (absolute frequency), ADo (absolute dominance), RDe (relative density), RF (relative frequency), and RDo (relative dominance) were calculated and value of SDR (summed dominance ratio) was determined.

### *D. citri* longevity test on weeds

Experiments were carried out by using five dominant weeds planted in pots. Each weed was inserted into nylon mesh cage (Ø 10 cm, height 30 cm) and 10 of five days old *D. citri* adults were released in the middle of the cage. They were previously starved for 24 hours before releasing. The cages were covered with sponge (Ø 10 cm, 2 cm thick). The number of *D. citri* survived on each weed was observed every 24 hours up to all of them in dead. The treatment was repeated 10 times.

### *Data analysis*

Analysis of variance at 95% significance level was done by using SPSS 10.0.5 (SPSS, 1999) to determine the effect of treatment. Duncan's multiple range test (DMRT) at 95% significance level was carried out to identify significant difference between treatments (Gomez and Gomez, 1983).



## RESULTS AND DISCUSSION

Results of vegetation analysis on weeds of citrus orchard found 16 species of weeds which consists of three species of grass, 10 species of broadleaf, and three species of sedges group (Table 1.). Based on these results, five most dominant weed species were taken for testing *D. citri* longevity; there were *Alternanthera philoeroides* of grass group, *Oleraceae Portulaca*, *Amaranthus spinosus*, *Ludwigia perrenis*, and *Boerhavia erecta* of broadleaf weeds with SDR values of 19.72%, 12.06%, 11.65%, 9.31% and 6.93% respectively. The analysis result also shows that the value of absolute density (ADe) was not necessarily generating high values of SDR, because the value of SDR was determined by three parameters: density (De), dominance (Do), and frequency (F). The highest value of SDR on *A. philoeroides* indicated that this weed was the most dominant weeds in the field.

*D. citri* was able to survive a maximum of eight days with the mean longevity of  $5.91 \pm 0.251$  days on the broadleaf weed *B. erecta*, but was not able to complete one life cycle (Figure 1.). It was significantly different to other weeds, *A. spinosus*, *A. philoeroides*, *L. perrenis* and *P. oleraceae* which were survived for 3.57, 0.20, 0.20 and 0.00 days respectively (Table 2.).

Table 1. Vegetation analysis on weeds of citrus orchards

Weeds species	Group	Ade	AF	ADo	RD	RF	RDo	SDR(%)
<i>Alternanthera philoeroides</i> *	G	15	3	20,5	14,29	12	32,87	19,72
<i>Digitaria longiflora</i>	G	4	1	2,6	3,81	4	4,17	3,99
<i>Physalis angulata</i>	B	1	1	0,6	0,95	4	0,96	1,97
<i>Cyperus iria</i>	S	4	2	1,3	3,81	8	2,08	4,63
<i>Amaranthus spinosus</i> *	B	14	3	6	13,33	12	9,62	11,65
<i>Ageratum conyzoides</i>	B	5	1	3,7	4,76	4	5,93	4,90
<i>Ludwigia perrenis</i> *	B	12	3	2,8	11,43	12	4,49	9,31
<i>Eclipta prostrata</i>	B	1	1	0,6	0,95	4	0,96	1,97
<i>Bomera leavis</i>	B	2	1	1,2	1,91	4	1,92	3,27
<i>Eleusine indica</i>	S	5	1	6,1	4,76	4	9,78	6,18
<i>Phyllanthus debilis</i>	B	3	2	0,7	2,86	8	1,12	3,99
<i>Boerhavia erecta</i> *	B	6	1	6,9	5,71	4	11,06	6,93
<i>Portulaca oleraceae</i> *	B	21	2	5,1	20,0	8	8,18	12,06
<i>Euphorbia hirta</i>	B	2	1	0,2	1,91	4	0,32	2,08
<i>Pentella repens</i>	G	8	1	3,9	7,62	4	6,25	5,96
<i>Hedrotis corymbasa</i>	S	2	1	0,1	1,91	4	0,16	2,02
Total		105	25	62,4	99,99	100	99,89	100,6

Notice : ADe (absolute density), AF (absolute frequency), ADo (absolute dominance ), Rde (relative density), RF (relative frequency), and RDo (relative dominance), SDR (summed dominance ratio), \* (five most dominance weeds), G (grass), B (broad leaf), S (sedge)



The ability to survive in non host plants indicated the existence of nutrients content in the weed that resembles its host plant nutrient content or the existence of certain substances that could stimulate *D. citri* to stay for feeding. The substances could be a various compounds of alcohol and aldehyde from the leaves of which were specific and volatile used by insect to find its host plant (Visser, 1986). However, the type and number of nutrient content was not as complete as in the host plant for surviving and completing a life cycle. Similar results were also obtained by Sudiono and Purnomo (2008) in Gemini virus isect vectors (*Bemecia tabaci*), and Hardiastono (2001) on Peanut Stripe Virus (PStV). *B. tabaci* were able to live on broadleaf weed *Ageratum conyzoides*. The weeds also serve as a source of inoculum since they were able to be infected by Gemini virus. While *Amaranthus spinosus*, *Bidens pilosa*, *Crotalaria incana*, *Glycine max* and *Physalis angulata* as alternative hosts and source of inoculums for PStV infection.




Figure 1. Percentage of *D. citri* survived on various dominant weeds

Table 2. The mean of longevity (days) of *D. citri* on various dominant weeds

Weeds species	Mean ± SE
<i>Boerhavia erecta</i>	5,91 ± 0,251 a
<i>Amaranthus spinosus</i>	3,57 ± 0,292 b
<i>Althernanthera philoxeroides</i>	0,20 ± 0,200 c
<i>Portulaca oleracea</i>	0,20 ± 0,200 c
<i>Ludwigia perrenis</i>	0,20 ± 0,200 c

Numbers followed by the same letter are not significantly different between treatments ( $P < 0,001$ )

Some of *D. citri* population (31%), obviously did not prefer to stay for feeding on any weeds treatments and just stayed on the cage. Presumably they did not recognize these five kinds of weeds as a host that can be used as a food source. While the absence of *D. citri* population on *P. Oleracea* likely could be caused by high water content in this weeds, which were structurally quite different from their host plants (citrus). The high water content also causes the low content of volatile compounds (specific odors) which were emitted, and were unable to attract *D. citri* stayed for feeding.

The implications of those survivorships of *D. citri* in some weed species is the presence of alternative host for *D. citri* when citrus plant is not available as a food source, either because there are no plants or when plants are treated with pesticides. The existence of alternative host will cause the population of *D. citri* is available throughout the season and serve as initial population for the next generation population. Role of *D. citri* as a propagative vector of CVPD pathogen will also make the disease inoculums always available throughout the season in field. This will further complicate the control measures against CVPD on citrus plant, since the disease transmission is highly dependent on the availability of disease inoculums and insect vector population in the field (Chen, 1998).

## CONCLUSION

1. Weeds were able as an alternative host for *D. citri*.
2. The mean of longevity of *D. citri* on broad leaf weeds *Bhoeravia erecta* was 5.91  $\pm$  0.251 days.

## REFERENCES

- Aubert B. 1990. Integrated activities for the control of *Huanglongbing*-greening and its vector *Diaphorina citri* Kuwayama in Asia. In: Aubert B, Tontyaporn S, Buangsuwon D (eds). Proceedings of the Fourth International Asia Pacific Conference on Citrus Rehabilitation, Chiang Mai, Thailand, 4-10 February 1990. Rome: FAO UNDP. pp133-144.
- BPS. 2009. Produksi buah-buahan di Indonesia. [http://www.bps.go.id/tab\\_sub/view.php?tabel=1&daftar=1&id\\_subyek=55&notab=2](http://www.bps.go.id/tab_sub/view.php?tabel=1&daftar=1&id_subyek=55&notab=2)
- Chen CN. 1998. Ecology of the insect vectors of citrus systemic diseases and their control in Taiwan. Food & fertilizer technology center. [www.agnet.org/library/data/eb/eb459a/eb459a.pdf](http://www.agnet.org/library/data/eb/eb459a/eb459a.pdf)
- Gomez KA, Gomez AA. 1983. Statistical procedures for agricultural research. Second edition. John Wiley and Sons. New York. 680 pp.



- Halbert SE, Manjunath KL. 2004. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87: 330-353.
- Hardiastono, T. 2001. Potensi gulma berdaun lebar pada pertanian kacang tanah (*Arachis hypogaea* L.) sebagai inang alternative penyakit bilur peanut stripe virus (PStV). *Habitat*. 12 (3): 139-146
- Irawan IGP, Sulistyowati L, Wijaya IN. 2003. Penyakit CVPD Pada Tanaman Jeruk (Analisis Baru Berbasis Bioteknologi). *Dirjen Perlindungan Hortikultura*.
- Rukmana dan Saputra. 1999. *Gulma dan Teknik Pengendalian*. Kanisius. 88 pp.
- SPSS Inc. 1999. *SPSS® for Windows™ Version 10.0.5*. Chicago: SPSS Inc.
- Sudiono dan Purnomo. 2008. Studi kisaran inang kutu kebul (*Bemisia tabaci* Genn.) di sentra sayuran dataran tinggi tanggamus. *Jurnal Penelitian Terapan*. 8 (3): 103-108.
- Visser, JH. 1986. Host odor perception in herbivores insects. *Annual Reviews of Entomology*. 31, 121-144
- Xu CF, Xia YH, Li KB, Ke C. 1988. Further study of the Transmission of citrus *Huanglongbing* by a psyllid *Diaphorina citri* kuwayana. *Proceeding of 10<sup>th</sup> Conference of IOCV*. Riverside. pp.243-248