

Sulfated exopolysaccharide
production and nutrient removal
by the marine diatom
Phaeodactylum tricorneratum growing
on palm oil mill effluent
by Muhamad Maulana Azimatun Nur

Submission date: 15-Feb-2021 11:04AM (UTC+0700)

Submission ID: 1509693739

File name: Nur2019_Article_SulfatedExopolysaccharideProdu_1.pdf (1.18M)

Word count: 9319

Character count: 45522



Sulfated exopolysaccharide production and nutrient removal by the marine diatom *Phaeodactylum tricornerutum* growing on palm oil mill effluent

Muhamad Maulana Azimatun Nur^{1,2} · Manasveni Kilnagar Swaminathan² · P. Boelen² · A. G. J. Buma²

Received: 5 July 2018 / Revised and accepted: 4 March 2019 / Published online: 16 March 2019
© The Author(s) 2019

Abstract

Palm oil mill effluent (POME) is the major source of environmental hazard in palm oil industry. Yet, due to its high nutrient content, POME may offer opportunities for the growth of algae as a source of value-added compounds such as sulfated extracellular polysaccharide (sEPS) while simultaneously removing valuable nutrients such as phosphate. The aim of this paper was to evaluate growth, total sEPS production, and nutrient removal by the diatom *Phaeodactylum tricornerutum* grown on POME under a range of experimental conditions (temperature, salinity, supplementation of extra nutrients). *Phaeodactylum tricornerutum* was found to grow well on a range of POME concentrations, with 30% POME as optimum concentration. Nitrate and urea addition enhanced both growth rate and final biomass, whereas phosphate significantly stimulated growth only at low temperature. Box-Behnken response surface methodology revealed that interactions between temperature and salinity, and between temperature and urea influenced sEPS production. The highest total sEPS (140 mg L⁻¹) concentration was recorded at 25 °C, 2.6% salinity, and 100 mg L⁻¹ urea addition. Our study shows that POME wastewater, supplemented with urea at relatively high temperatures, can be considered as a potential medium for *P. tricornerutum* to replace commercial nutrients while producing high amounts of sEPS and removing almost 90% of phosphorous from the wastewater.

Keywords POME wastewater · Temperature · Extracellular polysaccharide · *Phaeodactylum tricornerutum* · Box-Behnken · sEPS

Introduction

Palm oil mill effluent (POME) is agricultural wastewater generated from palm oil processing. Presently, Asian coconut palm oil (CPO) production comprises 87.8% of the total world production of 57.32 million t. Within this region, Indonesia is currently known as the largest CPO producer, followed by Malaysia (Faostat 2018). Direct discharge of POME into the environment causes severe pollution due to its high organic

matter content, high chemical oxygen demand (COD), biochemical oxygen demand (BOD), high nutrient levels, dark color, and strong smell (Nur et al. 2017). To date, the conventional ponding system is the most common method for treating POME (Liew et al. 2014). In Malaysia, more than 85% of palm oil mills have adopted the ponding system due to its low capital cost (Tong and Jaafar 2004). Nevertheless, the conventional ponding system has various limitations such as low nutrient removal efficiency, long retention times, and enhanced greenhouse gas emissions (Tabassum et al. 2015). The strategy to decrease adverse characteristics of POME by using microalgae while simultaneously producing value added products could be promising due to the fact that POME contains high micro- and macronutrient levels (Nur and Buma 2018). However, the nutrient removal efficiency by microalgae will also depend on cultivation conditions as well as on nutrient composition in terms of speciation and initial N/P ratio (Beuckels et al. 2015). Given the worldwide phosphorus shortage, nutrient conditions within cultivation systems should be optimal or otherwise optimized such that complete phosphate removal from the wastewater is achieved.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10811-019-01780-2>) contains supplementary material, which is available to authorized users.

✉ Muhamad Maulana Azimatun Nur
m.m.azimatun.nur@rug.nl; lanaazim@upnyk.ac.id

¹ Department of Chemical Engineering, Faculty of Industrial Technology, Universitas Pembangunan Nasional “Veteran”, Yogyakarta, Indonesia

² Department of Ocean Ecosystems, Energy and Sustainability Research Institute, Faculty of Science and Engineering, University of Groningen, Groningen, The Netherlands

Extracellular polysaccharide or exopolysaccharide (EPS) is a group of polysaccharide substances generated from secondary metabolic processes within algae, excreted under normal as well as under unfavorable conditions (De Jesus Raposo et al. 2013; Ates 2015). EPS may contain xylose, galactose, or glucose. In addition, sulfated exopolysaccharides (sEPS) are formed, which may be attached to lipids or proteins, depending on the cultivated algal species (Delattre et al. 2016). In general, the overall composition of sEPS is species specific. For example, sEPS produced by *P. tricornutum* contains 7.5–13.3% sulfate, and 1.4–6.3% uronic acids, while sEPS from *Chlorella stigmatophora* contains 7.8–9.4% sulfate and 3.7–9.0 uronic acid (Guzmán et al. 2003).

Recently, the interest in utilizing EPS from microalgae has increased. Several applications have been widely implemented in agricultural fields to improve soil properties, and thereby to enhance plant growth (Painter 1993). In the pharmaceutical field, sEPS from *P. tricornutum* is promising as an anti-inflammatory, antiviral, antiparasitic, anti-tumor, and hypocholesterolemic compound (Guzmán et al. 2003; De Jesus Raposo et al. 2013; Delattre et al. 2016). However, the production of microbial sEPS at a large scale, especially from microalgae, is limited due to the high production cost (Ates 2015). Alternatively, the cultivation of microalgae on POME with the aim to produce sEPS and carbohydrates could be promising, given the relatively high nutrient content of the wastewater, thereby making commercial nutrients redundant. Furthermore, the treated sEPS containing POME might be utilized for soil improvement in the vicinity of the palm oil plants.

It has been suggested that culture conditions and nutrient composition of the medium influence the sEPS production of microalgae (Guzmán-Murillo et al. 2007; Ekelhof and Melkonian 2017). Microalgae tend to produce higher sEPS under stressed conditions such as excess irradiance and supra-optimal temperatures, as a means to prevent cell damage via the action of exopolymers (Kumar et al. 2007; Delattre et al. 2016). Other factors such as nutritional conditions and salinity were also reported to influence EPS production and composition (Abdullahi et al. 2006; Díaz Bayona and Garcés 2014). However, based on our knowledge, the influence of temperature on EPS production was not yet explored. Furthermore, the interaction between these parameters was not well reported so far. Therefore, it can be assumed that total biomass, total sEPS, and intracellular carbohydrate productivity by *P. tricornutum* cultivated on POME would be influenced by environmental conditions such as salinity, nutrient availability and temperature. The aim of this research was to study the utilization of nutrient enriched POME and the interaction of urea addition, salinity, and temperature on growth, nutrient removal, sEPS, and biomass production of *P. tricornutum*. We chose urea addition because it guarantees complete phosphorus removal (Nur et al. 2019). Furthermore, excess nutrients from the wastewater that cause eutrophication could be treated by using microalgae

since it is cost-effective compared to chemical wastewater treatment (Smith et al. 1998; Delrue et al. 2016). In order to find optimal cultivation conditions, we employed response surface methodology (RSM) based from Box-Behnken design (BBD).

We chose the marine diatom *P. tricornutum* as test alga since *P. tricornutum* was shown previously to grow on POME (Nur and Buma 2018). Furthermore, sEPS composition of *P. tricornutum*, which mainly consist of glucose, mannose, and galactose, was also affected by environmental conditions such as salinity and nutrient addition (Abdullahi et al. 2006). Finally, in contrast to most diatoms, *P. tricornutum* has a cell wall that is rich in sulfated polysaccharides and poor in silica (Le Costaouëc et al. 2017). Therefore, *P. tricornutum* could be a promising candidate as a source of sEPS, when growing on POME in an optimum condition.

Material and methods

Wastewater preparation

POME was obtained from a small factory in Sumatra, Indonesia, after it had been released from an aerobic open pond lagoon. The wastewater was stored at $-19\text{ }^{\circ}\text{C}$ to prevent degradation over a long time period. The wastewater was thawed and filtered with GF/C glass fiber filters (Whatmann, 47 mm) to remove particulate matter. Subsequently, the filtrate was sterilized at $121\text{ }^{\circ}\text{C}$ for 15 min. This treated POME contained 1245 mg L^{-1} COD, 72.4 mg L^{-1} total dissolved nitrogen (TDN), and 7.93 mg L^{-1} total dissolved phosphorus (TDP), as determined using the assay kits LCK349 and LCK138 (Hach Lange).

Experimental setup

Precultures of *P. tricornutum* Bohlin (CCMP2558, NCMA, Maine, USA) were grown on a standard f/2 medium based on filtered natural oligotrophic seawater obtained from NIOZ Netherlands (adjusted to a salinity of 3.5% (w/v) with demineralized water) by the protocol of Guillard (1975) supplemented with silicate ($100\text{ }\mu\text{M}$). The algae were cultivated in a 16:8 h light dark cycle at a temperature of $25\text{ }^{\circ}\text{C}$. The cultures were illuminated with fluorescent lamps (see below) providing $300\text{ }\mu\text{mol photons m}^{-2}\text{ s}^{-1}$. The cultures were diluted with fresh Guillard medium if growth reached the stationary phase. The cultures were acclimated to the experimental conditions for at least 1 week prior to experimentation. In total, four experiments were done.

Determination of the optimal POME concentration for *P. tricornutum* growth

Six concentrations of POME (5–100% v/v) and one control (f/2) were prepared based on appropriate dilutions in sterilized

oligotrophic seawater. The pH of each concentration was adjusted to 8 by using 2 N NaOH or HCl, and the salinity was set to 3.5% (w/v) by using commercially available artificial sea salt (Reef Salt, Aqua Medic, Germany). All media were supplemented with silicate (100 μM). Two replicates (each 75 mL, in 100 mL Erlenmeyer flasks) were made for each concentration. *Phaeodactylum tricornutum* (2% v/v, which equals an initial cell density of around 15×10^4 cells mL^{-1}) was inoculated into each flask, and the cultures were incubated in a water bath at 25 °C and a light/dark cycle of 16 h:8 h. The water bath was equipped with a temperature controller and a U shaped lamp that contained 12 fluorescent lamps (six Biolux and six skywhite lamps, Osram) coupled to reflectors (Doublelux) and connected to dimmers (Osram) and set as described by Van de Poll et al. (2007). The culture flask was closed with a cotton stopper and gently shaken manually every day to resuspend the cells and to provide aeration to the cells. For all experiments, a saturating light intensity (300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) was used as determined earlier (Nur et al. 2019). Due to the variable irradiance penetration in the culture flasks as a result of the different POME fractions, the light intensity was adjusted using neutral density screens. The light intensity was measured in the center of culture flask by using a spherical light sensor (Biospherical Instruments QSL2101, USA) which is small enough to be placed inside the culture flasks. Every 24 h, 1 mL of sample was taken from each culture flask and direct cell counting was performed using a hemocytometer under a light microscope. At least 200 cells were counted using a hand counter. The cultivation was stopped at the end of the exponential phase (4–7 days).

Effects of different nutrient additions on *P. tricornutum* biomass and sEPS production when growing on 30% POME

Phaeodactylum tricornutum was cultured in 75 mL working volume in 100-mL sterilized Erlenmeyer flasks and placed in a water bath as described above. About 2% (v/v) of *P. tricornutum* culture was used as inoculate to autoclaved and filtered medium consisting of 30% v/v POME + 70% v/v filtered natural oligotrophic seawater. Cultures growing on 30% POME alone (no nutrient additions) served as controls. Nitrogen and phosphorus were added to the media to generate different N/P molar ratios (Table 2). All media were supplemented with silicate (100 μM). The experiments were carried out at 20 and 25 °C, initial pH was adjusted to 8.0 ± 0.2 by using 2 N HCl or 2 N NaOH, and salinity was adjusted to 3.5% w/v using artificial sea salt (Reef Salt, Aqua Medic, Germany). The light intensity was set to 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ inside the cultivation medium. Every 24 h, a 1 mL of sample was taken from each culture flask and direct cell counting was performed using a hemocytometer under a light microscope. At the end of the exponential growth phase

(7–15 days cultivation), the cultures were harvested for sEPS, biomass (dry weight), and intracellular carbohydrate.

Effect of salinity, temperature, and urea addition on growth rate, biomass, and sEPS production of *P. tricornutum* growing on 30% POME

General full factorial design (GFFD) with three variables was performed to determine the factor most affecting sEPS and biomass production as well as their interactions with respect to the growth rate. A further experiment based on FFD was employed using Box-Behnken design (BBD) response surface methodology (RSM) to study the optimum value of the variables for biomass and sEPS production. The empirical form of the second order polynomial model for BBD can be described as (Eq. 1)

$$y = \beta_0 + \sum \beta_i x_i + \sum \beta_{ii} x_i^2 + \sum \beta_{ij} x_i x_j \quad (1)$$

where y is the predicted value; and β_0 , β_i , β_{ii} , and β_{ij} are the constant, linear, quadratic, and the interaction coefficient, respectively.

The experimental design, based on two replicates, consisted of three factors, temperature (20, 22.5, 25 °C), salinity (0.5, 2.0, 3.5% w/v), and urea addition (0, 50, 100 mg L^{-1}). The alga was cultured in 75 mL working volume in a 100-mL sterilized Erlenmeyer flask and placed in the water bath equipped with U shaped lamps and temperature controller as described above. About 2% (v/v) of *P. tricornutum* culture was used as inoculate to the medium consisting of 30% v/v POME + 70% v/v seawater. Salinity was adjusted by using seawater and Milli-Q water. All media were supplemented with silicate (100 μM), and pH was adjusted to 8 ± 0.2 by using 2 N HCl or 2 N NaOH at the beginning of the cultivation. The light intensity was set to 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ inside the cultivation medium. About 1 mL of the sample was taken daily and preserved using 1% v/v Lugols iodine. The cell density was counted by using a hemocytometer until the cultures reached the end of the exponential phase (7 days). Then, cultures were harvested for analyses of final biomass (dry weight) and sEPS.

Effect of nutrient enrichment on nutrient removal efficiency of POME

Phaeodactylum tricornutum was cultured in 75-mL working volume in 100-mL sterilized Erlenmeyer flasks and placed in a water bath as described above. About 2% (v/v) of *P. tricornutum* culture was used as inoculate to autoclaved and filtered medium consisting of 30% v/v POME + 70% v/v filtered natural oligotrophic seawater. Cultures growing on 30% POME alone (no nutrient additions) served as control. Different nutrient enrichments were employed, 50 mg L^{-1} of

urea (U), and 3 mg L⁻¹ of phosphate (P). All media were supplemented with silicate (100 μM). The experiments were carried out at 20 °C, initial pH was adjusted to 8.0 ± 0.2 by using 2 N HCl or 2 N NaOH, and salinity was adjusted to 3.5‰ (w/v) using artificial sea salt (Reef Salt, Aqua Medic, Germany). The light intensity was set to 300 μmol photons m⁻² s⁻¹ inside the cultivation medium. At the beginning of cultivation and the end of the exponential growth phase (7–9 days), the culture was filtered using GF/F glassfiber filters and nutrients were analyzed in the filtrate to obtain nutrient removal efficiency (for analysis, see description below).

Analyses

Growth rate

The growth rate was calculated from the linear regression of the natural logarithm of cell density versus time (Eq. 2)

$$\mu = \frac{\ln\left(\frac{X_2}{X_1}\right)}{t_2 - t_1} \quad (2)$$

where μ is the growth rate (day⁻¹), and X_2 and X_1 are the cell densities (cells mL⁻¹) at time 2 (t_2 , day) and time 1 (t_1 , day) respectively.

Biomass based on dry weight

Determination of cell dry weight was done gravimetrically. About 30 mL of the sample was harvested by filtering over pre-dried and pre-weighed GF/C filters. The filters were washed with 0.5 M NH₃HCO₃ according to Zhu and Lee (1997). Then, the filter was dried at 95 °C until a constant weight was reached.

Carbohydrate and sEPS determination

To analyze the intracellular carbohydrate content of *P. tricornutum*, 1 mL of the sample was centrifuged at 13,000 rpm for 5 min. The supernatant was kept for sEPS determination. The pellet was washed twice by using cold Milli-Q water to remove remaining supernatant. The collected pellet was stored at -20 °C until later analysis. Carbohydrate extraction was carried out by using 1 mL double distilled water (Milli-Q) at 80 °C for 1 h (Van Oijen et al. 2005). The sample was then centrifuged at 5000 rpm for 5 min at 22 °C. The supernatant was collected to analyze total carbohydrates, using the total carbohydrate assay kit (Sigma, MAK104-1KT) at 490 nm on a Multilabel counter (Victor 1420, PerkinElmer) according to the Dubois method (Dubois et al. 1956). D-Glucose was used as a standard curve.

The determination of sEPS was done by colorimetry as explained by Ramus (1977) and Guzmán-Murillo et al. (2007) with slight modifications. Briefly, 1 mL of cell-free culture supernatant was gently mixed with 4 mL 0.5 M acetic acid and 500 μL Alcian Blue 8GX (1 mg mL⁻¹ in acetic acid 0.5M, pH 2.5), vortexed at 3000 rpm for 20 s, and then incubated overnight at room temperature (22 °C). The samples were centrifuged at 3000×g for 30 min at 22 °C. The sample was read at 610 nm on a Hach DR 3900, which did not interfere with the maximum peak of wastewater absorption at 320–330 nm. The difference in absorbance between the blank medium (30% v/v POME or Milli-Q water) and the samples was taken as proportional polyanion concentration. Dextran sulfate was used as a standard curve.

The volumetric carbohydrate productivity was calculated based on Eq. 3.

$$P_c = \frac{(X_t - X_0) \cdot C_c}{t} \quad (3)$$

where P_c is the carbohydrate productivity (mg L⁻¹ day⁻¹), X_t is the final biomass concentration (mg L⁻¹), X_0 is the initial biomass concentration (mg L⁻¹), C_c is the carbohydrate content per total biomass dry weight (w/w), and t is the total duration of the cultivation (days).

Nutrient analysis

For nutrient analysis, about 5–10 mL of the sample were filtered using GF/F glassfiber filter (Whatman), after which the filtrate was used for dissolved nutrient analyses as described below. Total phosphorus and orthophosphate were analyzed using the assay kit LCK349 (0.05–1.50 mg L⁻¹ orthophosphate, 0.15–4.50 mg L⁻¹ TDP, provided by Hach, The Netherlands). Samples were diluted using Milli-Q water if necessary. For TDP analysis, 2 mL of the sample was carefully pipetted to the tube containing reagent LCK394 and hydrolyzed to a temperature block (Hach LT 200) at 100 °C for 60 min. After cooling down to room temperature, the tube was shaken vigorously and 0.2 mL of reagent was added. The orthophosphate determination procedure was identical to TDP analysis except for the hydrolysis step. For total dissolved nitrogen analysis, samples were diluted with Milli-Q water if necessary due to the range of TDN provided by the assay kit (1–16 mg L⁻¹) from LCK138 (Hach, The Netherlands). Briefly, 1.3 mL of the sample and reagents was added to a reaction tube and hydrolyzed at 120 °C for 30 min in the heater block. After cooling down, the kit reagent was added to the tube and shaken. Then, 0.5 mL of the sample was pipetted in a tube containing kit reagents (LCK138). For nitrate analysis, LCK339 (Hach) was used (0.23–13.5 mg L⁻¹ NO₃-N) and the sample was diluted with Milli-Q if necessary. About 1 mL of the sample was pipetted to the tube containing

Table 1 Growth rate of *P. tricornutum* cultivated on f/2 and on the mixture of seawater (SW) and filtered POME concentrations at 25 °C and 300 μmol photons m⁻² s⁻¹ (measured inside the culture). Average values of duplicate cultures are shown. SD is shown after ± symbol. Values that do not share a common lowercase letter in the same column are significantly different (*P* < 0.05)

Media	NO ₃ ⁻ -N (μM)	PO ₄ -P (μM)	N/P molar ratio ^a	Growth rate (day ⁻¹)
f/2	880.00	36.90	24:1	1.10 ± 0.04a
5% POME +95% SW	36.79	2.89	13:1	0.30 ± 0.06b
10% POME +90% SW	73.57	5.78	13:1	0.48 ± 0.02bc
30% POME +70% SW	220.71	17.33	13:1	0.66 ± 0.12c
50% POME +50% SW	367.86	28.88	13:1	0.63 ± 0.05c
70% POME +30% SW	515.00	40.44	13:1	0.61 ± 0.01c
100% POME	735.71	57.77	13:1	0.61 ± 0.01c

^aThe molar ratio is calculated from dissolved inorganic nitrogen as nitrate and dissolved phosphorus as phosphate in the media

reagents (LCK339) after which 0.2 mL of kit reagent was added. After 15 min, the sample was measured using spectrophotometry. Measurements were done using a Hach Lange 3900 DR spectrophotometer following the programs listed in the machine.

The efficiency of nutrient removal was calculated based on Eq. 4.


$$E = \frac{n_0 - n_t}{n_0} \times 100 \tag{4}$$

where *E* is the nutrient removal efficiency (%), *n*₀ is the nutrient concentration (mg L⁻¹) of the sample at the beginning of cultivation, and *n*_t is the nutrient concentration (mg L⁻¹) of the sample at the end of cultivation.

Statistical analysis

Minitab ver. 18. (Demo version) was used for statistical analysis, GFFD and BBD design, and the evaluation. Differences between treatments were analyzed with one-way or two-way analysis of variance (ANOVA) with *P* value of 0.05. Post hoc tests (Tukey HSD) were performed for pair-wise comparisons. The experimental results were obtained based on at least two replicates as expressed in the standard deviations (± SD).

Fig. 1 Growth profile of *P. tricornutum* cultivated at 25 °C and 300 μmol photons m⁻² s⁻¹ under different POME concentration



Results

Effect of POME fraction on the growth rate of *P. tricornutum*

The highest growth rate (1.10 day⁻¹) was recorded for *P. tricornutum* growing on f/2 medium, i.e., artificial medium without POME. All POME fractions showed substantially lower growth rates, ranging between 0.30 and 0.66 day⁻¹ (Table 1, Fig. 1). Maximum growth was observed at 30% POME: Above this fraction, growth rates decreased slightly. Therefore, 30% v/v of POME was used for all subsequent experiments.

Effects of nutrient enrichment on growth rate, total biomass, carbohydrate productivity, and sEPS


The highest temperature (25 °C) resulted in overall higher growth rates as compared with the 20 °C cultures (*P* < 0.05) (Table 2). At both temperatures, the growth rates after nutrient enrichment (both nitrogen and phosphorus) on 30% POME were not significantly different compared to f/2 medium. The lowest growth rate was found in the control cultures based on 30% POME without nutrient addition (C) for both temperatures (Table 2) (*P* < 0.05). At 25 °C, POME with urea addition

Table 2 Growth rate, total biomass, total sEPS, carbohydrate productivity, and carbohydrate content of *P. tricornutum* cultivated at 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, on f/2 only and on 30% POME supplemented with different nutrients. Mean values are from two replicates. SD are shown after the \pm symbol. F2, f/2 medium no POME; NPT, POME, nitrate, phosphate, and trace elements; NP, POME, nitrate, and phosphate; UP, POME, urea, and phosphate; U, POME and urea; C, POME only. Values that do not share a common lowercase letter in the same column are significantly different ($P < 0.05$)

Medium	Temp ($^{\circ}\text{C}$)	External nitrogen source	TDIN (μM)	$\text{PO}_4\text{-P}$ (μM)	N/P molar ratio ^a	μ (day^{-1})	Total biomass (g L^{-1})	Carbohydrate productivity ($\text{mg L}^{-1} \text{day}^{-1}$)	Carbohydrate content (%)	Total sEPS production (mg L^{-1})	Total sEPS normalized to biomass (g g^{-1})
F2	20	$\text{NO}_3\text{-N}$	880.00	36.90	24:1	0.84a \pm 0.04	0.35a \pm 0.06	5.45a \pm 0.13	24.06a \pm 3.88	78.89a \pm 11.21	0.23a \pm 0.01
NPT	20	$\text{NO}_3\text{-N}$	435.00	101.67	4:1	0.87a \pm 0.01	0.25a \pm 0.04	5.60ab \pm 1.65	32.66a \pm 5.07	56.06b \pm 0.50	0.22a \pm 0.03
NP	20	$\text{NO}_3\text{-N}$	435.00	101.67	4:1	0.88a \pm 0.02	0.27 a \pm 0.00	5.15a \pm 0.17	28.32a \pm 0.64	61.56b \pm 2.59	0.23a \pm 0.01
UP	20	Urea	435.00	101.67	4:1	0.86a \pm 0.01	0.21a \pm 0.10	5.48a \pm 0.69	41.86a \pm 14.61	59.97b \pm 3.68	0.31a \pm 0.13
U	20	Urea	1863.57	17.33	108:1	0.69b \pm 0.04	0.18ab \pm 0.05	8.63bcd \pm 1.19	34.38a \pm 4.37	71.94ab \pm 2.68	0.43ab \pm 0.09
C	20	–	220.71	17.33	13:1	0.29c \pm 0.04	0.10b \pm 0.01	5.87abc \pm 1.80	41.90a \pm 0.24	78.93a \pm 6.44	0.83c \pm 0.23
F2	25	$\text{NO}_3\text{-N}$	880.00	36.90	24:1	1.10d \pm 0.04	0.29a \pm 0.02	9.41d \pm 0.10	48.49b \pm 2.17	101.25cd \pm 7.03	0.35d \pm 0.04
NPT	25	$\text{NO}_3\text{-N}$	435.00	101.67	4:1	1.09de \pm 0.06	0.31a \pm 0.09	9.53d \pm 0.07	46.61b \pm 12.75	94.64d \pm 1.00	0.31d \pm 0.09
NP	25	$\text{NO}_3\text{-N}$	435.00	101.67	4:1	1.07de \pm 0.09	0.28a \pm 0.07	8.82 cd \pm 0.13	49.42b \pm 13.51	96.89d \pm 4.18	0.36d \pm 0.08
UP	25	Urea	435.00	101.67	4:1	1.02de \pm 0.00	0.26a \pm 0.00	8.03abcd \pm 0.33	46.81b \pm 1.41	95.65d \pm 8.12	0.37d \pm 0.03
U	25	Urea	1863.57	17.33	108:1	0.88e \pm 0.01	0.17ab \pm 0.03	9.17d \pm 1.00	38.95b \pm 2.15	131.18c \pm 0.75	0.80ef \pm 0.13
C	25	–	220.71	17.33	13:1	0.41f \pm 0.07	0.11b \pm 0.04	7.43abcd \pm 0.07	52.68b \pm 22.07	100.52cd \pm 29.20	0.96f \pm 0.12

^a The ratio is calculated from the molar of total dissolved inorganic nitrogen (TDIN) from nitrate or urea and dissolved phosphorus as phosphate from external nutrient addition and/or POME

Fig. 2 Pareto chart showing the effects of parameters on the growth rate of *P. tricornutum*. The vertical line indicates the significance of the effects at 95% confidence level. A is temperature (°C), B is salinity (%), and C is urea addition (mg L⁻¹)



(U) alone (no phosphate addition) resulted in a growth rate averaging $0.88 \pm 0.04 \text{ day}^{-1}$, which was significantly higher compared to C ($0.41 \pm 0.07 \text{ day}^{-1}$). Furthermore, the growth rate at 25 °C was not significantly different between urea addition alone and nitrate/urea plus phosphate addition (NPT, NP, UP) ($P > 0.05$). However, at 20 °C, the growth rate after urea addition alone was significantly different compared to f/2, NPT, NP, and UP medium ($P < 0.05$). Based on this case, the addition of phosphorus alone could also influence growth rate at low temperature. In our research, the addition of 1 mg L⁻¹ phosphate alone to POME resulted at 20 °C in a growth rate as high as the urea addition alone (data not shown).

With respect to total dry biomass production, the addition of nitrogen and phosphorus resulted in significantly different values compared to the POME control C at both temperatures ($P > 0.05$) (Table 2). Furthermore, the total biomass production after the addition of urea alone (U) was not significantly different compared to NPT, NP, and UP at both temperatures ($P > 0.05$). Also, no significant difference was found between both temperatures (Table 2) ($P > 0.05$).

Temperature significantly influenced carbohydrate content and total sEPS production ($P < 0.05$), while nutrient enrichment on 30% POME did not significantly influence total sEPS production at both temperatures (Table 2). Total sEPS normalized to biomass was found to be significantly higher with the addition of urea-only (U) compared to combined nutrient additions (NPT, NP, UP) at 25 °C. Furthermore, the addition of urea-only (U) resulted in no significantly different carbohydrate productivity compared to the other nutrient additions (NPT, NP, UP) at the high temperature ($P < 0.05$), although a trend was observed of elevated sEPS levels for the control and urea-only additions.

Since the urea-only conditions (U) showed positive results (growth, biomass, and sEPS higher than other nutrient additions) as compared with the 30% POME alone cultures, this

addition was chosen for the last experimental series. In this way, phosphate addition was avoided.

Optimization of total biomass and total sEPS as a function of salinity, temperature, and urea addition

Based on the FFD statistical analysis, all interactive variables influenced the growth rate of *P. tricornutum* ($P < 0.05$) (Fig. 2). Increased salinity or urea concentration enhanced the growth rate, and the interactive effect of temperature-urea and temperature-salinity also enhanced the growth rate (Fig. 3). Based on the Pareto chart (Fig. 2), the interactive effect of temperature, salinity, and urea was the most influencing factor with respect to the growth rate, followed by urea addition and salinity. Therefore, temperature, urea, and salinity were chosen for the optimization of the total biomass and total sEPS based on BBM RSM (Table 3).

BBD showed that higher temperatures increased total sEPS ($P < 0.05$), while the addition of urea alone had no effect on total sEPS ($P > 0.05$) (Table 4, Fig. 4, S1). The optimal salinity was recorded at 2.0–2.5% (w/v) ($P < 0.05$). The interactive effect of temperature-salinity and the interactive effect of temperature-urea influenced the total sEPS ($P < 0.05$) (Table 4, Fig. 4, S1).


As for total biomass, urea levels exceeding 40–80 mg L⁻¹, temperatures above 22–23 °C, and a salinity exceeding 2.0–3.5% (w/v) lowered total biomass ($P < 0.05$) (Fig. 5). Furthermore, the interactive effect between variables did not significantly influence biomass ($P > 0.05$).

Nutrient removal efficiency


TDP and TDN removal efficiency were highly dependent on the nutrient enrichment used (Fig. 6). The highest TDP removal

Fig. 3 Response surface plot (3D) of growth rate as a function of temperature, salinity, and urea addition at hold value **a)** 50 mg L⁻¹ urea addition, **b)** 3.5% (w/v) salinity, and **c)** temperature 22.5 °C


a)



b)



c)



efficiency ($88.2 \pm 2.3\%$) was recorded for POME supplemented with urea (U) alone (no phosphate addition). This value was

significantly different from the control C (POME without nutrient $68.8 \pm 3.07\%$) and from the P (phosphate addition) (54.3

Table 3 Design of the experiment of BBD with the results. Mean ± SD values of the experimental results are from two replicates

Temperature (°C)	Salinity (%)	Urea addition (mg L ⁻¹)	Total biomass (g L ⁻¹)		Total sEPS (mg L ⁻¹)	
			Experimental	Predicted	Experimental	Predicted
20	0.5	50	0.06 ± 0.00	0.07	43.48 ± 1.14	48.18
20	3.5	50	0.18 ± 0.05	0.18	71.94 ± 2.68	62.44
20	2.0	0	0.08 ± 0.01	0.11	96.88 ± 8.92	103.87
20	2.0	100	0.13 ± 0.00	0.13	87.17 ± 0.69	84.98
22.5	0.5	0	0.08 ± 0.01	0.02	41.18 ± 1.26	30.68
22.5	3.5	0	0.12 ± 0.00	0.14	87.33 ± 1.17	85.44
22.5	0.5	100	0.11 ± 0.01	0.07	50.41 ± 2.95	46.40
22.5	3.5	100	0.19 ± 0.02	0.17	65.44 ± 0.08	78.38
22.5	2.0	50	0.24 ± 0.11	0.27	99.31 ± 3.43	101.65
25	0.5	50	0.02 ± 0.00	0.02	33.74 ± 1.68	46.83
25	3.5	50	0.17 ± 0.03	0.14	131.71 ± 0.75	119.31
25	2.0	0	0.05 ± 0.02	0.05	94.45 ± 26.77	108.41
25	2.0	100	0.12 ± 0.01	0.11	141.06 ± 0.69	135.97

± 5.1%) (*P* < 0.05). For TDN removal efficiency, the addition of nutrients was nowhere significantly different compared to the control (*P* > 0.05). The highest TDN removal was found for POME enriched with urea (45.2 ± 3.1%), followed by the control (18.4 ± 12.2%), and P addition (13.7 ± 11.4%).

Discussion

The present study shows that optimal growth of *P. tricornutum* is achieved when growing on 30% POME supplemented with urea. Without nutrient addition, the observed higher growth rate at 30% POME (v/v) as compared with 5 and 10% POME (v/v) could be due to the relatively higher nutrient concentrations in the 30% POME cultures (Table 1). As reported previously, POME contains high levels of micro nutrients such as iron, zinc, potassium, and magnesium, which could boost algal growth (Habib et al. 2003). Furthermore, POME may also contain acetic acid which could be utilized by *P. tricornutum* by applying mixotrophic growth (Villanova et al. 2017; Nur and Buma 2018).

In contrast, at the highest POME fractions (> 30% v/v), the slightly lower growth rates could not be explained by lower light penetration due to high turbidity in POME, as irradiance was set to saturating levels in all cultures (300 μmol photons m⁻² s⁻¹) (Nur et al. 2019). Instead, the presence of potentially toxic phenolic compounds could provide a valid explanation, since these are still contained in digested POME (Chantho et al. 2016). The total concentration of phenolic compounds in POME can reach up to 303 mg L⁻¹ as demonstrated before (Neoh et al. 2013), yet unfortunately these compounds were not measured during our experiments. Marine microalgae such as *Dunaliella salina* and *P. tricornutum* were described to be more sensitive to phenolic compounds from


digested POME than freshwater algae such as *Chlorella vulgaris* (Nur et al. 2016; Duan et al. 2017). Overall, a 30% POME fraction was found to be most suitable to be used as growth medium for *P. tricornutum* for all subsequent experiments in this study. However, the optimal fraction is determined by the quality and the nutritional content of POME (COD, BOD, nitrogen, phosphorus, lignin) which may vary depending on factories and seasonal crops that could lead to deviations in these substances of around ± 20–50% (Poh et al. 2010).

Nutrient additions were found to significantly affect growth and final biomass of *P. tricornutum* growing on POME. *P. tricornutum* could grow well on 30% POME (0.66 ± 0.12 day⁻¹) without adding external nutrients, likely due to the favorable N/P ratio being close to Redfield (13:1) as found in a previous study (Nur et al. 2016; Hadiyanto et al. 2017). However, this growth rate was significantly lower than the artificial f/2 medium, which might be due to nutrient speciation

Table 4 Estimated regression uncoded coefficient of Box-Behnken design RSM for total sEPS production (mg L⁻¹)

Variables	Coefficient	<i>P</i> value	Remarks
Constant	777	< 0.01	Significant
Temp (°C)	-63.10	< 0.01	Significant
Salinity (%)	2.5	< 0.01	Significant
Urea (mg L ⁻¹)	-1.85	0.534	
Temp (°C). Temp (°C)	1.25	0.228	
Salinity (%).Salinity (%)	-17.90	< 0.01	Significant
Urea (mg L ⁻¹). Urea (mg L ⁻¹)	-0.00	0.856	
Temp (°C). Salinity (%)	3.88	< 0.05	Significant
Temp (°C).Urea (mg L ⁻¹)	0.09	< 0.05	Significant
Salinity (%).Urea (mg L ⁻¹)	-0.07	0.242	

Fig. 4 Response surface plots (3D) showing the effects of temperature ($^{\circ}\text{C}$), urea (mg L^{-1}), and salinity (%) on total sEPS generated by *P. tricoratum* at hold value 22.5°C (a) 50 mg L^{-1} (b), and 2.0% (c)



issues in the treated POME which contains relatively high organic nitrogen (Onyia et al. 2001). As a result, the addition of nutrients, in particular nitrate/urea and phosphate, significantly enhanced the growth rate at both temperatures, reaching close to $f/2$ levels (Table 2). The addition of nitrogen—in this case urea—alone (U) promoted growth and total biomass to levels similar to those of nitrogen plus phosphate (NPT, NP, UP) at high temperature (Table 2). This implies that the phosphorus availability in POME does not limit the growth of *P. tricoratum*. At the same time, the addition of urea alone was found to guarantee almost complete phosphorus removal (Fig. 6). A similar result was reported earlier; at high temperatures, phosphorus exhaustion did not decrease growth rate, compared to low temperatures where phosphorus limitation inhibits growth, probably due to the fact that cells require a higher amount of nutrients with decreasing temperature (Rhee and Gotham 1981; Delgadillo-Mirquez et al. 2016).

As shown in the present study, urea can be an excellent nitrogen source for *P. tricoratum*, boosting growth rate,

biomass, and phosphate removal when cultivated on POME. Therefore, compared to phosphorus addition, urea addition is more sustainable since the media containing phosphorus released to rivers could enhance eutrophication of water bodies (Poh et al. 2010). Also, the phosphorus source is more expensive than the nitrogen source which was added from cheap fertilizer (urea). However, the addition of urea may not be the best option to remove phosphorus in the large scale since urea will be left over in the medium after harvesting of the algae, when not carefully supplemented. To solve this, the media containing any excess urea could be reused as growth medium for second-stage algal growth, or recycled as palm tree fertilizer, while the sEPS produced in the filtrate could also serve as soil fertilizer (Painter 1993). Besides this, the optimum condition for urea needs to be further explored since high urea concentrations might result in lower growth rate, possibly due to urea toxicity (Collos and Harrison 2014).

Salinity and the interaction between temperature and salinity or urea significantly influenced the growth rate (Table 3).

This outcome is of interest when *P. tricornutum* is considered for large-scale production systems. At 0.5% salinity, *P. tricornutum* showed a lower growth rate compared to 2.0 and 3.5% (w/v). Furthermore, the addition of urea did not promote growth, when salinity was low and temperature was high, possibly due to low salinity stress in *P. tricornutum*. Liang et al. (2014) reported that optimum growth and photosynthetic efficiency for *P. tricornutum* were recorded at 2.0–4.0% (w/v) salinity. This condition was also supported by our finding that the optimum condition for total biomass production was found at 2.0–3.0% (w/v) salinity, 40–80 mg L⁻¹ urea addition, at 22–23 °C (Fig. 5).

Total sEPS production and sEPS content were found to be influenced by temperature (Table 2, Fig. 4). When temperature was relatively low, total sEPS was significantly reduced. An explanation for this could be that *P. tricornutum* accumulates sEPS when the culture is in a stressed condition to protect cells from damage (Delattre et al. 2016). Previous studies using the green alga *Graesiella* revealed that total EPS production increased with temperature (Mezhoud et al. 2014).

In the present study, cultivation at 25 °C also resulted in an elevated carbohydrate content. This matches an earlier study

where carbohydrate content was enhanced when *P. tricornutum* was grown at high irradiance, high temperature, around 27 °C, and low nitrogen availability, reflecting unfavorable conditions (Buono et al. 2016). In our experiments, carbohydrate content and total sEPS content in *P. tricornutum* seemed linked. Previous studies showed that the intracellular carbohydrate production of green algae was in line with the EPS production, as a response to nutrients, toxic compounds, temperature, and light intensity (El-Sheekh et al. 2012; García-Cubero et al. 2018).

Total sEPS production was also influenced by nutrient availability. Although not significant, the addition of nitrogen and phosphate to 30% POME seemed to result in lower total sEPS production compared to control (C) cultures for both temperatures (Table 2). This implies that phosphate limitation could stimulate sEPS production. The availability of nutrients with the addition of nitrogen increased total biomass but decreased total sEPS normalized to biomass (Table 2). At control C (POME without nutrients), low growth rates and total biomass were found, indicating that the cultures were nitrogen limited, which could result in a higher total sEPS normalized to biomass: sEPS levels were 0.83 ± 0.23 g g⁻¹ algal biomass at low

Fig. 5 Response surface plots (3D) showing the effects of temperature (°C), urea (mg L⁻¹), and salinity (%) on total biomass (g L⁻¹) generated by *P. tricornutum* at hold value 50 mg L⁻¹ (a), 22.5 °C (b), and 2.0% (c)





Fig. 6 Nutrient removal by *P. tricornutum* (at 20 °C) growing on 30% POME with phosphate or urea addition. P = 3 mg L⁻¹ phosphate addition, U = 50 mg L⁻¹ urea addition, C = control POME without nutrient enrichment. Asterisk (*) indicates significantly different from other values ($P < 0.05$)



temperature and $0.96 \pm 0.12 \text{ g g}^{-1}$ at high temperature. These values were 2-fold and 1.2-fold higher compared to the urea-only cultures (Table 2). This result was supported by previous studies who found that the total exopolysaccharide (normalized to biomass) increased when *Cylindrotheca closterium* and *Porphyridium marinum* were grown in nitrogen-limited medium (Staats et al. 2000; Soanen et al. 2016).

In this work, the interactive effect of temperature and urea influenced total sEPS production. The addition of urea at high temperature promoted total sEPS production (Fig. 4). It seems that total sEPS at high temperature with the addition of urea was more dependent on the increase in total biomass production rather than total sEPS normalized to biomass. This is in accordance with earlier studies who reported that media enriched with nitrogen resulted in higher total EPS production (Lupi et al. 1994; Guerrini et al. 2000; Magaletti et al. 2004; Ekelhof and Melkonian 2017), although the total sEPS normalized to biomass was slightly lower compared to the control (POME only) (Table 2). Furthermore, the higher total sEPS normalized to biomass as found at the higher temperature (Table 2) might be explained by the interaction of temperature and urea that may become toxic at high temperatures, possibly due to the rapid hydrolysis of urea to ammonium by the cellular enzyme urease (Lomas and Glibert 1999; Glibert et al. 2008; Ni 2014). In support of this, EPS production by algae was increased when toxic compounds were added to the medium to protect cells from damage (El-Sheekh et al. 2012). In summary, in order to obtain optimal conditions for total sEPS production by *P. tricornutum*, cultivation conditions are proposed to be 25 °C, 2.6% salinity, and 100 mg L⁻¹ urea addition. In this way, 140 mg L⁻¹ sEPS can be obtained based on the RSM approach (Fig. 4, Table 4). However, to make sEPS production more sustainable and cheap, the biomass that contained value-added products (fucoxanthin; carbohydrate)

could also be used as the source of fine chemicals and bioethanol by utilizing the bio-refinery concept. In this case, the optimum biomass and sEPS could be produced by the diatom for 7 days cultivation by adding 50 mg L⁻¹ urea on 30% POME. By using wastewater and adding 50 mg L of urea, the cost of fertilizer for the growth could be greatly reduced.

In conclusion, the present study has shown that *P. tricornutum* is a suitable candidate for effective cultivation on 30% POME while producing sEPS at the large scale, when salinity requirements are met at 2.0–3.5% (w/v). To obtain nearly complete phosphorus removal, high growth rates, and total sEPS, 25 °C cultivation temperature needs to be accompanied by urea addition.

Funding information This project is funded by Lembaga Pengelola Dana Pendidikan (LPDP), Kementerian Keuangan, Republik Indonesia (Reference no. PRJ-72/LPDP.3/2016).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abdullahi AS, Underwood GJC, Gretz MR (2006) Extracellular matrix assembly in diatoms (Bacillariophyceae). V. Environmental effects on polysaccharide synthesis in the model diatom, *Phaeodactylum tricornutum*. J Phycol 42:363–378
- Ates O (2015) Systems biology of microbial exopolysaccharides production. Front Bioeng Biotech 3:200

- Beuckels A, Smolders E, Muylaert K (2015) Nitrogen availability influences phosphorus removal in microalgae-based wastewater treatment. *Water Res* 77:98–106
- Buono S, Colucci A, Angelini A, Langellotti AL, Massa M, Martello A, Fogliano V, Dibenedetto A (2016) Productivity and biochemical composition of *Tetrademus obliquus* and *Phaeodactylum tricorutum*: effects of different cultivation approaches. *J Appl Phycol* 28:3179–3192
- Chantho P, Musikavong C, Suttinun O (2016) Removal of phenolic compounds from palm oil mill effluent by thermophilic *Bacillus thermoleovorans* strain A2 and their effect on anaerobic digestion. *Int Biodeterior Biodegrad* 115:293–301
- Collos Y, Harrison PJ (2014) Acclimation and toxicity of high ammonium concentrations to unicellular algae. *Mar Pollut Bull* 80:8–23
- De Jesus Raposo MF, De Morais RMSC, De Morais AMMB (2013) Bioactivity and applications of sulphated polysaccharides from marine microalgae. *Mar Drugs* 11:233–252
- Delattre C, Pierre G, Laroche C, Michaud P (2016) Production, extraction and characterization of microalgal and cyanobacterial exopolysaccharides. *Biotechnol Adv* 34:1159–1179
- Delgadillo-Mirquez L, Lopes F, Taidi B, Pareau D (2016) Nitrogen and phosphate removal from wastewater with a mixed microalgae and bacteria culture. *Biotechnol Rep* 11:18–26
- Delnue F, Álvarez-Díaz PD, Fon-Sing S, Fleury G, Sassi JF (2016) The environmental biorefinery: using microalgae to remediate wastewater, a win-win paradigm. *Energies* 9:132
- Díaz Bayona KC, Garcés LA (2014) Effect of different media on exopolysaccharide and biomass production by the green microalga *Botryococcus braunii*. *J Appl Phycol* 26:2087–2095
- Duan W, Meng F, Lin Y, Wang G (2017) Toxicological effects of phenol on four marine microalgae. *Environ Toxicol Pharmacol* 52:170–176
- Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F (1956) Colorimetric method for determination of sugars and related substances. *Anal Chem* 28:350–356
- Ekelhof A, Melkonian M (2017) Enhanced extracellular polysaccharide production and growth by microalga *Netrium digitus* in a porous substrate bioreactor. *Algal Res* 28:184–191
- El-Sheekh MM, Khairy HM, El-Shenody R (2012) Algal production of extra and intra-cellular polysaccharides as an adaptive response to the toxin crude extract of *Microcystis aeruginosa*. *Iran J Environ Health Sci Eng* 9:10
- Faostat (2018) <http://www.fao.org/faostat/en/#data/QD/visualize>
- García-Cubero R, Cabanelas ITD, Sijtsma L, Kleinegris DMM, Barbosa MJ (2018) Production of exopolysaccharide by *Botryococcus braunii* CICALA 778 under laboratory simulated Mediterranean climate conditions. *Algal Res* 29:330–336
- Glibert PM, Azanza R, Burford M, Furuya K, Abal E, al-Azri A, al-Yamani F, Andersen P, Anderson DM, Beardall J, Berg GM, Brand L, Bronk D, Brookes J, Burkholder JAM, Cembella A, Cochlan WP, Collier JL, Collos Y, Diaz R, Doblin M, Drennen T, Dyhrman S, Fukuyo Y, Furnas M, Galloway J, Granéli E, Ha DV, Hallegraeff G, Harrison J, Harrison PJ, Heil CA, Heimann K, Howarth R, Jauzein C, Kana AA, Kana TM, Kim H, Kudela R, Legrand C, Mallin M, Mulholland M, Murray S, O'Neil J, Pitcher G, Qi Y, Rabalais N, Raine R, Seitzinger S, Salomon PS, Solomon C, Stoecker DK, Usup G, Wilson J, Yin K, Zhou M, Zhu M (2008) Ocean urea fertilization for carbon credits poses high ecological risks. *Mar Pollut Bull* 56:1049–1056
- Guerrini F, Cangini M, Boni L, Trost P, Pistocchi R (2000) Metabolic responses of the diatom *Achnanthes brevipes* (Bacillariophyceae) to nutrient limitation. *J Phycol* 36:882–890
- Guillard RRL (1975) Culture of phytoplankton for feeding marine invertebrates. In: Smith WL, Chanley MH (eds) Culture of marine invertebrate animals. Plenum Press, New York, pp 26–60
- Guzmán S, Gato A, Lamela M, Freire-Garabal M, Calleja JM (2003) Anti-inflammatory and immunomodulatory activities of polysaccharide from *Chlorella stigmatophora* and *Phaeodactylum tricorutum*. *Phytother Res* 17:665–670
- Guzmán-Murillo MA, López-Bolaños CC, Ledesma-Verdejo T, Roldan-Libenson G, Cadena-Roa MA, Ascencio F (2007) Effects of fertilizer-based culture media on the production of exocellular polysaccharides and cellular superoxide dismutase by *Phaeodactylum tricorutum* (Bohlin). *J Appl Phycol* 19:33–41
- Habib MAB, Yusoff FM, Phang SM, Kamarudin MS, Mohamed S (2003) Growth and nutritional values of *Molina micrura* fed on *Chlorella vulgaris* grown in digested palm oil mill effluent. *Asian Fish Sci* 16:107–119
- Hadiyanto H, Soetrisnanto D, Silviana S, Mahdi MZ, Titisari YN (2017) Evaluation of growth and biomass productivity of marine microalga *Nannochloropsis* sp. cultured in palm oil mill effluent (POME). *Philipp J Sci* 146:355–360
- Kumar AS, Mody K, Jha B (2007) Bacterial exopolysaccharides—a perception. *J Basic Microbiol* 47:103–107
- Le Costaouëc T, Unamunzaga C, Mantecon L, Helbert W (2017) New structural insights into the cell-wall polysaccharide of the diatom *Phaeodactylum tricorutum*. *Algal Res* 26:172–179
- Liang Y, Sun M, Tian C, Cao C, Li Z (2014) Effects of salinity stress on the growth and chlorophyll fluorescence of *Phaeodactylum tricorutum* and *Chaetoceros gracilis* (Bacillariophyceae). *Bot Mar* 57:469–476
- Liew WL, Kassim MA, Muda K, Loh SK, Affam AC (2014) Conventional methods and emerging wastewater polishing technologies for palm oil mill effluent treatment: a review. *J Environ Manag* 149:222–235
- Lomas MW, Glibert PM (1999) Interactions between NH_4^+ and NO_3^- uptake and assimilation: comparison of diatoms and dinoflagellates at several growth temperatures. *Mar Biol* 133:541–551
- Lupi FM, Fernandes HML, Tomé MM, Sá-Correia I, Novais JM (1994) Influence of nitrogen source and photoperiod on exopolysaccharide synthesis by the microalga *Botryococcus braunii* UC 58. *Enzym Microb Technol* 16:546–550
- Magaletti E, Urbani R, Sist P, Ferrari CR, Cicero AM (2004) Abundance and chemical characterization of extracellular carbohydrates released by the marine diatom *Cylindrotheca fusiformis* under N- and P-limitation. *Eur J Phycol* 39:133–142
- Mezhoud N, Zili F, Bouzidi N, Helaoui F, Ammar J, Ouada HB (2014) The effects of temperature and light intensity on growth, reproduction and EPS synthesis of a thermophilic strain related to the genus *Graesiella*. *Bioprocess Biosyst Eng* 37:2271–2280
- Neoh CH, Yahya A, Adnan R, Abdul Majid Z, Ibrahim Z (2013) Optimization of decolorization of palm oil mill effluent (POME) by growing cultures of *Aspergillus fumigatus* using response surface methodology. *Environ Sci Pollut Res* 20:2912–2923
- Ni WM (2014) Impact of nutrients and temperature on algal growth and relative abundance of Xixi wetland, China. *Appl Mech Mater* 665:426–429
- Nur MMA, Buma AGJ (2018) Opportunities and challenges of microalgal cultivation on wastewater, with special focus on palm oil mill effluent and the production of high value compounds. *Waste Biomass Valor*. <https://doi.org/10.1007/s12649-018-0256-3>
- Nur MMA, Setyoningrum TM, Budiaman IGS, Kristanto D (2016) Utilization of microalgae cultivated in palm oil mill wastewater to produce lipid and carbohydrate by employing microwave-assisted irradiation. *Recent Innov Chem Eng* 9:107–116
- Nur MMA, Setyoningrum TM, Budiaman IGS (2017) Potency of *Botryococcus braunii* cultivated on palm oil mill effluent wastewater as a source of biofuel. *Environ Eng Res* 22:417–425
- Nur MMA, Muizelaar W, Boelen P, Buma AGJ (2019) Environmental and nutrient conditions influence fucoxanthin productivity of the marine diatom *Phaeodactylum tricorutum* grown on palm oil mill effluent. *J Appl Phycol* 31:111–122

- Onyla CO, Uyub AM, Akunna JC, Norulaini NA, Omar AKM (2001) Increasing the fertilizer value of palm oil mill sludge: bioaugmentation in nitrification. *Water Sci Technol* 44:157–162
- Painter TJ (1993) Carbohydrate polymers in desert reclamation: the potential of microalgal biofertilizers. *Carbohydr Polym* 20:77–86
- Poh PE, Yong WJ, Chong MF (2010) Palm oil mill effluent (POME) characteristic in high crop season and the applicability of high-rate anaerobic bioreactors for the treatment of POME. *Ind Eng Chem Res* 49:11732–11740
- Ramus J (1977) Alcian blue: a quantitative aqueous assay for algal acid and sulfated polysaccharides. *J Phycol* 13:345–348
- Rhee G-Y, Gotham IJ (1981) The effect of environmental factors on phytoplankton growth: temperature and the interactions of temperature with nutrient limitation. *Limnol Oceanogr* 26:635–648
- Smith VH, Tilman GD, Nekola JC (1998) Eutrophication: impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environ Pollut* 100:179–196
- Soanen N, Da Silva E, Gardarin C, Michaud P, Laroche C (2016) Improvement of exopolysaccharide production by *Porphyridium marinum*. *Bioresour Technol* 213:231–238
- Staats N, Stal LJ, Mur LR (2000) Exopolysaccharide production by the epipelagic diatom *Cylindrotheca closterium*: effects of nutrient conditions. *J Exp Mar Biol Ecol* 249:13–27
- Tabassum S, Zhang Y, Zhang Z (2015) An integrated method for palm oil mill effluent (POME) treatment for achieving zero liquid discharge—a pilot study. *J Clean Prod* 95:148–155
- Tong SL, Jaafar AB (2004) Waste to energy: methane recovery from anaerobic digestion of palm oil mill effluent. *Energy Smart* 4:1–8
- Van de Poll WH, Visser RJW, Buma AGJ (2007) Acclimation to a dynamic irradiance regime changes excessive irradiance sensitivity of *Emiliania huxleyi* and *Thalassiosira weissflogii*. *Limnol Oceanogr* 52:1430–1438
- Van Oijen T, Veldhuis MJW, Gorbunov MY, Nishioka J, Van Leeuwe MA, De Baar HJW (2005) Enhanced carbohydrate production by Southern Ocean phytoplankton in response to in situ iron fertilization. *Mar Chem* 93:33–52
- Villanova V, Fortunato AE, Singh D, Bo DD, Conte M, Obata T, Jouhet J, Fernie AR, Marechal E, Falciatore A, Pagliardini J, le Monnier A, Poolman M, Curien G, Petroustos D, Finazzi G (2017) Investigating mixotrophic metabolism in the model diatom *Phaeodactylum tricorutum*. *Phil Trans Roy Soc B* 372:20160404
- Zhu CJ, Lee YK (1997) Determination of biomass dry weight of marine microalgae. *J Appl Phycol* 9:189–194

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Sulfated exopolysaccharide production and nutrient removal by the marine diatom *Phaeodactylum tricoratum* growing on palm oil mill effluent

ORIGINALITY REPORT

15%

SIMILARITY INDEX

15%

INTERNET SOURCES

%

PUBLICATIONS

%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

15%

★ link.springer.com

Internet Source

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%