

III.C.1.a. The Understanding of Volcanoclastics Model in Tebing

by Basuki Rahmad

Submission date: 05-Apr-2023 03:44PM (UTC+0700)

Submission ID: 2056454937

File name: .C.1.a._The_Understanding_of_Volcanoclastics_Model_in_Tebing.pdf (772.58K)

Word count: 2090

Character count: 12309

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/342313548>

The Understanding of Volcanoclastics Model in Tebing Breksi Geotourism By Digital Outcrop Model

Conference Paper · January 2019

CITATIONS

0

READS

88

4 authors, including:



Muchamad Ocky Bayu Nugroho

Universitas Pembangunan Nasional "Veteran" Yogyakarta

16 PUBLICATIONS 3 CITATIONS

SEE PROFILE



Bambang Yuwono

Universitas Pembangunan Nasional "Veteran" Yogyakarta

27 PUBLICATIONS 77 CITATIONS

SEE PROFILE



All content following this page was uploaded by Muchamad Ocky Bayu Nugroho on 19 June 2020.

The user has requested enhancement of the downloaded file.

The Understanding of Volcanoclastics Model in Tebing Breksi Geotourism By Digital Outcrop Model

Muchamad Ocky Bayu NUGROHO¹, Muhamad SYAIFUDIN¹, Bambang YUWONO², Basuki RAHMAD¹

¹Department of Geological Engineering, Faculty of Technology Mineral, University of Pembangunan Nasional
"Veteran" Yogyakarta, Indonesia

²Department of Informatics Engineering, Faculty of Industrial Technology, University of Pembangunan Nasional
"Veteran" Yogyakarta, Indonesia

Email the first author: bayu.ocky@gmail.com ; bayu.ocky@upnyk.ac.id

ABSTRACT

Tebing Breksi Geotourism Object is one of geotourism object in Yogyakarta which part of UNESCO Gunung Sewu Geoheritage. Geologically, Tebing Breksi is included in Semilir Formation that consists of volcanoclastic sediments such as Tuff, Tuffaceous sandstone, Tuffaceous claystone and lapilli (Surono, 2009). This research is trying to apply digital technology to geological science through 3D Outcrop Model in order to understanding history, sedimentary process and depositional environment easier. The Digital outcrop model was build up by photo that acquired by drone with systematic captured. Sedimentology data was captured by conventional method and measuring section that represents all layers. Based on model and data, it shows the dynamics of sedimentary process from bottom to top and could be identified from sedimentary structures and lithofacies. Bottom part shows bedding structures that describe grain size changing from sandstone to conglomerate and refer to model (Kendall, 2004) match on Supra Fan Lobes, the middle part displaying a bedding structure and coarsening upward stacking pattern and interpreted as inner fan channel facies and the upper part is new sequence package interpreted as basin plain. As a whole, the volcanoclastics sediment in Tebing Breksi area was from great Semilir Eruption in Early Miocene (Smyth, 2011) which was deposited in submarine fan system to basin plain (Kendall, 2004). This model is being developed with additional data to be an Augmented Reality (AR) smartphone application for other purposes.

Keywords: Tebing Breksi, Digital Outcrop Model, Volcanoclastic, Geotourism..

INTRODUCTION

The application of technology in geological science is very useful both in academic and industrial terms. DOM (Digital Outcrop Model) or VOM (Virtual Outcrop Model) is one of the examples. DOM is a surface outcrop that represents 3D view mostly in a form of textured polygon mesh. DOM allows for interpretation and measurement of different geological features, e.g. orientation of geological surfaces, width and thickness of layers (Bellian et al., 2005). The quantity of identifiable and measurable geological features highly depends on the outcrop model resolution and accuracy (Buckley et al., 2008). DOM is helping a geologist to imagine and interpret a outcrop easier because it geometry and resolution that completely covered.

Semilir Formation is known widely as volcanoclastic deposit that well outcropped in Tebing Breksi geotourism object Yogyakarta. Formed from rock mining remnants which makes a good shape and geometry in a cliff. This research addresses the field scale architecture and dimension of volcanoclastic deposits into DOM and the objective is to utilize a model for the understanding of volcanoclastic deposits easier.

GEOLOGICAL BACKGROUND

Indonesia archipelago is well-known for great volcanic activity due to plate activities. Java island has a row of volcanoes elongates from west to east called "volcanic arc" that formed of subduction slab since the Early Cenozoic (Hall, 2002; Smyth et al., 2008b). The Eocene to Early Miocene Volcanic arc in East Java is parallel to and south of modern/present volcanic arc (Smyth et al., 2011) where Semilir Mountain was present (Fig. 1). Southern Mountain volcanic arc has produced thick volcanic and volcanoclastic deposits that well preserved in Central Java and East Java, one of them is in Tebing Breksi. According to volcanic activity which had been started in Early Miocene, the Southern Mountain Volcanic Arc was a small volcanic island surrounded by carbonate volcanoclastic shelves separated by intra-arc basin (Smyth et al., 2011).

Volcanoclastic Deposits of Semilir Formation

Semilir Formation is well preserved and outcropped in several spots in Southern Mountain Area, such as Baturagung, Wonogiri, Tepus, and Candi Ijo. Several authors notes if in each spot of Semilir Formation has specific sedimentation character that representing volcano facies. Semilir volcanic activity was extensive, explosive and of Pillian Type with andesite-rhyolite composition include thick ashes, pumice lithic,

breccias, lava flows and crystal lithic tuff (Smyth, 2005). The volcanic rocks are characterized by features typical of terrestrial air-fall, pyroclastic surge and flow deposits, crystal layering, well-sorted granular laminations, diffuse bedding, breccias (with metre-scale pumice blocks), thick mantling ashes, and abundant fragments of charcoal. Some places there are found flame, traction, irregular bases and slump folds indicating an unstable marine slope, so it could be interpreted as erupted in subaerial setting in which pyroclastic flow entering the sea and some redeposited as lahar in distal settings (Smyth et al., 2011).




Figure 1. Modern Volcanic arc and Southern Mountain Volcanic Arc (Smyth et.al, 2011)

METHOD

This research use the integrated method of references review, fieldwork and information technology. References review was very important due to imagine a historical geology for model building while fieldwork was conducted to characterise the outcrop deposits include litology, sedimentary structure and layer position. Using information technology in term "DOM" makes Understanding more easier and comprehensive. Aquisition data using professional UAV have been done with horizontally and vertically schematic capture (Fig.2). Editing, compiling and processing graphic are the next step that supported with several software such as *Adobe Photoshop*, *CorelDraw* and *Zephyr 3D*.

The model or DOM has been integrated with field data and then be analyzed and compared with volcanoclastic model from references.




Figure 2. Schematics Capturing of Tebing Breksi and Professional UAV (Nugroho et.al, 2019)

DISCUSSION

Stratigraphic Analysis

Stratigraphic section has covered all layer that shows several sequences of sedimentation event. Sedimentary structure, texture and material composition is the main parameter that will be integrated with DOM. Semilir Formation in Tebing Breksi area is epiclastics deposit that formed by reworking pyroclastic or epiclastic previous deposit with a submarine-fan deposit system. Major portions of submarine-fan sequences are composed of turbidites that occur as channel-fill, lobes, and sheet sands (Shanmugam, 1988).

As shown in Fig. 3, the stratigraphic section of Tebing Breksi can be divided into 2 depositional sequences that interpreted from sediment supply due to subaqueous condition.

The lowest part of section dominated with conglomerate and massive tuffaceous sandstone. Parallel lamination and graded bedding are sedimentary structures that dominant in sight address to lowstand system tract that interpreted as suprafan lobes facies (Kendall, 2004). In the middle part shown as stacking tuffaceous sandstone interbedded with tuffaceous claystone that dominantly massive, graded bedding and parallel lamination sedimentary structures. The section is clearly display about transgressive system tract on middle channel fan facies (Kendall, 2004). Over previous section is quite clearly showing a high system tract deposits that characterized by coarse to medium tuffaceous sandstone with fining upward stacking's trend and interpreted as boundary of sequence. Wavy lamination is shown in upper part of section, parallel lamination, graded bedding and massive structures are dominantly shown in almost every beds, so section could be predicted as Inner channel facies (Kendall, 2004).

Above the sequence boundary, we find more dominant tuffaceous claystone, layer of tuffaceous breccia and tuffaceous pebbly sandstone. It could be interpreted as transgressive system tract due to stacking pattern was finning upward and deposited in basin plain facies.




Figure 3. Stratigraphic Section of Tebing Breksi from bottom to top (from left to right)

DOM Integration and Volcaniclastic Model

The model has a minimum sedimentary structures due to resolution that difficult to paleocurrent interpretation, but the dipping average of bedding plan about 10-13° with dominantly east direction concluding if paleocurrent direction is N 270°-320° E and correlated with sedimentary structure field data. Volcanoclastic model of Tebing Breksi Area was build from DOM integrated with stratigraphic section and references. We divided Tebing Breksi into 4 stacking sequences and have each depositional mechanism. Overall, we selected 1 DOM from the best one because another DOM have minimum resolution due to weathering and discoloration.

On Figure 4, Red line shows bedding structures that describes grain size changing from sandstone to conglomerate and refer to model (Kendall, 2004) match on Supra Fan Lobes, (Wood, 2017). Yellow line in above, has minimum structure in lower part but lamination and bedding structure in upper part. Two kind of that structure separated with tuff layer (clay size) in mean massive transgressive moment. Generally, this section has finning upward stacking pattern and it correlates to middle fan channel facies (Kendal, 2004) and yellow circle on (Wood, 2017) model.

On Blue line overall there is no specific structure and rather difficult identified from DOM, but in detail, we

can see a vague parallel bedding and color gradation. Related with field data in the upper part that has graded bedding structure and coarsening upward stacking pattern, so it displaying inner fan channel facies (Kendall, 2004) and blue circle on (Wood, 2017) model. The top section (green line) look very different compared and clearly showing a bedding structure although has been discoloration. Generally, it shows a special sedimentary package that could be mean as a new sequence.





Figure 4. DOM integrated with Volcaniclastic Submarine Model (Kendall, 2004 & Wood, 2017)

CONCLUSION

DOM (Digital Outcrop Model) is one of tools that can use for geosciences purposes recently and for example is building geological model in some areas. Building a DOM using a UAV through several stages, they are data acquisition, processing and visualization. UAV DOMs are not a replacement for field observations, although, they can efficiently provide supplemental measurements and enable guided interpretation data (i.e., sedimentary logs, cross-section sketches). In this study, we used field data (stratigraphic section) to improving and detailing geologic information for building a model.

ACKNOWLEDGMENTS

Alhamdulillah, we would like to thank God Almighty Allah SWT and Prophet Muhammad, LPPM UPN "Veteran" Yogyakarta for funding this research, Geological Engineering and Informatic Engineering Department also all those who helped in this research.

REFERENCES

- Bellian, J.A., Kerrans, C., Jennette, D.C., Digital Outcrop Models: Applications Of Terrestrial Scanning Lidar Technology In Stratigraphic Modeling, *Journal Of Sedimentary Research*, Vol. 75, No. 2, March, 166–176 (2005).
- Buckley, S.J., Howell, J.A., Enge, H.D., Kurz, T.H., Terrestrial laser scanning in geology: data acquisition, processing and accuracy considerations, *Journal of the Geological Society, London*, Vol. 165, 625–638 (2008).
- Hall, R., Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. *Journal of Asian Earth Sciences* 20, 353–434 (2002).
- Hall, R., Smyth, H.R., Cenozoic arc processes in Indonesia: identification of the key influences on the stratigraphic record in active volcanic arcs. In: Draut, A., Clift, P.D., Scholl, D.W. (Eds.), *Formation and Applications of the Sedimentary Record in Arc Collision Zones: Special Paper*. Geological Society of America, Boulder, 27–54 (2008).
- Nugroho, M.O.B., Syaifudin, M., Yuwono, B., Sinangaseta, G., Aplikasi Alat Bantu Foto Udara (Drone) Dan Pengeditan Foto Untuk Mendukung Pengambilan Data Geologi Pada Tebing Curam, Studi Kasus : Tebing Breksi, Yogyakarta, *Proceedings Seminar Nasional LPPM UPN "Veteran" Yogyakarta* (2019).
- Shanmugam, G., Moiola, R.J., *Submarine Fans: Characteristics, Models, Classification and Reservoir Potential*, Earth-Science Reviews, No.24, Elsevier Science Publishers B.V, Netherland, 383–428 (1988).
- Smyth, H., Hall, R., Hamilton, J.P., Kinny, P., East Java: Cenozoic basins, volcanoes and ancient basement, Indonesian Petroleum Association. *Proceedings of the 30th Indonesian Petroleum Association Convention*. Indonesian Petroleum Association, Jakarta, 251–266 (2005).
- Smyth, H., Crowley, Q.G., Hall, R., Hamilton, J.P., Kinny, P., Schmidt, D.N. A Toba-scale eruption in the Early Miocene: The Semilir eruption, East Java, Indonesia, *Lithos*, Elsevier Publisher, 198–211 (2011)
- Surono., Litostratigrafi Pegunungan Selatan Bagian Timur Daerah Istimewa Yogyakarta Dan Jawa Tengah, *Jurnal Sumber Daya Geologi* Vol.19 No.3, Juni, 209–221 (2009)

III.C.1.a. The Understanding of Volcanoclastics Model in Tebing

ORIGINALITY REPORT

17%
SIMILARITY INDEX

17%
INTERNET SOURCES

0%
PUBLICATIONS

0%
STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

17%
★ www.researchgate.net
Internet Source

Exclude quotes On
Exclude bibliography On

Exclude matches < 10%