

URBAN DEVELOPMENT
AND INFRASTRUCTURE

Wayan Suparta, PhD
Editor

Urban Development and Lifestyle


NOVA

URBAN DEVELOPMENT AND INFRASTRUCTURE

URBAN DEVELOPMENT AND LIFESTYLE

WAYAN SUPARTA

EDITOR



nova
science publishers
New York

The logo for Nova Science Publishers features the word "nova" in a large, bold, lowercase sans-serif font. Below it, the words "science publishers" are written in a smaller, lowercase sans-serif font, and "New York" is written in an even smaller font at the bottom. A thick, curved arrow starts from the left side of the "nova" text and points towards the right, arching over the top of the text.

Copyright © 2020 by Nova Science Publishers, Inc.

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic, tape, mechanical photocopying, recording or otherwise without the written permission of the Publisher.

We have partnered with Copyright Clearance Center to make it easy for you to obtain permissions to reuse content from this publication. Simply navigate to this publication's page on Nova's website and locate the "Get Permission" button below the title description. This button is linked directly to the title's permission page on copyright.com. Alternatively, you can visit copyright.com and search by title, ISBN, or ISSN.

For further questions about using the service on copyright.com, please contact:

Copyright Clearance Center

Phone: +1-(978) 750-8400

Fax: +1-(978) 750-4470

E-mail: info@copyright.com.

NOTICE TO THE READER

The Publisher has taken reasonable care in the preparation of this book, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained in this book. The Publisher shall not be liable for any special, consequential, or exemplary damages resulting, in whole or in part, from the readers' use of, or reliance upon, this material. Any parts of this book based on government reports are so indicated and copyright is claimed for those parts to the extent applicable to compilations of such works.

Independent verification should be sought for any data, advice or recommendations contained in this book. In addition, no responsibility is assumed by the Publisher for any injury and/or damage to persons or property arising from any methods, products, instructions, ideas or otherwise contained in this publication.

This publication is designed to provide accurate and authoritative information with regard to the subject matter covered herein. It is sold with the clear understanding that the Publisher is not engaged in rendering legal or any other professional services. If legal or any other expert assistance is required, the services of a competent person should be sought. FROM A DECLARATION OF PARTICIPANTS JOINTLY ADOPTED BY A COMMITTEE OF THE AMERICAN BAR ASSOCIATION AND A COMMITTEE OF PUBLISHERS.

Additional color graphics may be available in the e-book version of this book.

Library of Congress Cataloging-in-Publication Data

ISBN: 978-1-53618-560-7

Names: Suparta, Wayan, editor.

Title: Urban development and lifestyle / [edited by] Wayan Suparta,
Universitas Pembangunan Jaya Tangerang Selatan, Banten, Indonesia.

Description: Hauppauge : Nova Science Publishers, 2020. | Series: Urban
development and infrastructure | Conference proceedings. | Includes
bibliographical references and index. |

Identifiers: LCCN 2020039650 (print) | LCCN 2020039651 (ebook) | ISBN
9781536185607 (hardcover) | ISBN 9781536186314 (adobe pdf)


Subjects: LCSH: Sustainable urban development--Congresses. | City
planning--Congresses. | Community development--Environmental
aspects--Congresses. | Intelligent transportation systems--Congresses.

Classification: LCC HT241 .U6973 2020 (print) | LCC HT241 (ebook) | DDC
307.1/416--dc23

LC record available at <https://lcn.loc.gov/2020039650>

LC ebook record available at <https://lcn.loc.gov/2020039651>

Published by Nova Science Publishers, Inc. † New York



As Professor at the Universitas Pembangunan Jaya, with daily activities in lecturing, doing research, as well as water resources development planning, I really praise the Nova Science Publishers for publishing selected papers from “2020 International Conference on Urban Sustainability, Environment, and Engineering (CUSME 2020)”. Hence, this publication would be useful for professionals, researchers, scholar, policymakers, and NGO. I believe that currently, many professionals would like to give more attention on development of sustainable urban. In addition, this publication could be used as reference for City authorities to make appropriate policy choices to protect the provision of equitable housing, health, and transportation services.

Prof. Ir. Frederik Josep Putuhena M.Sc., Ph.D
Center for Urban Studies – Universitas Pembangunan Jaya



Urban Development and Lifestyle are trend issues for the cities around the world. Learning from experiences is the most effective way to support the cities to be sustainable developed. This book offers the knowledge sharing among countries which covers variety of cities’ issues. It also provides the great lessons for researchers, officers and policy makers on coping with several urban problems.

Associate Professor Sarintip Tantanee, Ph.D.
Director
Center of Excellence on Energy Technology and Environment
(CETE)
Faculty of Engineering, Naresuan University, Thailand

CONTENTS

Preface		xiii
Urban Psychology, Cultural, and Communication Studies		1
Chapter 1	Ecological Behavior Model: Explained By Rational and Moral Perspective <i>Hetti Rahmawati, Siti Sendari and Yuni Rahmawati</i>	3
Chapter 2	Mapping Digital Political Communication for Millennials in Indonesia <i>Naurissa Biasini, Emma Rachmawati and Yosaphat Danis Murtiharso</i>	13
Chapter 3	Case Study of Urban Baby Boomer in Hoax Messages Distribution through Whatsapp Application in Indonesia <i>Suci Marini Novianty and Emma Rachmawati</i>	25
Chapter 4	Phubbing in Urban Families (A Case Study on the Families in South Tangerang) <i>Sri Wijayanti and Nathaniel Antonio Parulin</i>	35
Chapter 5	Communication for Risk Reduction in Natural Disaster (Case Study: National Disaster Management Agency – Indonesia) <i>Reni Dyanasari, Fasya Syifa Mutma and Melisa Arisanty</i>	45
Urban Education and Community Services		59
Chapter 6	Development of Social Media-Based Online Shop (Instagram and Facebook) in Economic Learning <i>Iskandar Agie Hanggara</i>	61
Chapter 7	ICT Integration in English Language Teaching in a University: Academic Perspectives <i>Endang Darsih</i>	71

Chapter 8	The Sustainable Community Development in Learning Activities for Children at Suburban Area <i>Fitriyah Nurhidayah, Johannes Hamonangan Siregar and Chaerul Anwar</i>	79
Chapter 9	Introducing Science, Technology, Engineering, Arts, and Math through Creative Exercise to Stimulate a Child's Creativity and Motoric Sense in an Urban Environment <i>Ismail Alif Siregar</i>	93
	Urban Economics and Lifestyle	103
Chapter 10	The Influence of Income and Financial Literacy on Financial Satisfaction through Financial Behavior as a Mediating Variable <i>Khairina Natsir, Made Setini and Agus Zainul Arifin</i>	105
Chapter 11	Marketing and Promoting the Concept of Malaysian Heritage Garden <i>Ahmad Zamil Zakaria, Melasutra Md Dali and Hazreena Hussein</i>	117
Chapter 12	Propensity for Sustainable Entrepreneurship of MSEs Owner in Yogyakarta, Indonesia <i>Kartika Nuringsih and Nuryasman MN</i>	131
Chapter 13	Strategic Facilities Management Approach in Australian Public Healthcare Organisation <i>Yuhainis Abdul Talib, Nor Aini Salleh and Kharizam Ismail</i>	141
Chapter 14	The Impact of Corporate Social Responsibility Implementation on the Value of Construction Firms in Indonesia <i>Muhammad Tony Nawawi and Agus Zainul Arifin</i>	151
Chapter 15	Value of Innovation: Creative Business Strategy of Nyatu Rubber Tree in Central Kalimantan <i>Vivy Kristinae, Made Wardana, I Gusti Ayu Kt Giantari and Agoes Ganesha Rahyuda</i>	167
Chapter 16	Promotion and Brand Trust in Affecting Repurchase Intention on E-Commerce Go-Jek Application <i>I Wayan Gede Antok Setiawan Jodi, Bagus Nyoman Kusuma Putra and I Made Surya Prayoga</i>	181
Chapter 17	Service Quality: Sales Strategy Towards Increasing Customer Satisfaction in Indonesian Modern Markets <i>Hastuti Naibaho and Yohanes Totok Suyoto</i>	189
Chapter 18	Factors Affecting Indonesian Sharia Bank Based on its Performances <i>Yanuar Yanuar and Eliza Christabella Phuanerys</i>	199

Chapter 19	The Influence of Financial Decision and Corporate Social Responsibility on Value of the Firm: Evidence from Manufacturing Companies in Indonesia <i>I Gede Adiputra, Nyoman Suprastha and Kevin Andreas</i>	207
Chapter 20	The Effect of Environmental Accounting and Fundamental Factors on Company Value <i>Irma P. Sofia and Sila Ninin W.</i>	217
Urban Architecture and Green Technology		231
Chapter 21	Atrium Design as Noise Control in Buildings <i>Santi Widiastuti and Jennie Kusumaningrum</i>	233
Chapter 22	IoT (Internet of Things) for Participatory Planning and Sustainable Urban Planning <i>S. Hariyani, E. B. Kurniawan, F. Usman and F. Shoimah</i>	247
Chapter 23	Amphibious Architecture: An Alternative Floodproof Design for Urban Flood Mitigation in Palembang, Indonesia <i>Harrini Mutiara Hapsari Wahyu, Rizka Drastiani and Sri Lilianti Komariah</i>	259
Agricultural Technology and Cultivation		269
Chapter 24	The Effect of Stem Cutting Type and Plastic-Covered on Bougainvillea Growth in Green Open Space <i>Heti Herastuti and E. K. Siwi Hardiastuti</i>	271
Chapter 25	Propagation of Arrowroot Plants <i>In Vitro</i> for the Agroforestry Environment Development <i>Endah Wahyurini and Susilowati</i>	281
Chapter 26	Potential of Kirinyuh (<i>Chromolaena odorata</i>) and Cow Manure to Increase the Nitrogen Uptake of Tomatoes (<i>Lycopersicum esculentum L.</i>) on Sandy Beach Soil <i>Lelanti Peniwiratri and Dyah Arbiwati</i>	289
Chapter 27	Optimum NPK Uptake of Paddy Soil By Applying Legume Residue and Combining Organic-Inorganic Fertilizer <i>Oktavia S. Padmini</i>	299
Chapter 28	The Role of Guano Phosphate in Improving the Quality of Compost from Household and Mushroom Media Waste in Supporting the Zero Waste Concept <i>Tuti Setyaningrum and Dyah Arbiwati</i>	307

Chapter 29	Increasing the Performance of Bougainvillea with Top Grafting for the Green Line in Urban Environment <i>Tutut Wirawati, Heti Herastuti and M. Husain Kasim</i>	317
Chapter 30	The Efforts to Improve the Quality of Growing Bulbil Iles-Iles (<i>Amorphophallus muelleri</i> Blume) with Oligo Chitosan Immersion <i>Sumarwoto and Sugeng Priyanto</i>	327
Chapter 31	Land Suitability Evaluation for Cassava Development in Integrated Farming System in Monggol <i>Purbudi Wahyuni, Dyah Sugandini, Didi Saidi, Sari Bahagiarti K and Olga Sisca Novaryan Scandiskti</i>	337
Chapter 32	The Screening of Growth, Yield Component, and Resistance of Various Sweet Corn Lines against Downy Mildew on S-3 Generation <i>Bambang Supriyanta, O. S. Padmini, M. Kundarto and Danar Wicaksono</i>	347
Chapter 33	Bamboo Construction Strengthening Techniques with Metal Materials in the Furniture Industry <i>Teddy Mohamad Darajat</i>	355
Urban Computing, Technology, and Smart Transportation		363
Chapter 34	Sefaira and Autodesk FormIt360: A Comparison of Energy-Efficient Building Simulation <i>Beta Paramita and Sarah Luziani</i>	365
Chapter 35	Fuzzy Time Series to Predict the Volume of Yield of Arrowroot <i>Rifki Indra Perwira, Danang Yudhiantoro and Endah Wahyurini</i>	375
Chapter 36	Computational Fluid Simulation of Self Bed Feedstock in Rice Husk Bubbling Fluidized Bed Gasification <i>Mummyyis Pramono, Hafif Dafiqurrohman, Kania Amelia Safitri, Iwan Setyawan and Adi Surjosatyo</i>	385
Chapter 37	Images Matching Using the SURF Method on the Face of Buddha Statue <i>Linda Marlinda, Supriadi Rustad, Ruri Sukobasuki, Fikri Budiman and Muhamad Fatchan</i>	397
Chapter 38	Analysis of Performance Proximity Sensor SRF-05 and Maxsonar-EZ2 Implemented Using Arduino <i>Erwan Eko Prasetyo and Farid Ma'ruf</i>	405
Chapter 39	Smartgrid Technology and Development in Thailand <i>Jirawadee Polprasert, Sarawut Wattanawongpitak, Suchart Yammen and Thitipong Samakpong</i>	415

Chapter 40	Product and Visual Development of Portable Dynamic Message Sign <i>Desi Dwi Kristanto, Resdiansyah and Prio Handoko</i>	425
Chapter 41	Magnesium-Based Solid Storage for Fuel Cell Vehicle Application <i>Zulkarnain Jalil, Adi Rahwanto and Erfan Handoko</i>	435
Chapter 42	The Important Factors for Sustainable Padang - Pariaman Train Operation <i>Purnawan and Widia Safira</i>	443
Climate Change, Disaster, and Environmental		453
Chapter 43	The Relation Pedestrian Age and Gender to Walkability in Commercial Area of Pangkalpinang City <i>Revy Safitri and Ririn Amelia</i>	455
Chapter 44	Synthesis CdS/Pt-TiO ₂ with Enhanced Its Performance for Photocatalytic Degradation of Palm Wastewater Treatment <i>Ratnawati, Singgih Hartanto, Yuli Amalia Husnil and Christin Rina Ratri</i>	463
Chapter 45	The Efficiency of Produced Water Treatment Using Combination of Coagulation Process and Membrane Bioreactor <i>Ayu Utami, Wibiana W. Nandari, Ekha Yogafanny, E. A. Kristiati, Tedy A. Cahyadi and Nur Ali Amri</i>	475
Chapter 46	The Application Practice of Solar Energy Generation in Indonesia's Textile Industry <i>Herawati Rubiana, Mochammad Chaeron and Apriani Soepardi</i>	485
Chapter 47	Towards Sustainable Food Landscape: Generating and Protecting Spatial Context <i>Hedista Rani Pranata, Dalhar Susanto and Toga H. Panjaitan</i>	495
Chapter 48	Perception and Adaptation of Farmers on Climate Change (A Case Study at Clove Farmer in Samigaluh Kulonprogo, Yogyakarta) <i>Antik Suprihanti, Dwi Aulia Puspitaningrum, Eko Amiadji Julianto and Herwin Lukito</i>	505
About the Editor		515
Index		519

Chapter 26

**POTENTIAL OF KIRINYUH (*CHROMOLAENA ODORATA*)
AND COW MANURE TO INCREASE THE NITROGEN
UPTAKE OF TOMATOES (*LYCOPERSICUM
ESCULENTUM L.*) ON SANDY BEACH SOIL**

Lelanti Peniwiratri and Dyah Arbiwati*

Department of Soil Science, Faculty of Agriculture,
UPN "Veteran" Yogyakarta Indonesia

ABSTRACT

This research was conducted to determine the Nitrogen uptake of tomatoes (*Lycopersicum esculentum L*) on the sandy beach soil which was of kirinyuh (*Chromolaena odorata*) and cow manure application. The research method used was a complete randomized design consisting of two factors. The first factor is the giving of organic material consisting of 4 levels, i.e., without kirinyuh + cow manure (B0), 30 tons/ha kirinyuh (B1), 30 tons/ha cow manure (B2) and 30 tons/ha mixture of cow manure and kirinyuh. The second factor is the application time of organic material consisting of 3 levels, i.e., at the time planting (T1), 20 days before planting (T2) and at 40 days before planting (T3). Each treatment was replicated three times. The results showed that cow manure and kirinyuh were able to increase the available Nitrogen of sandy beach soil and Nitrogen uptake of tomato. Then 30 tons/ha mixture of cow manure and kirinyuh applicated at 20 days before planting affects the hightest of Nitrogen uptake of tomatoes.

Keywords: Kirinyuh, Cow manure, Nitrogen uptake, Tomato

INTRODUCTION

Nitrogen as one of the macro nutrients is needed by plants for growth. Nitrogen derived from fertilizers and free air can be absorbed by plants in the form of NH_4^+ and NO_3^- [1]. The element is much needed by tomatoes in the vegetative phase to form proteins, chlorophyll, and other organic compounds [2]. Tomatoes are very responsive in the provision of N nutrients, therefore in the tropics these nutrients often show a real effect. When the N supply is sufficient, plant leaves grow large and expand the surface available for photosynthesis, accelerating the conversion of carbohydrates into proteins which are subsequently converted into protoplasm, functioning as vital substances of various enzymes that guide the entire process of plant metabolism [3].

Tomatoes (*Lycopersicon esculentum* L) are a horticultural commodity that is multipurpose and has the potential to be developed in Indonesia. Tomato production in Yogyakarta in 2015 reached 12.5 tons [4] and each year is increasing to keep pace with the increasingly high needs of the community. This plant can grow well in the highlands (more than 700 m above sea level), medium plains (200 m - 700 m above sea level), and lowlands (less than 200 m above sea level) [5]. Growing media that are suitable for tomatoes are clay or dusty clay textures, acumbly structure containing a lot of organic matters and easily binding water, so that not many nutrients are leached. Tomatoes are horticultural plants that can grow in the lowlands and highlands and have the potential to be developed. Tomatoes require loose soil, acidity levels of 5-6, soil contains little sand, and contains a lot of humus, regular, and sufficient irrigation, and the addition of Nitrogen to support vegetative growth [6]. For optimum growth, tomatoes require N 200-230 kg/ha [7].

The sandy beach soil is one of the types of land that has the potential as a medium for growing tomatoes because of its wide spread in Indonesia and has not been widely used, but in its use, the sandy beach soil has obstacles in both physical and chemical properties. Sandy beach soil has soil properties such as sandy texture, single grain structure to crumbs, loose consistency when dried, a low moisture holding capacity, high water escape ability, and pH ranges from 6-7 [8]. This soil also has low organic matter content, low cation exchange capacity. Total nutrient content is actually high but the available nutrient content is low, especially Nitrogen [9, 10].

The low availability of Nitrogen in sandy beach soil can a use stunted tomato growth. Tomatoes need large amounts of Nitrogen for optimal growth. In one season, tomatoes need Nitrogen given in the form of urea of 175 kg/ha [11]. If specifically calculated (45% N in urea), the Nitrogen given to tomatoes is 78.75 kg N/ha. If these needs are met by an artificial N fertilizer, it will require a large cost and, in the long run, will have a negative impact on the environment.

To overcome these obstacles, manipulation needs to be done, so that the sandy beach soil can be used as a medium for growing tomatoes. Manipulation that can be done is the addition of organic fertilizers in the form of cow manure and kirinyuh together, and set the application time of organic matter. The addition of organic fertilizers is expected to provide benefits for improving the quality of soil structure. With improved soil structure and good pore balance and spread, soil aggregates can provide a dense balance and pore space that is more beneficial for tomato growth.

Cow manure is a fertilizer that comes from cattle cages, in the form of solid manure mixed with food scraps and urine. All waste products from cows in fresh or composted form in a solid or liquid form have natural properties and do not damage the soil, and provide macro elements (nitrogen, phosphorus, potassium, calcium, and sulfur) and the micro elements (iron, zinc, boron, cobalt, and molybdenum). Cow manure is useful for improving physical, chemical, and biological soil properties, increasing water retention capacity, soil microbiological activity, cation exchange capacity value and adding nutrients [12].

Kirinyuh (*Chromolaena odorata*) is a potential weed to be used as a source of organic fertilizers because of high biomass production. The production of *Chromolaena odorata* biomass is 18.7 tons/ha in the fresh form and 3.7 kg/ha in the dried form, consisting of content of 103.4 kg/ha of N, 15.4 kg/ha of P, 80.9 kg/ha of K, and 63.9 kg/ha of Ca. Kirinyuh can be used as a source of green fertilizers, increase the content of N elements in the soil, and also reduce the use of synthetic fertilizers [13]. The result of the study by Nurhajati [14], it was found that Kirinyuh that grew alongside the road obtained fresh materials of 3 kg/ha. At the age of 6 months, *Chromolaena odorata* produces biomass of 11.2 tons/ha, and after 3 years of age is able to produce biomass of 277.7 tons/ha. Weed biomass has a high nutrient content (2.65% N, 0.53% P and 1.9% K) [15]. If pruned, then 3 (three) months, it will grow back and even can produce 4 tons/ha or the equivalent of 1.2 tons/ha of dry material containing artificial fertilizers (73 kg Urea, 97 kg SP-36 and 84 kg KCl) [16].

The release of Nitrogen from organic matter depends on the physical and chemical properties of the organic matter at the time of application and the community of the remodeled organisms. The content of N, lignin and polyphenols is the main factor that determines whether or not organic matter decomposes easily and releases N. The right application time of organic matter can help the availability of necessary N nutrients and will increase the Nitrogen uptake by plants [17].

This study aims to determine the potential of Kirinyuh weed and cow manure based on the right time of application to increase Nitrogen uptake of tomatoes on sandy beach soil.

METHODS

This research was conducted in Srigading Village, Sanden District, Bantul Regency, D. I. Yogyakarta. The analysis was carried out at the Laboratory of Plant Nutrition and Fertilizer Technology, Faculty of Agriculture, UPN "Veteran" Yogyakarta.

This research is a pot experiment, with a completely randomized two-factor design method. The first factor is giving of organic matter consisting of 4 levels, i.e., without kirinyuh and cow manure (B0), 30 tons/ha of kirinyuh or equivalent to 474 grams/pot (B1), 30 tons/ha of cow manure or equivalent to 474 grams/pot (B2) and 30 tons/ha mixture of cow manure and kirinyuh or equivalent to 237 grams/pot. The second factor is the application time of organic matter consisting of 3 levels, i.e., at the time of planting (T1), 20 days before planting (T2), and 40 days before planting (T3). Each treatment was replicated 3 (three) times. Therefore, the number of trials is 36 pots

Table 1. Physical and chemical properties of sandy beach soil

Soil Properties	Values	Rate [18]
C – Organic (%)	0,14	Very low
N – Total (%)	0,206	Very low
N – available (%)	0,013	Very low
P – available (ppm)	9,73	Low
K – available (me %)	0,237	Very low
pH H ₂ O	6,75	Neutral
CEC (me %)	5,9	Low
Fraction :		
Sand (%)	90,96	High
Silt (%)	5,84	Low
Clay (%)	3,2	Low
Texture class		Sand
Bulk density (g/cm ³)	1,63	
Particle density (g/cm ³)	2,73	

Table 2. Chemical compositions of Kirinyuh and cow manure

Organic Matters	Composition		C/N
	C-organic (%)	N Content (%)	
Kirinyuh	31,00	2,56	12,10
Cow Manure	17,74	1,42	12,49

The sandy beach soil taken from the field was air dried and sieved through a 2 mm. This soil is put in a plastic pot, in which each pot is equivalent to 8 kg of absolute dried soil. The treatment is given by giving cow manure and kirinyuh based on the treatments, namely without giving cow manure and kirinyuh (control), giving 474 grams/pot or equal to 30 tons/ha, giving cow manure 474 grams/pot or equal to 30 tons/ha, and a mixture of cow and kirinyuh manure of 237 grams/pot (50%: 50% ratio). Application of cow and kirinyuh manure is given based on the treatment, namely at the time of planting, 20 days before planting, and 40 days before planting. After being mixed evenly between the soils, manure and kirinyuh are then given water until a field capacity condition is reached. Soil moisture is maintained in the condition of field capacity, namely by weighing the water. The amount of water is determined by weighing the experiment pots every day. The difference between the weight and contents added.

As a bioessay to determine the response of cultivated plants to soil conditions due to treatment, planting tomato seedlings is carried out on the soils in the treated pots. Planting is carried out on soil samples with an equivalent weight of 8.3 kg of soil in the field condition. Before planting, the tomato seedlings are made nurseries after the age of 20 days by taking several plants that grow well and are relatively uniform. Planting tomatoes is planted with 3 plants per pot. At the age of 7 days after planting thinning is done by leaving one of the best plants per pot. Plants maintained reach a maximum vegetative phase. After the maximum vegetative phase is reached, soil sampling is carried out to test the Nitrogen availability of

sandy beach soil, plant tissue sampling for plant growth observation, nutrient test for plant tissue nitrogen, and plant tissue absorption, which is calculated using the following formula:

$$N - \text{uptake (g/pot)} = N \text{ tissue content (\%)} \times \text{Plant dry weight (g)} \quad (1)$$

To find out the effect of treatment on the parameters of the experiment, Analysis of Variance with significant of 5% was used, while to find out the comparison between treatments, Duncan's Multiple Range Test (DMRT) with significant of 5% was used [19].

RESULTS AND DISCUSSION

The results of the soil analysis at the study site (Table 1) showed that the beach sandy soil used in the study was dominated by 90.96% sand fraction (sand texture). With this texture, the surface area is small and has more macro pores, so the ability of the soil to bind water is relatively low. This soil has a macro pore space that is very easy for water and air movement, so the soil porosity is high. This soil also has good aeration because it has a bulk density of 1.63 g/cm³ and a particle density of 2.73 g/cm³. Chemical fertility constraints include : very low organic C (0.14%), low CEC (5.9 me%) as well as other nutrients, namely total Nitrogen and very low N-available respectively 0.206% and 0.013%, therefore, it is necessary to make efforts to improve the fertility of this sandy beach soil. From Table 2, it can be seen that the Kirinyuh contains very high nitrogen of 2.56%, very high C-Organic content of 31% and low C/N of 12.10. Cow manure also contains very high Nitrogen of 1.42%, high organic C content of 17.74% and low C/N of 12.49. From these data, both kirinyuh and cow manure have the potentials to improve the soil fertility of sandy beach soil.

Nutrient uptake is an indicator of soil response to fertilization. The response of nitrogen uptake of tomatoes to the application of cow manure and kirinyuh at various times of application can be seen from the parameters of N-available soil, plant growth and N-uptake observed at the maximum vegetative age (Table 3). From Table 3, the results of the analysis of variance of 5% significance level, it can be seen that the treatment of organic matter at various times of application significantly affects the N-available of sandy beach soil, plant dry weight, tissue N content and N-uptake of tomatoes. This condition indicates that the addition of cow manure and kirinyuh with the time of application is able to significantly increase these parameters. This is due to nutrient N produced by cow manure and kirinyuh by setting the time of application, which is able to meet nutrient requirements in the maximum vegetative phase of tomatoes.

From Table 3, it can be seen that the mixture of cow and kirinyuh (B3) manure treatments showed the highest N-Available result of 0.07% and was significantly different from the B0, B1, B2 treatments. The treatment of a mixture of cow manure and kirinyuh (B3) is the best treatment for increasing N-available sandy beach soil, where kirinyuh is able to play a role in improving the quality of cow manure as a source of organic matter, whereas cow manure acts as a decomposer to accelerate the process of over hauling organic matter by microorganisms, the rate of decomposition increases so that the process of organic N mineralization into NH₄⁺ and NO₃⁻ goes well, then the need for tomatoes for N-available can be fulfilled. Mean while, without the use of cow manure and kirinyuh (B0), it actually reduces

the N-availablesandy beach soils. This is due to the absence of cow manure which acts as a decomposer. Decomposer microorganisms found in cow manure are able to add microorganisms found in the soil. As a result of the absence of cow manure, the number of organic matter remover microorganisms is only small, the process of overhauling organic material becomes slow, and the number of the available N produced is little. The application of cow and kirinyuh manure at 20 days before planting (T2) is able to increase the highest available N and is significantly different from the simultaneous planting time (T1) because T2 is the best time for manure and kirinyuh to release the maximum N nutrients contained in cow manure and kirinyuh, while giving organic matter at 40 days before planting (T3) actually gives results that are not significantly different.

The increase in available N in the soil appears to respond to plant growth as indicated by the increase in plant dry weight (biomass) and plant tissue N-uptake (Table 3), which is in this case tomatoes uptake N from the soil in the available form (NO_3^- and NH_4^+) because of the addition of N from kirinyuh and cow manure (Table 2). This increase in nutrient uptake is due to an improved crop root system as evidenced by the increase in total dry weight of plants in Table 3. Such increase is due to the increased N availability of sandy beach soil. This is because photosynthates in plant tissues that are formed due to the application of a mixture of cow manure and kirinyuh given are widely used for cell division. This is consistent with the results of research [20] which states that an increase in total N-soil can increase the N availability of soil, which increases plants N uptake because N is absorbed by plants in the form of the available N (NO_3^- and NH_4^+) of soil. In fact, this is due to the addition of N from the cow manure and kirinyuh

Table 3. Effects of cow and kirinyuh manure at various times of application to N-available soil, plant dry weight, tissue N-content and N-uptake of tomatoes

Treatments	N-Available (%)	Dry weight (g)	Tissue N Content (%)	N-Uptake (g/plant)
Organic Matter				
B0	0.04 c	5.92 c	0.34 c	0.15 c
B1	0.06 b	5.98 bc	1.05 b	0.28 b
B2	0.06 b	6.11 ab	1.09 b	0.29 b
B3	0.07 a	6.27 a	1.38 a	0.51 a
Application Time				
T1	0.05 q	5.94 q	0.59 q	0.14 q
T2	0.07 p	6.15 p	1.13 p	0.44 p
T3	0.06 p	6.12 p	1.17 p	0.34 p
Interaction	(-)	(-)	(-)	(-)

Description: Average followed by the same letter in the same row or column show there is non significant on Duncan's Multiple Range Test at the level of 5%.

B0: without kirinyuh and animal manure

B1: 30 tons/ha kirinyuh

B2: 30 tons/ha cow manure

B3: 30 tons/ha mixture of kirinyuh and cow manure

T1: at the time as planting

T2: 20 days before planting

T3: 40 days before planting

From Table 3, the uptake of tomato plants to the maximum vegetative age shows that the mixed treatment of cow manure and kirinyuh (B3) application at 20 days before planting (T2) shows the highest N uptake of tomato plants and is significantly different from the treatment of B0, B1, B2. This is because the mixture of manure and kirinyuh is the best treatment that is able to meet the availability of nitrogen nutrients in the soil that plants need to uptake and use as food through the process of photosynthesis to produce photosynthates that are used for plant growth, namely the dry weight of plants. The higher the photosynthate produced, the higher the photosynthate transplanted so that the dry weight of the plants will increase (Table 3). Meanwhile, without giving cow manure and kirinyuh (B0) produces the lowest N uptake. This is due to the absence of cow manure which acts as a decomposer. Decomposer microorganisms contained in cow manure will result in the slowing process of the organic material overhauling, little available N produced, and the least N uptake proven to be the smallest dry weight of tomato plants (B0). The N is needed by plants in quite large. Nitrogen in plants functions in the process of plant growth, affects the action of chlorophyll, and increases protein levels in the body of the plants. Meanwhile, the need for nitrogen for tomato plants lasts throughout life. Nitrogen uptake is not the same at each phase of tomato growth, so this plant requires continuous availability of nitrogen at all stages of growth. Therefore, if there is no addition of N in the soil (B0), plant growth will be inhibited.

ACKNOWLEDGMENT

The paper is part of the results of the applied research LPPM (2019) entitled Response of Tomato (*Lycopersicon esculentum* L.) Nutrient Uptake in Sandy Beach Soil Which Was Given Kirinyuh Weeds (*Chromolaena odorata*) and Cow Manure. Thanks go to the Chairman of LPPM of UPN "Veteran" Yogyakarta.

CONCLUSION

The application of cow manure and kirinyuh (*Chromolaena odorata*) can increase the available N of sandy beach soil, dry weight, N content and N-uptake of tomatoes. The 30 tons/ha mixture of cow manure and kirinyuh (*Chromolaena odorata*) that applied at 20 days before planting affects the highest N-uptake of tomatoes.

REFERENCES

- [1] Winarso, S. (2003). *Kesuburan Tanah Dasar Kesehatan Dan Kualitas Tanah [Soil Fertility Basic Health and Soil Quality]*. Jember: Gava Media. 189 hlm.
- [2] Adil, W. H., Sunarlim, N., & dan Roostika, I. (2005). Pengaruh Tiga Jenis Pupuk Nitrogen terhadap Tanaman Sayuran [Effect of Three Types of Nitrogen Fertilizers on Vegetable Plants]. *Biodiversitas*, 7 (1), 77-80.
- [3] Lakitan, B. (2007). *Dasar-dasar Fisiologi Tumbuhan [Fundamentals of Plant Physiology]*. Jakarta: PT Raja Grafindo Persada. Anonymous. 2009. *Data Produksi*

- Tomat [Tomato Production Data]. <http://www.bps.go.id>. Downloaded on January 9, 2018. Central Statistics Office.
- [4] Adiyoga, W., Suherman, R. & Soetiarso, T. A. (2004). *Laporan Akhir Profil Komoditas Tomat [Final Report on Tomato Commodity Profile]*. Project/Project Participatory Agricultural Technology Assessment Project (PPATP), Research Center.
- [5] Susila, A. D. (2006). *Panduan Budidaya Tanaman Sayuran [Guide on Vegetable Cultivation]*. Department of Agronomy and Horticulture. Bogor Agricultural Institute. Page 115. Accessed on 4 February 2018.
- [6] Darmawijaya, M. I. (1997). *Klasifikasi Tanah Dasar Teori Bagi Penelitian Tanah dan Pelaksanaan Pertanian di Indonesia [Classification of Soil Theories for Soil Research and Agricultural Implementation in Indonesia]*. Gadjah Mada University Press. Yogyakarta.
- [7] Anonymous. (2009). *Data Produksi Tomat [Tomato Production Data]*. Central Statistics Bureau.
- [8] Budiyanto, G. (2009). *Bahan Organik dan Pengelolaan Nitrogen Lahan Pasir [Organic Materials and Nitrogen Management Sand Land]*. Unpad Press.
- [9] Sunardi. & Sarjono, Y. (2007). Penentuan Kandungan Unsur Makro Pada Lahan Pasir Pantai Samas Bantul Dengan Metode Analisis Aktivasi Neutron (AAN) [Determination of the Macro Element Content in the Samas Bantul Coastal Land by the Neutron Activation Analysis (AAN) Method]. *Proceeding PPI-PDIPTN 2007 BATAN* p: 123-129.
- [10] Anonymous. (2007). *Panduan Lengkap Budi Daya Tomat [Complete Guide to Tomato Cultivation]*. Agromedia. Jakarta. Agromedia Editor
- [11] Syekhfani. (2000). The Significance of Organic Materials for Soil Fertility, *Journal of Organic Fertilizer Research*.
- [12] Wardhani. (2006). *Aplikasi Mulsa Chromolaena odorata dan Cendawan Mikoriza arbuskular pada Tanah Latosol untuk Pertumbuhan dan Produksi Preuraria javanica [Application of Mulch Chromolaena odorata and Arbuskular mycorrhizal fungi on Latosol soil for growth and production of Preuraria javanica]*. Thesis of the Faculty of Animal Husbandry, Bogor Agricultural University. Accessed on 16 November 2017.
- [13] Nurhajati, H. (1982). *Pengaruh Pemberian Pupuk Hijau dan Kapur pada Tanah Podsolik Merah Kuning terhadap Ketersediaan Fosfor dan Produksi Tanaman Jagung [The Effect of Provision of Green and Lime Fertilizer on Red Yellow Podsolic Soils on Phosphorus Availability and Corn Production]*. Doctoral Dissertation. Graduate School of Undergraduate. Bogor: Bogor Agricultural University.
- [14] Suntoro, Syekhfani, Handayanto, E., & Sumarno. (1998). *Penggunaan Bahan Pangkasan Krinyuh (Chromolaena odorata) untuk meningkatkan ketersediaan P, K, Ca dan Mg pada Oxic Dystrudepth di Jumapolo, Karanganyar, Jawa Tengah [The use of Kirinyuh Pruning Materials (Chromolaena odorata) to increase the availability of P, K, Ca dan Mg at Oxic Dystrudepth in Jumapolo, Karanganyar, Central Java]*. *BIOMA*, 13(1), 20 – 26.
- [15] Hadi, M., Hidayat, W., & Baskoro, K. (2000). Pembuatan Kertas Anti Rayap Ramah Lingkungan dengan Memanfaatkan Ekstrak Daun Kirinyuh (*Chromolaena odorata* L.) [Making Environmentally Friendly Anti-Termite Paper Using Extracts of Kirinyuh leaf (*Chromolaena odorata* L.)]. *BIOMA*, 6(2), 12-18.


- [16] Handayanto, Nuraini, E. Y. & Syekhfani. (2000). *Stimulasi Dekomposisi Akibat Penambahan Bahan Organik Baru [Decomposition Stimulation Due to Addition of New Organic Materials]*. Proceedings of the National Congress VII. BLACK. Bandung.
- [17] Anonim. (2009). *Analisa Kimia Tanah, Tanaman, Air dan Pupuk [Chemical Analysis of Soil, Plants, Water and Fertilizer]*. Technical Guidelines. Soil Research Center. Center for Agricultural Research and Development. Agriculture Department.
- [18] Gomez, K. A. & Gomez, A. A. (1995). *Prosedur Statistik untuk Penelitian Pertanian [Statistical Procedures for Agricultural Research]*. Jakarta: UI Press.
- [19] Syukur & Harsono. (2008). Pengaruh Pemberian Pupuk Kandang dan NPK terhadap Beberapa Sifat Kimia dan Fisika Tanah Pasir Pantai Samas Bantul [Effect of Applying Manure and NPK on Some Chemical and Physical Properties of The Samas Bantul Beach Sand Soil]. *Jurnal Ilmu Tanah dan Lingkungan*, 8, 138-145.

ABOUT THE EDITOR


Wayan Suparta

(Chief Editor)


Associate Professor
University of Malaya



Wayan Suparta, PhD has experience as a high school teacher, lecturer, and researcher for over 25 years. His latest position is Associate Professor at University of Malaya, Kuala Lumpur, Malaysia, where previously it was a principal lecturer in the Department of Informatics, Universitas Pembangunan Jaya Tangerang, Indonesia (6 Feb 2019 – 5 Feb 2020), lecturer at the Dept of Electrical Engineering, Sanata Dharma University, Yogyakarta (Feb 2018 - Jul 2018). Previously, he was an Adjunct Professor in the Department of Civil Engineering at the Yogyakarta University of Technology (Sep 2017 - Jan 2018). He is also an associate faculty member at the Doctor of Computer Science (DCS), Bina Nusantara University (BINUS) Jakarta. Before returning to Indonesia, he worked at Universiti Kebangsaan Malaysia (UKM) as a lecturer and researcher from the rank of Senior Lecturer to Associate Professor from July 2008 to April 2017. He began his career as a teacher of Physics, Electronics, and Computers in high school from SMUK Cor Jesu Malang (1994-1997), and a teacher at SMUK Santo Aloysius Bandung (1997-2000). Appointed as a Lecturer in Electrical Engineering at College Legenda Group, Malaysia (2000-2004). After get a Ph.D. in March 2008, he was appointed as a Post-Doctoral Fellow (2007-2008), subsequently appointed as a Senior Lecturer (1 July 2008 - 3 April 2012). His teaching area is electronics/electrical engineering, informatics, physics, and statistics, while his research is focused on GPS/GNSS Technology, Communication Systems (Remote Sensing), Smart Systems, and Artificial Neural Network and Data Mining Applications for Meteorology and Climate Change.

MEMBERS OF THE EDITORIAL TEAM

Johannes Hamonangan Siregar, PhD is currently a lecturer of Information Systems Program Study at Universitas Pembangunan Jaya, Tangerang, Indonesia. He is also Scientific Coordinator for Information Systems, member of the Board of University Senate and the Board of Widayakala University Journal. He teaches Foundations of Mathematical Logic, Database Systems, Knowledge Management, Information Security and Network Administration. He considers his students to find smart solutions for Information System challenges that can be overcome through understanding mathematics. In the past, he was mathematics lecturer at Surya College of Education, Indonesia in 2013. He also has work experience with Japanese industry. In 2006, he was a system analyst to develop mobile phone at NEC Company Projects. Then he joined the University of Tsukuba e-learning project in 2010. Based on his educational and industry experiences, he has research interest in algorithms for database systems, mathematical modelling in information systems and mathematics education.



Nur Uddin, PhD is an assistant professor at Department of Informatics, Universitas Pembangunan Jaya in Tangerang Selatan, Indonesia. He received PhD in Engineering Cybernetics from Norwegian University of Science and Technology (NTNU) in Trondheim, Norway in 2016, master degree in Mechanical Engineering from Gyeongsang National University (GNU) in Jinju, South Korea in 2009, and bachelor degree in Aeronautics Engineering from Institut Teknologi Bandung (ITB) in Bandung, Indonesia in 2002. He has five years industrial work experiences in an oil service company and airlines. He began academic carrier in 2017. His research areas are in system dynamics, control system, robotics, artificial intelligence, and computer vision. His current research is autonomous mobile robots.



Hendy Tannady, PhD - His career as practitioner began in Quality Engineering division at mining shoes manufacturer in 2008, then he worked in Production Planning Inventory Control at one of seafood processing industry and became a Market Research Analyst at Binus university in 2009. Career in academic began in 2010 as Head of the Integrated Industrial Engineering Laboratory at Binus University. During his work at Binus University, he was actively involved in preparing the process of re-accreditation B to A and ABET international accreditation. Since 2013 to June 2019, he was trusted by Universitas Bunda Mulia as Head of Industrial Engineering Study Program, concurrently he was Manager of the Research Department (February 2018 – January 2019). Since July 2019, he has been trusted as the Dean of Faculty of Humanities and Business at Universitas Pembangunan Jaya. His research is focusing on Operations Management, Management Science, Industrial Psychology and Organizational Behaviour.



Yohanes Totok Suyoto, PhD is a management lecture and researcher at Universitas Pembangunan Jaya, South Tangerang, Indonesia. He has also served as a Head of Management Department in the university for almost two years (August 2, 2018 – present) from which he undertook his passion on researching in Marketing. Previously, he was a lecture at Pelita Harapan University, Surabaya, for eight years (August 2010 – July 2018) and Widya Mandala Catholic University, Surabaya, for six years (September 2003 – January 2009).

Since 2012, he also served as a member of professional association, Indonesia Marketing Association/IMA. He obtained his Ph.D in Management science from Airlangga University, Surabaya. As a senior lecture, he has given lectures in Management areas, e.g., Strategic Management, Change Management, Marketing Management, Strategic Marketing, Global Marketing, Consumer Behavior, Creative Management and Customer Value.



Frederik Josep Putuhena, PhD is a full professor and began his professional carrier as Tutor at University of Indonesia, and promoted as lecturer for Water Resources Engineering in 1976. He received British Council Award for study at University of Newcastle upon Tyne, UK, and received Master of Science in Engineering Hydrology. His experience as Water Resources Planner was gained, while conducting various river basins planning projects in Indonesia. In 1991, he received award from Indonesia Planning Board, for Doctoral study in SUNY at Buffalo, USA. Later, he taught at Pancasila

University, Jakarta. Also, assigned as Coordinator for Capacity Building in Water Resources Sector in Indonesia (1997 – 2000). Then (2003 – 2016) he was assigned as lecturer in the Faculty of Engineering, Universiti of Malaysia Sarawak (UNIMAS). Currently, he is teaching at Universitas Pembangunan Jaya, Jakarta. Overall, the courses cover Numerical Analysis, Statistics, Hydrology, Hydraulics, Urban Drainage, Water Resources Engineering, Hydraulic Structures, and Engineering Economics.

As Professor at the Universitas Pembangunan Jaya, with daily activities in lecturing, doing research, as well as water resources development planning, I really praise Nova Science Publishers for publishing selected papers from "2020 International Conference on Urban Sustainability, Environment, and Engineering (CUSME 2020)". This publication would be useful for professionals, researchers, scholar, policymakers, and NGO. I believe that currently, many professionals would like to give more attention to the development of sustainable urban. In addition, this publication could be used as a reference for city authorities to make appropriate policy choices to protect the provision of equitable housing, health, and transportation services.

Prof. Ir. Frederik Josep Putuhena M.Sc., Ph.D
Center for Urban Studies - Universitas Pembangunan Jaya

Urban Development and Lifestyle are trend issues for cities around the world. Learning from experience is the most effective way to support cities to be sustainable developed. This book offers knowledge sharing among countries and covers a variety of cities' issues. It also provides great lessons for researchers, officers and policy-makers who are coping with several urban problems.

Associate Professor Sarintip Tantanee, Ph.D.
Director

Center of Excellence on Energy Technology and Environment (CETE)
Faculty of Engineering, Naresuan University, Thailand


nova
science publishers

www.novapublishers.com

ISBN 978-1-53618-560-7



9 781536 185607