

Dwi Fitri Yudiantoro <d_fitriyudiantoro@upnyk.ac.id>

[JGEET] Article Review Request

1 pesan

JGEET (J. Geoscience Eng. Environ. Technol.) <admin@journal.uir.ac.id>
Kepada: Dwi Fitri Yudiantoro <d_fitriyudiantoro@upnyk.ac.id>

7 Desember 2023 pukul 08.45

Dwi Fitri Yudiantoro:

I believe that you would serve as an excellent reviewer of the manuscript, "Petrology and Mineral Chemistry of Ultramafic Rock in Baula - Pomalaa Ophiolite Complex, Southeast Sulawesi," which has been submitted to Journal of Geoscience, Engineering, Environment, and Technology. The submission's abstract is inserted below, and I hope that you will consider undertaking this important task for us.

Please log into the journal web site by 2024-01-03 to indicate whether you will undertake the review or not, as well as to access the submission and to record your review and recommendation.

The review itself is due 2024-01-03.

Submission URL: <https://journal.uir.ac.id/index.php/JGEET/reviewer/submission?submissionId=14491&reviewId=7586&key=pHvUDw>

Thank you for considering this request.

JGEET (J. Geoscience Eng. Environ. Technol.)
Department of Geological Engineering, Faculty of Engineering, Universitas Islam Riau
jgeet@journal.uir.ac.id

"Petrology and Mineral Chemistry of Ultramafic Rock in Baula - Pomalaa Ophiolite Complex, Southeast Sulawesi"

Ultramafic rocks compose the Baula and Pomalaa Ophiolitic Complexes. Peridotite, composed primarily of olivine and pyroxene minerals, dominates ultramafic rocks. Chemical and petrological research has focused on minerals, such as olivine, pyroxene, and spinel. This study examines the tectonic setting and temperature of ultramafic rock formation. Twelve ultramafic rock samples were examined using geothermometers made of pyroxene, petrographic examination, and coexisting olivine and spinel analyses. SEM and petrographic analysis of pyroxene lamellae and mylonite-ultramylonite structures allowed for the measurement of the geothermometer of ultramafic rocks. Using SEM-EDS, the coexistence of olivine and spinel was analyzed to determine the type of ultramafic tectonic setting. In the coexistence of olivine and spinel, olivine and spinel oxide compounds as tectonic setting markers in the form of Fo and Cr# values. Ultramafic rocks have different temperature levels and the first one starts at a high temperature of 1000-1200°C. It is also characterized by thin, elongated lamellae. Instead, large lamellae characterize orthopyroxene at medium temperatures (800–1000°C). The presence of ultramylonite structures likewise characterizes this phase. Lamella irregular, anhedral, and broader forms are typical of low temperatures (500–800°C). The presence of mylonite structures can also distinguish this phase. The distribution of #Fo ranged from 0.87 to 0.92, and Cr# values ranged from 0.13-0.19. According to coexisting olivine and spinel analysis. On the Olivine-Spinel Mantle Array (OSMA), the Fo and Cr# plot indicates that the peridotites tectonic setting was from the ocean floor and the magmatism was from MORB (Mid Oceanic Ridge Basalt).

JOURNAL OF GEOSCIENCE, ENGINEERING, ENVIRONMENT, AND TECHNOLOGY

Department of Geological Engineering

Jl. Kaharuddin Nasution No 113 Perhentian Marpoyan, Pekanbaru, Riau 28284

e-mail : jgeet@journal.uir.ac.id, web : <http://journal.uir.ac.id/index.php/JGEET>

RESEARCH ARTICLE

Petrology and Mineral Chemistry of Ultramafic Rock in Baula -
Pomalaa Ophiolite Complex, Southeast Sulawesi

Rio Irhan Mais Cendrajaya¹, Laode Ihksan Juarsan², Masri^{1,*}, Al Rubaiyn², Syahrul³, Neni¹,
Suci Ramadani¹, Hasria¹

¹ Geological Engineering, Halu Oleo University, Kendari, Indonesia

² Geophysical Engineering, Halu Oleo University, Kendari, Indonesia

³ Mining Engineering, Universitas Sembilanbelas November, Kolaka, Indonesia

* Corresponding author: masri@uho.ac.id
Tel.: +6282-296-451-988
Received: Oct 1, 2016; Accepted: Nov 20, 2016.
DOI: 10.24273/jgeet.2016.1.2.001

Abstract

Ultramafic rocks compose the Baula and Pomalaa Ophiolitic Complexes. Peridotite, composed primarily of olivine and pyroxene minerals, dominates ultramafic rocks. Chemical and petrological research has focused on minerals, such as olivine, pyroxene, and spinel. This study examines the tectonic setting and temperature of ultramafic rock formation. Twelve ultramafic rock samples were examined using geothermometers made of pyroxene, petrographic examination, and coexisting olivine and spinel analyses. SEM and petrographic analysis of pyroxene lamellae and mylonite-ultramylonite structures allowed for the measurement of the geothermometer of ultramafic rocks. Using SEM-EDS, the coexistence of olivine and spinel was analyzed to determine the type of ultramafic tectonic setting. In the coexistence of olivine and spinel, olivine and spinel oxide compounds as tectonic setting markers in the form of Fo and Cr# values. Ultramafic rocks have different temperature levels and the first one starts at a high temperature of 1000-1200°C. It is also characterized by thin, elongated lamellae. Instead, large lamellae characterize orthopyroxene at medium temperatures (800-1000°C). The presence of ultramylonite structures likewise characterizes this phase. Lamella irregular, anhedral, and broader forms are typical of low temperatures (500-800°C). The presence of mylonite structures can also distinguish this phase. The distribution of #Fo ranged from 0.87 to 0.92, and Cr# values ranged from 0.13-0.19. According to coexisting olivine and spinel analysis. On the Olivine-Spinel Mantle Array (OSMA), the Fo and Cr# plot indicates that the peridotites tectonic setting was from the ocean floor and the magmatism was from MORB (Mid Oceanic Ridge Basalt).

Keywords: pyroxene lamella, geothermometer, olivin-spinel coexisting, MORB peridotite, Kolaka

1 Introduction

In Indonesia, Ophiolite complexes are located in Meratus Mountains, eastern and southeastern Sulawesi, Halmahera, and Papua. Some ophiolite complexes are associated with melange, such as in Ciletuh (West Java), Karangsambung (Central Java), and Bantimala (Sulawesi) (Suroño and Hartono, 2013). The ophiolite complex on Sulawesi Island is known as the East Sulawesi Ophiolite Belt (ESOB) or the East Sulawesi Ophiolite Belt (LOST). The LOST extends for 500 km along the eastern arm to the southeastern arm (Parkinson, 1998). The LOST can be traced from the East Arm of Sulawesi, starting from Poh, Bunta, to Ampana and Morowali. In the central part of Sulawesi, LOST is distributed in Kolonodale, Bungku, and Kendari. The west coast of the Southeast Arm of Sulawesi, such as Lasusua, Kolaka, and Kabaena (Kadariusman et al., 2004, Hamilton, 1979).

The ophiolites of eastern Sulawesi are Late Cretaceous-Eocene in age and were displaced during the Oligocene-Miocene period by the Sula Spurs collision (Parkinson, 1998). The ophiolite was displaced along a back-arc basin shear fault (Monnier et al., 1995) with an obduction mechanism (Husein et al., 2014). The ophiolite sequence consists of peridotite cumulates, microgabbro, sheeted dolerite, and MORB basaltic (Kadariusman et al., 2004). Peridotites are dunite, harzburgite, lherzolite and

pyroxenite, while microgabbro and basalt are only found locally in a few places (Suroño, 2013). REE element studies on peridotite clinopyroxene indicate a mid-ocean ridge (MOR) origin rather than a suprasubduction zone. (Kadariusman et al., 2004).

The displacement of ultramafic rocks is important for the study of petrology and petrogenesis because mineralogical parameters related to changes in temperature and pressure during the rock formation process (geothermometer) and displacement can tell us about the early genesis of peridotite rock formation and its tectonic evolution (Lindsley and Andersen, 2012). The formation temperature of ultramafic rocks can be identified based on pyroxene minerals that exhibit lamellar exsolution textures between clinopyroxene and orthopyroxene (Yellappa et al., 2021, Koizumi et al., 2014, Lindsley, 1983). The exsolution process usually occurs due to lamellar growth in the crystal of origin during the cooling process. The determination of the tectonic environment of ultramafic rocks is based on the presence of coexisting olivine and spinel minerals (Olfindo et al., 2020, Payot et al., 2018, Arai, 1994).

The research location for the study of petrology and mineral chemistry of ultramafic rocks is Pomalaa and Baula, Kolaka Regency, Southeast Sulawesi Province. In this area, there are widely exposed ultramafic rocks, which have

Commented [S1]: Petrochemistry of Ultramafic Rock in Ophiolite Complex...

Commented [S2]: yang diukur suhu terbentuknya batuan beku atau proses termetamorfkan ? atau pensesaran juga ubahan ?

Commented [S3]: Lebih dispesifikan termasuk jenis ofiolit yang mana ? . ESOB atau LOST. sehingga ofiolit daerah penelitian jelas letak tatanan tektoniknya. Juga karakteristik ofiolitnya dibandingkan dengan ofiolit yang lain.

generally been serpentinized (Cendrajaya, 2017, Simandjuntak et al., 1993). The absence of relevant research related to the mineral chemistry and petrology of ultramafic rocks in this area is the basis for proposing research.

In this study, the formation temperature of ultramafic rocks was analyzed by making petrographic observations and analyzing pyroxene lamellae (geothermometer) using SEM. This data was then combined with the results of chemical analysis of coexisting olivine-spinel to obtain an overview of the tectonic environment of the study area.

2 Methodology

A descriptive approach to field geology methods, supported by laboratory analysis, is used to conduct this research. The percentage of mineral composition in petrographic observations has been the subject of quantitative research. Megascopically and microscopically detailed descriptions of ultramafic rock samples serve as primary data.

Fresh ultramafic rocks in the study area are the focus of research. The object was then concentrated on the orthopyroxene-clinopyroxene minerals, which form lamellar exsolution textures, as well as the coexisting olivine and spinel minerals. Thin sections of ultramafic rocks show the presence of these textures. The microscopic description of ultramafic rocks is based on the work of Streckeisen (1976).

The petrographic analysis uses Nikon Eclipse E-100 polarizing microscope at the Geological Engineering Laboratory of Halu Oleo University, with thin sections prepared at the Petrology Laboratory of Hasanuddin University, Makassar. Observed optical properties include crystal shape and size, cleavage, relief, index of refraction, absorbs and pleochroic color, interference color, optical orientation, twinning, and extinguishing angle. This description seeks to identify various characteristics of the olivine and pyroxene minerals present in the thin section, such as specific mineral types, distinctive textures, and other distinguishing characteristics.

If two pyroxene minerals are observed coexisting and forming lamellar exsolution, the formation temperature (geothermometer) of ultramafic rocks can be determined. The two pyroxene minerals analyzed contain clinopyroxene lamellae within an orthopyroxene mineral host. SEM and polarizing microscopy were used to examine these two pyroxene minerals. The formation temperature of ultramafic rocks requires a descriptive analysis based on the texture and chemical composition of lamellar pyroxene.

A Phenom ProX SEM equipped with an EDS detector was used to determine the mineral's chemical composition. Utilizing the EDS detector, chemical data on minerals was gathered. With 15 kV of accelerating voltage and 30 mA of probe current, mineral chemistry tests were conducted in a high vacuum environment. There are sixteen firing points (beams), each of which is composed of a lamellar-forming orthopyroxene-clinopyroxene mineral and a coexisting olivine-spinel mineral. The elements measured include Oxygen (O), Magnesium (Mg), Silicon (Si), Iron (Fe), Nickel (Ni), Calcium (Ca), Aluminum (Al), Manganese (Mn), Chromium (Cr), Titanium (Ti), Sodium (Na), Potassium (K), Zinc (Zn), and Vanadium (V) (V). The chemical data is presented as oxide compounds with weight percent (wt percent) units, including SiO₂, MgO, FeO, NiO, CaO, MnO, Al₂O₃, Cr₂O₃, TiO₂, K₂O, Na₂O, ZnO, and V₂O₃. The weight percent (wt percent) data were analyzed and processed in order to determine the mole percent (percent mol) of Mg

(percent En), Fe (percent Fs), Ca (percent Wo), Cr#, Mg#, and others.

Together-growing olivine and spinel minerals (coexisting) were used to determine the tectonic environment of ultramafic rocks in the study area. Using SEM-EDS analysis, chemical data were obtained based on the work of Arai (1994).

3 Geology

3.1 Geomorphology

The geomorphology of the Baula-Pomalaa area, Kolaka Regency, consists of hills, valleys, and lowlands at 0-800 meters above sea level elevation. Based on van Zuidam (1986) classification, the study area is divided into three geomorphological units, i.e. structural morphology (55%), which is spread in the central and northeast-southeast part of the study area, denudational morphology (40%) which is distributed in the west-southwest part of the study area, and fluvial system (5%) which is spread in the northwest part of the study area. River systems consist of rectangular and dendritic patterns. Rectangular type is found almost throughout the study area, with igneous and metamorphic rocks. The dendritic flow pattern is located southwest to northwest of the study area, consisting of conglomerate rocks and alluvial deposits.

3.2 Stratigraphy

The stratigraphy of the Baula-Pomalaa area comprises ultramafic rocks, metamorphic rocks, monomic conglomerates, polyimic conglomerates, and alluvial deposits. Ultramafic rocks consist of peridotite. The constituent minerals of ultramafic rocks are olivine, pyroxene, serpentine, and opaque minerals. Both ultramafic and metamorphic rocks are separated by structural contact of fault. In general, ultramafic rocks have undergone partial or complete serpentinization and have been weathered into laterite deposits. Ultramafic rocks occupy ~40% of the study area (Figure 1).

Metamorphic rocks in the study area consist of serpentinite and schist. Serpentinities, alteration rocks of ultramafic rocks, are composed of serpentine minerals such as lizardite and chrysotile. Muscovite-quartz schists dominate the schists in the study area and belong to the greenschist facies. The schists are scattered in the eastern part of the study area.

Monomic conglomerates are commonly found in intra-mountain depressions and rivers with >10m fault scarp. The grain fragments are dominated by ultramafic rocks and serpentinite (>90%). The grain size is sand-boulder (>30 cm), dark brown-brown in color, non-carbonate cement, and poor sorted and generally distributed in the southern part and slightly in the north of the study area.

The polyimic conglomerate is found in low-elevation areas in the south-southwest and slightly north of the study area. Its grain consists of ultramafic, schist, serpentinite, and limestone lithics with boulder-sand size (<30 cm), light grey-brown, non-carbonate cement, and poorly sorted. In some locations, these rocks have not consolidated well.

Alluvial deposits are found in the north and along the coastline of the study area. This area is generally for residential use. The alluvium unit consists of brownish-grey to reddish clay-boulder loose material.

Commented [S4]: Sebaiknya menggunakan kata positif bukan menggunakan kata negative (BELUM)

Commented [S5]: serpentin bukan termasuk mineral penyusun uktra basa, tetapi mineral ubahan. atau mineral metamorf. atau serpentinisasi

Commented [S6]: sudah disebut batuan alterasi. Dijelaskan secara jelas geologi dari batuan ultra basa nya. Batuan ultra basa atau metamorf atau batuan ubahan.

Commented [S7]: bagaimana hubungan batuan ultrabasa, metamorf dan konglomerat secara geologi




Figure 1 Geologic map and distribution of ultramafic rock sample in the study area

3.3 Geological Structure

Geological structures that develop in the study area are strike-slip and thrust faults (Figure 1). The regional tectonics influences the formation of the geological structure of the study area in Sulawesi, which has the direction of the regional structure northwest-southeast. Thrust faults are generally northwest-south-southeast and northeast-southwest, and strike-slip faults are northwest-southeast and northeast-southwest. The formation of

Table 1 Representation of primary and secondary minerals from 12 thin sections of ultramafic rocks. Mineral symbols based on Whitney and Evans (2009). Superscript 'R' = relic, 'X' = lamellar exsolution

Sample	Main Minerals	Metamorphic/retrograde Minerals	Lithology
BDM-01	Ol ^R + Opx + Cpx	Srp + Tlc	Lerzolit
EWO-01	Ol ^R + Opx + Cpx + Cr-Spl	Srp + Tlc + Mag	Harzburgit
PSRD-01	Ol ^R + Opx + Cpx	Srp + Chl + Mag	Olivin Websterit
HUKO-H16	Ol + Opx + Cpx ^X + Cr-Spl	Srp + Chl + Tr	Lerzolit
HUKO-H11	Ol ^R + Opx + Cpx	Srp + Tlc + Chl	Lerzolit
PMS-01	Ol + Opx + Cpx ^X + Cr-Spl	Srp + Chl + Tr	Harzburgit
SOP-H12	Ol + Opx + Cpx + Cr-Spl	Srp + Chl	Harzburgit
SOP-H6	Ol ^R + Opx + Cpx + Cr-Spl	Srp + Tlc + Mag	Lerzolit
VALE-H14	Ol + Opx + Cpx + Cr-Spl	Srp + Chl + Mag	Lerzolit
VALE-H18	Ol ^R + Opx + Cpx ^X + Cr-Spl	Srp + Tlc + Chl	Lerzolit
VALE-H6	Ol ^R + Opx + Cpx + Cr-Spl	Srp + Tlc + Mag	Lerzolit
VALE-02	Ol + Opx + Cpx + Cr-Spl	Srp + Chl + Tlc	Olivin Websterit

Harzburgite (EWO 01, PMS 01, and SOP H12) is generally composed of olivine (45-70%), orthopyroxene (20-40%), spinel (<5%), and clinopyroxene (5-15%). Harzburgite has generally undergone partial serpentinization (Figure 2a and 2d). Orthopyroxene is present as 1-5 mm subhedral-anhedral crystals. The center of orthopyroxene crystals is generally serpentinized, while the rim, usually in contact with olivine, is altered to tremolite and talc (Figure 2a). Clinopyroxene is present as lamellar exsolution alongside orthopyroxene. The spinel present generally coexisting with olivine and is of the spinel chromite type. Olivine typically has euhedral-anhedral crystals measuring 0.2-3 mm, altered to serpentine, forming vein and mesh textures (Figure 2d). At some contacts between olivine and orthopyroxene, olivine is also present as monilite and ultramylonite (Figure 2a). Ultramylonites indicate an intensive deformation process in the shear zone (Matsiyak and Trepman, 2015). The magnetite present is

structures in the study area is closely related to the collision direction between Southeast Sulawesi and the Buton Microcontinent Fragment which occurred during the Early Miocene and continued until the Late Miocene.

4 Results

4.1 Petrography of Ultramafic Rocks

The ultramafic rocks in the Baula Pomalaa Ofiolite Complex consist of harzburgite, lherzolite and olivine websterite, mostly serpentinized. Ultramafic rocks are structurally adjacent to metamorphic rocks; in some locations, the ultramafic rocks are fully serpentinized. Conglomerates are divided into polyimic and monomic conglomerates, with a predominance of ultramafic grains or fragments. Monomic conglomerates are commonly found in intra-mountain depressions and rivers with >10m inclines. The grain fragments are dominated by ultramafic and serpentinite rocks (>90%). Polyimic conglomerates are found in low elevation areas with ultramafic, schist, serpentinite and limestone grains.

Based on petrographic analysis of 12 thin sections, ultramafic rocks can be grouped into three types. Harzburgite was identified in samples SOP-H12, EWO-01, and PMS-01, lherzolite was identified in samples BDM-01, HUKO-H11, HUKO-H16, SOP-H6, VALE-H14, VALE-H18, and VALE-H6, while olivine websterite was identified in samples PSRD-01 and VALE-02. Generally, olivine is present both as relicts and in subhedral crystal. In addition to primary minerals, secondary minerals characteristic of retrograde metamorphism are also present, characterized by the abundance of serpentine, talc, and chlorite (Table 1).

a by-product of the serpentinization process (Maulana et al., 2015).

Lherzolites (HUKO H16 and VALE H14) are generally composed of olivine (40-50%), orthopyroxene (20-30%), clinopyroxene (25-30%), and spinel (<5%). The serpentinization process of lherzolite in the study area is more intensive than that of harzburgite (Figure 2f). Orthopyroxene is present as subhedral-anhedral crystals of 0.5-5 mm in size. Tremolite-talc and olivine ultramylonite are present at the rim of orthopyroxene in contact with olivine (Figure 2b). One of the orthopyroxene crystals was also found to have a kinkband texture, and some had undulate extinction (Figure 2b). These textures are characteristic of rigid/brittle deformation (Kadarusman et al., 2004). Clinopyroxene is present as subhedral-anhedral crystals of 1-5 mm size and as lamellar exsolution within the host mineral orthopyroxene (Figure 2b and 2e). Hornblende was found to start partially replacing

Commented [S9]: agar disebutkan pada sari

Commented [S10]: akibat pelapukan atau termetamorfkan ?

Commented [S11]: secara regional polimik. Jelaskan bila monomik ?

Commented [S8]: Legend batuan tidak jelas

Commented [S12]: pembagian ini berdasarkan apa ? dijelaskan

Commented [S13]: ditinjau lagi...apakah batuan ultra basa atau metamorf ?

Commented [S14]: bila batuan sudah termetamorfkan apakah namanya masih batuan beku ? dan seterusnya

Commented [S15]: termetamorfkan ?

Commented [S16]: sebaiknya ortopirosken langsung disebut hipersten, klinopirosken juga langsung disebutkan jenis mineralnya.

Commented [S19]: mineral ini bukan kelompok ultra basa. Bagaimanakan menjelaskannya, terkait suhu dan proses geologinya

Commented [S17]: mengapa yang diubah hanya olivine saja ?

Commented [S18]: monilite adalah mineral hasil dari proses pensesaran. Apa arti dari batuan ini ? masih tetap ultra basa atau metamorf hingga juga pensesaran.

clinopyroxene (Figure 2c). Spinel chromite type is found generally coexisting with olivine (Figure 2c). Olivine typically has euhedral-anhedral crystals measuring 0.1-3 mm, altered to serpentine and chlorite, forming vein and mesh textures (Figure 2c and 2f).

Olivine websterites (PSRD 01 and VALE 02) are generally composed of olivine (15-40%), orthopyroxene (35-50%), clinopyroxene (20-30%), and spinel (<5%). Enstatite-type orthopyroxene is present predominantly in this rock compared to clinopyroxene (Figure 2h). Orthopyroxene is present as subhedral-anhedral crystals measuring 0.5-5 mm. Serpentine and talc are commonly present in the rim of orthopyroxene as alteration minerals. Some orthopyroxene crystals are also indicated as protomylonite characterized by undulate extinction (Figure 2g). Clinopyroxene is a 0.2-2 mm subhedral-anhedral crystal and lamellar exsolution within host orthopyroxene mineral (Figure 2g). Spinel, identified as spinel chromite, is present and coexisting with orthopyroxene and olivine (Figure 2h). Generally, orthopyroxene coexisting with >2mm spinel. Olivine has typically 0.1-3 mm in size of euhedral-anhedral crystals, serpentinized and chloritized, forming vein and mesh textures (Figure 2h).

The microstructures present in the ultramafic rock samples consist of milonite, ultramylonite, and kinkband (Figure 2a, 2b, and 2g). The process that allows these structures is deformation control (Matysiak and Trepmann, 2015). The olivine that forms mylonite and ultramylonite are interpreted as the result of the recrystallization process and the previous olivine minerals during the deformation process. The difference between milonite and ultramylonite can be seen in their crystal size. The milonite present can be seen in Figure 2g. The olivine ultramylonite is generally in the rim between olivine and pyroxene, and some of the olivine is altered (Figure 2a and 2b). Some of the orthopyroxene crystals in the sample were identified as kinkbands, especially pyroxene crystals with dimensions >1mm. The orthopyroxene crystals (Figure 2b), which underwent lamellar exsolution by clinopyroxene, appear to have a kinkband structure. This kinkband structures characterizes a brittle deformation

process (Burnley et al., 2013, Drue, 2011, Kadarusman et al., 2004).

5 Mineral Chemistry of Ultramafic Rocks

5.1 Olivin

Olivine mineral chemistry data (4 oxygen) were normalised, assuming all Fe is Fe²⁺. All olivine minerals analyzed were forsterite ([Fe,Mg]₂SiO₄) with Fo contents varying from 0.87 - 0.92 (Table 2).

5.2 Orthopyroxene

The chemical data of orthopyroxene minerals in ultramafic rocks of the Baula and Pomalaa Regional Ophiolite Complex can be seen in Table 3. The orthopyroxene mineral chemistry data is normalized (6 oxygen) with a total cation of 4 apfu. Orthopyroxene is present in all ultramafic rock incisions and is of the Mg-rich enstatite type with compositions En_{87,1-88,5}, Fs_{9,1-10,8}, Wo_{2,05-2,9}. X_{Mg} range ~0.92.

5.3 Clinopyroxene

The chemical data of clinopyroxene minerals of ultramafic rocks of the Ophiolite Complex of Baula and Pomalaa Regions can be seen in Table 3. The mineral chemistry data of orthopyroxene is normalized (6 oxygen) with a total cation of 4 apfu. Mg-rich clinopyroxene is present in some ultramafic rock incisions and is predominantly of the augite and diopsid types. Augite has a composition of En_{50,8-55,2}, Fs_{4,6-5}, Wo_{38,6-44,1} with X_{Mg} ~0.92. Diopsids have compositions En_{47,08}, Fs_{4,66}, Wo_{48,26} with X_{Mg} ~0.91.

5.4 Chromite-spinel

The chemical data of spinel mineral of ultramafic rock of the Ophiolite Complex of Baula and Pomalaa Area can be seen in Table 2. The spinel mineral chemistry data (4 oxygen) is normalized with a total cation of 3 apfu. All spinel minerals analyzed at the study sites are chromite-spinel type ([Mg,Fe][Cr,Al]₂O₄) with Cr# <0.2.

Commented [S20]: ultra basa sudah mengalami proses geologi yang lain

Commented [S21]: belum pernah disebutkan pada diskripsi sebelumnya




Figure 2 (a) Microphotography of harzburgite (PMS 01) under cross-polarization, showing serpentinized olivine (ol) and orthopyroxene (opx). Ultramylonite (uml) olivine is present on the rim of orthopyroxene. Alterations of orthopyroxene, tremolite (tr) and talc (tlc) are also present in the rim of orthopyroxene (opx). (b) Orthopyroxene (opx) on lherzolite (HUKO H16) as host mineral shows a *kinkband* texture (due to brittle deformation) with clinopyroxene (cpx) as lamellar exsolution. Tremolite (tr) and talc (tlc) present as rim alteration of orthopyroxene (opx). Olivine (ol) is partially altered to serpentine (srp), and in contact with orthopyroxene (opx), olivine ultramylonite (uml) is formed due to intensive deformation processes. (c) Lherzolite (HUKO H16) shows some chromite-spinel (cr-spl) crystals coexisting with serpentinized olivine (ol). Clinopyroxene (cpx) appears to be replaced by hornblende (hbl). (d) Harzburgite (SOP H12), with olivine (ol) partially altered to serpentine (srp), showing serpentine (srp) mesh texture. (e) Lamellar exsolution of orthopyroxene (opx) and clinopyroxene (cpx) in Lherzolite (HUKO H16). (f) Lherzolite (VALE 14) showing a chromite spinel-bearing (cr-spl) serpentinized lherzolite. Serpentine (srp) replaces olivine (ol) and exhibits a mesh texture. Chlorite (chl) is present as an alteration mineral. Magnetite (mag) is abundantly distributed and associated with serpentine veins. This magnetite distribution indicates a high degree of serpentinization (Ray et al., 2010) (g) Cumulate orthopyroxene (opx), clinopyroxene (cpx) and olivine (ol) in olivine websterite VALE 02, orthopyroxene (opx) is seen as protomylonite (red arrow) with wavy blackout features, and visible mylonite structure in red circle. (h) Irregular chromite spinel (cr-spl) coexisting with serpentinized olivine (ol) and enstatite-type orthopyroxene (opx) in sample VALE 02.

Commented [S22]: hornblend ? atau tremolit ?

Commented [S23]: talc bukan kelompok ultra basa, juga serpentin dan mineral lainnya

Commented [S24]: klorit juga belum dijelaskan sebelumnya. dan bukan termasuk mineral penyusun batuan ultra basa

Table 2 Chemical data of coexisting olivine (ol) and spinel (spl) minerals (HUKO H16)

HUKO H16	Ol	Spl	Ol	Spl	Ol	Spl	Ol	Spl
SiO ₂	40,00	0,30	36,32	0,38	38,57	0,47	44,58	0,81
TiO ₂	0,00	0,00	0,00	0,03	0,00	0,03	0,00	0,00
Al ₂ O ₃	0,00	53,64	0,22	49,72	0,43	53,11	0,15	49,16
Cr ₂ O ₃	1,22	11,85	0,02	17,86	0,08	15,25	0,02	15,69
FeO	8,45	12,89	12,86	13,80	10,70	12,37	7,12	11,37
MnO	0,07	0,27	0,07	0,15	0,02	0,11	0,10	0,34
MgO	51,30	20,31	50,34	17,09	48,74	17,97	46,85	21,68
NiO	0,19	0,00	0,31	0,00	0,25	0,00	0,36	0,00
CaO	0,01	0,07	0,28	0,00	0,28	0,00	0,25	0,01
Na ₂ O	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
K ₂ O	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
ZnO	0,00	0,22	0,00	0,00	0,00	0,00	0,00	0,26
V ₂ O ₃	0,00	0,02	0,00	0,00	0,00	0,00	0,00	0,00
Total	101,24	99,57	100,42	99,03	99,07	99,31	99,43	99,32
Si	0,96	0,01	0,89	0,01	0,95	0,01	1,10	0,02
Ti	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Al	0,00	1,66	0,01	1,59	0,01	1,67	0,00	1,53
Cr	0,02	0,25	0,00	0,38	0,00	0,32	0,00	0,33
Fe ³⁺	0,00	0,08	0,00	0,00	0,00	0,00	0,00	0,09
Fe ²⁺	0,17	0,21	0,26	0,31	0,21	0,28	0,15	0,16
Mn	0,00	0,01	0,00	0,00	0,00	0,00	0,00	0,01
Mg	1,84	0,80	1,83	0,69	1,80	0,72	1,73	0,86
Ni	0,00	0,00	0,01	0,00	0,00	0,00	0,01	0,00
Ca	0,00	0,00	0,01	0,00	0,01	0,00	0,01	0,00
Na	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
K	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Zn	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,01
V	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Total	3,00	3,00	3,00	3,00	3,00	3,00	3,00	3,00
Fo	0,91		0,87		0,89		0,92	
Cr/(Cr+Al)		0,13		0,19		0,16		0,18
Cr/(Cr+Al+Fe ³⁺)		0,12		0,19		0,16		0,17
Al/(Cr+Al+Fe ³⁺)		0,84		0,81		0,84		0,78
Fe ³⁺ /(Cr+Al+Fe ³⁺)		0,04		0,00		0,00		0,05
Mg/(Mg+Fe ³⁺)		0,91		1,00		1,00		0,90

6 DISCUSSION

6.1 Geothermometer

The geothermometer or formation temperature of ultramafic rocks in this study was determined using lamellar exsolution between orthopyroxene and clinopyroxene (Lindsley, 1983). The formation temperature of ultramafic rocks can be obtained through lamellar exsolution of orthopyroxene and clinopyroxene, provided that the content of wolastony + enstatite + ferrophylite amounts to ≥90%. The diagram used in determining the formation temperature is the enstatite-diopsid-hedenbergite-ferrosilite (En-Di-Hd-Fs) quadrilateral diagram with a pressure of 5 kbar (Lindsley and Andersen, 2012). The quadrilateral diagram of En-Di-Hd-Fs at a pressure of 5 kbar is used based on the research

of Kadarusman et al. (2004) on the metamorphism process of amphibolite facies in the ophiolite complex in Sulawesi that occurred at a pressure of about 4 kbar.

Based on the %mol En, Fs, Wo data of orthopyroxene and clinopyroxene lamellar exsolution in Table 3 plotted in the En-Di-Hd-Fs quadrilateral diagram at a pressure of 5 kbar, three temperature groups were obtained. These three temperature groups are located on different temperature lines (Figure 3). Low-temperature rock formation data has a temperature interval of 500-800°C. Medium temperature rock formation data which has a temperature interval of 800-1000°C. High-temperature rock formation data has a temperature interval of 1000-1200°C.

Commented [S25]: penggunaan metode ini dikaji ulang kembali karena batuan ultra basa telah mengalami lebih dari satu kali proses geologi

Table 3 Chemical data of orthopyroxene (opx) and clinopyroxene (cpx) minerals forming lamellar exsolution (PMS 01)

PMS 01	Cpx1	Opx1	Cpx2	Opx2	Cpx3	Opx3	Cpx4	Opx4
SiO ₂	52,02	55,77	52,94	53,31	52,30	55,64	52,86	54,95
TiO ₂	0,70	0,25	0,58	0,26	0,56	0,11	0,65	0,28

Al ₂ O ₃	4,04	3,66	3,02	3,98	3,49	3,43	3,98	4,17
Cr ₂ O ₃	1,16	1,06	0,80	0,59	0,83	0,76	1,27	0,89
FeO	2,93	6,77	2,80	6,79	4,01	6,08	3,16	6,02
MnO	0,05	0,06	0,02	0,11	0,10	0,10	0,05	0,09
MgO	17,39	30,55	15,88	34,25	20,19	32,23	17,89	32,81
NiO	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
CaO	20,95	1,00	22,65	1,51	19,65	1,48	21,64	1,21
Na ₂ O	0,26	0,00	0,18	0,01	0,00	0,13	0,21	0,01
K ₂ O	0,02	0,00	0,01	0,02	0,00	0,01	0,02	0,00
ZnO	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
V ₂ O ₃	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Total	99,52	99,12	98,88	100,83	101,13	99,97	101,73	100,43
Si	1,90	1,97	1,96	1,82	1,87	1,93	1,89	1,89
Ti	0,02	0,01	0,02	0,01	0,02	0,00	0,02	0,01
Al	0,17	0,15	0,13	0,16	0,15	0,14	0,17	0,17
Cr	0,03	0,03	0,02	0,02	0,02	0,02	0,04	0,02
Fe ³⁺	0,00	0,00	0,00	0,17	0,07	0,00	0,00	0,01
Fe ²⁺	0,09	0,20	0,09	0,02	0,05	0,18	0,09	0,17
Mn	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Mg	0,95	1,61	0,88	1,74	1,07	1,67	0,95	1,68
Ni	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Ca	0,82	0,04	0,90	0,06	0,75	0,05	0,83	0,04
Na	0,02	0,00	0,01	0,00	0,00	0,01	0,01	0,00
K	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Zn	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
V	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Total	4,00	4,00	4,00	4,00	4,00	3,99	4,00	4,00
En	51,01	87,12	47,08	87,50	55,22	87,81	50,80	88,54
Fs	4,82	10,83	4,66	9,73	6,15	9,29	5,03	9,11
Wo	44,17	2,05	48,26	2,77	38,63	2,90	44,16	2,35
Mg#	0,91	0,89	0,91	0,99	0,95	0,90	0,91	0,91




Figure 3 Plot of %mol En, Fs, Wo on Lindsley (1983) En-Di-Hd-Fs quadrilateral diagram at 5 kbar pressure

The low-temperature rock formation data, which has a temperature interval of 500-800°C, is interpreted as the formation temperature of the milonite structure. The lamellar texture shown in the low-temperature data (Figure 4) is interpreted to have been disturbed by deformation. Deformation is characterized by the irregular shape of the lamellae and wider dimensions compared to other lamellae. The addition of calcium (Ca) influences the irregular shape and dimension of lamellae. The medium-

temperature rock formation data, which has a temperature interval of 800-1000°C, is interpreted as the formation temperature of the ultramylonite structure. The lamellar texture that has been deformed is broader compared to the lamellar texture at high temperatures. The high-temperature rock formation data, with a temperature interval of 1000-1200°C, is interpreted as the initial temperature of ultramafic rock formation. The lamellar texture shown in this high-temperature data has not been disturbed by deformation and is characterized by a thin and straight lamellar.




Figure 4 SEM EDS observation of point PMS 01 showing some lamellar textures

6.2 Petrogenesis of Ultramafic Rocks

According to petrographic analysis, peridotite dominates ultramafic rocks in the study area. The rocks found are peridotite (harzburgite and lherzolite) and pyroxenite (olivine websterite). The absence of garnet in the ultramafic rocks of the Baula and Pomalaa Ophiolite Complexes indicates that the ultramafic rocks have yet to undergo a high degree of metamorphism. The dominant presence of serpentines in all samples indicates intensive metamorphism at low temperatures. Chlorite and serpentine characterize the continuation of retrograde metamorphism at low temperatures (Frost et al., 2013, Arai, 1994). The study area also found tremolite, talc, and hornblende as alteration minerals of pyroxene. The formation of amphiboles (tremolite and hornblende) and talc requires H₂O and SiO₂ (Maulana et al., 2015). Serpentine+tremolite+magnetite+talc, especially at the pyroxene rim, indicates a reaction between pyroxene minerals and seawater (Zeng et al., 2012). Another source could also be the decomposition of water from previous serpentine formations. These processes likely occurred during the uplift of the ophiolite from the upper mantle to the surface (Arai et al., 2008). The absence of plagioclase in ultramafic rocks indicates that the uplift process of these ultramafic rocks occurred rapidly (Maulana et al., 2015).

The tectonic environment of ultramafic rocks (Figure 5) was determined by putting the Cr# spinel and Fo olivine data from the mineral chemistry of coexisting olivine and spinel (Table 2) onto the Olivine Spinel Mantle Array plot (Arai, 1994). Based on these data, it can be seen that the pattern of olivine spinel chemical data in the study area is similar to the four types of theoretical tectonic environments: ocean floor, oceanic hotspot, Japan arcs, and continent. Based on the chemical data pattern of these tectonic environments, in the oceanic hotspot type, the weight percent of TiO₂ is ~1 wt%. The Japan arcs type has several Cr# values >0.5 and Fo <0.87. In the continent type, there are Cr# values >0.5 even up to 0.75; in the ocean floor type, the Fo values range from 0.88-0.92 and Cr# 0.1-0.6. The chemical data distribution of the study area has TiO₂ ~0.03 wt%, Fo 0.87-0.92, Cr# 0.13-0.19, and none of Fo <0.87. Based on the distribution pattern of mineral chemical data, it can be interpreted that the ultramafic rocks in the study area are suitable for the ocean floor type.

The Al₂O₃ vs TiO₂ chemical data of the peridotite spinel minerals of the study area were plotted using plot from Kamenetsky et al. (2001) to determine the tectonic setting of the peridotite rocks. The data was then compared with spinel mineral data on Kabaena and Ampana lherzolites (Kadarusman et al., 2004), which have similar distribution patterns (Figure 6). Al₂O₃ spinel data distribution in the study area is 46.16-53.64 wt% and TiO₂ ~0.03 wt%. Based on these data, ultramafic rocks are interpreted as the MORB peridotite type.




Figure 5 Plot of Cr# spinel vs Fo olivine of ultramafic rocks of Baula-Pomalaa Ophiolite Complex using OSMA diagram. The red shading color is the result of this study and the yellow shading color is the data from Arai (1994).




Figure 6 Plot Al₂O₃ vs TiO₂ in spinel peridotite of the study area using graphs from Kamenetsky et al. (2001) compared with data of Al₂O₃ vs TiO₂ in spinel peridotite of Kabaena and Ampana from Kadarusman et al. (2004)

6.3 Tectonic Setting

Based on petrographic analysis, the Baula Pomalaa Ophiolite Complex comprises only ultramafic rocks. The ultramafic rocks encountered are peridotite (harzburgite and lherzolite) and pyroxenite (olivine websterite). The absence of gabbro, basalt, sheeted dykes, and other ophiolite sequences indicates that the ultramafic rocks in the study area have undergone a very significant tectonic process and produced dismembered ophiolitic sequence (Cendrajaya, 2017, Syahrul, 2017, Maulana et al., 2015).

The speed of the ocean floor spreading results in different characteristics of the ophiolites (Poli and Schmidt, 2002, Pearce et al., 1984). The ophiolite complexes formed through fast-spreading centers are characterized by harzburgite, poor clinopyroxene, and dunite (Maulana et al., 2015). The Baula Pomalaa Ophiolite Complex generally consists of harzburgite and lherzolite with abundant

Commented [S26]: focus penelitian sebenarnya menjelaskan magmatisme atau metamorfisme..?

Commented [S27]: terlebih pendapat ini...serpentin termasuk batuan metamorf atau batuan alterasi

Commented [S28]: proses metamorf dan alterasi pasca ultra basa terjadinya bagaimana ?

Commented [S29]: proses di dasar laut, kapan termetamorfkan dan berubah ? bagaimana penjelasannya ?

Commented [S30]: hindari penjelasan bukti negative (Tidak adanya...)

clinopyroxene minerals. The dismembered ophiolitic sequence and the abundance of clinopyroxene in the ultramafic rocks indicate that the ophiolite sequence formed through slow spreading center mechanism (Gong et al., 2016, Maulana et al., 2015). The pyroxenite (olivine websterite) also indicates that the Baula Pomalaa Ophiolite Complex formed at a slow spreading center.

The formation process of ultramafic rocks can be interpreted based on the formation temperature of lamellar exsolution between orthopyroxene and clinopyroxene minerals. Ultramafic rocks in the study area formed at a temperature of ~1200°C. This temperature is the crystallization temperature of pyroxene minerals. The mylonite and ultramylonite structures found were formed at different pressures and temperatures. During the emplacement of ultramafic rock mass onto the metamorphic basement, mylonite structures were formed at a temperature of ~500-800°C, and under other conditions ultramylonite structures were also formed at a temperature of ~800-1000°C.

The chemical data of the ultramafic rocks of the Baula Pomalaa Ophiolite Complex show some similarities with several other ophiolite complexes on Sulawesi Island. The ultramafic rocks of the Southeast Arm of Sulawesi are part of the Cretaceous-aged East Sulawesi Ophiolite (ESO) (Surono, 2013). The data distribution plot of Al₂O₃ vs TiO₂ spinel peridotite of the study area has similarities with the Kabaena and Ampana patterns, which have previously been studied by Kadarusman et al. (2004) and belongs to the East Sulawesi Ophiolite, i.e. the peridotite at the site is sourced from the MORB peridotite. Based on the data plot of Cr# vs Fo and Al₂O₃ vs TiO₂ (Figure 6), it can be concluded that the tectonic setting of ultramafic rocks in the study area is ocean floor with magmatism sourced from MORB (Mid Oceanic Ridge Basalt).

7 Conclusion

Ultramafic rocks in the Baula-Pomalaa Ophiolite Complex are dominated by peridotite (harzburgite, lherzolite) and pyroxenite (olivine websterite). Ultramafic rocks have undergone serpentinization and low-degree metamorphism as evidenced by serpentine, talc, and amphibole minerals (tremolite and hornblende). Based on the analysis results, the formation temperature of ultramafic rocks is divided into three groups, i.e., high temperature (1000-1200°C) as the initial rock formation temperature characterized by thin elongated lamellas. Intermediate temperatures (800-1000°C) are characterized by a relatively wide type of lamellae in orthopyroxene. The presence of ultramylonite structures also characterizes this phase. Anhedral and wider irregular lamella types characterize low temperatures (500-800°C). The presence of mylonite structures also characterizes this phase. Coexisting olivine and spinel analysis shows a distribution of Fo values ranging from 0.87-0.92 and Cr# values ranging from 0.13-0.19. The results of the tectonic setting plot on the Olivine-Spinel Mantle Array (OSMA) obtained that the tectonic setting of ultramafic rocks in the study area belongs to the ocean floor with magmatism sourced from MORB (Mid Oceanic Ridge Basalt).

8 Acknowledgements

This research is supported by Halu Oleo University through the Institute for Research and Community Services Funding, through the 2023 Penelitian Dosen pemula (PDP) Research Scheme.

References

- Arai, S. 1994. Characterization of spinel peridotites by olivine-spinel compositional relationships: Review and interpretation. *Chemical Geology*, 113(3-4). 191-204. [https://doi.org/10.1016/0009-2541\(94\)90066-3](https://doi.org/10.1016/0009-2541(94)90066-3)
- Arai, S., Tamura, A., Ishimaru, S., Kadoshima, K., Lee, Y.-I. & Hisada, K.-i. 2008. Petrology of the Yugu peridotites in the Gyeonggi Massif, South Korea: Implications for its origin and hydration process. *Island Arc*, 17(4). 485-501. <https://doi.org/10.1111/j.1440-1738.2008.00633.x>
- Burnley, P. C., Cline, C. J. & Drue, A. 2013. Kinking in Mg₂GeO₄ olivine: An EBSD study. *American Mineralogist*, 98(5-6). 927-931. <https://doi.org/10.2138/am.2013.4224>
- Cendrajaya, R. I. M. 2017. Studi petrogenesis dan mineralisasi Kompleks Ofolit Baula, Kolaka, Sulawesi Tenggara. Thesis, Institut Teknologi Bandung.
- Drue, A. G. 2011. Microstructural characterization of kinked germanate olivine grains. Theses, University of Nevada.
- Frost, B. R., Evans, K. A., Swapp, S. M., Beard, J. S. & Mothersole, F. E. 2013. The process of serpentinization in dunite from New Caledonia. *Lithos*, 178. 24-39. <https://doi.org/10.1016/j.lithos.2013.02.002>
- Gong, X.-H., Shi, R.-D., Griffin, W. L., Huang, Q.-S., Xiong, Q., Chen, S.-S., Zhang, M. & O'Reilly, S. Y. 2016. Recycling of ancient subduction-modified mantle domains in the Purang ophiolite (southwestern Tibet). *Lithos*, 262. 11-26. <https://doi.org/10.1016/j.lithos.2016.06.025>
- Hamilton, W. B. 1979. Tectonics of the Indonesian Region (Report No. 1078), Professional Paper, U.S. Govt. Print. Off. 345 p. <https://doi.org/10.3133/pp1078>
- Husein, S., Novian, M. I. & Barianto, D. H. 2014. Geological structures and tectonic reconstruction of Luwuk, East Sulawesi Proceedings, Indonesian Petroleum Association: Thirty-Eight Annual Convention & Exhibition Jakarta, Indonesian Petroleum Association. IPA 14-G-137.
- Kadarusman, A., Miyashita, S., Maruyama, S., Parkinson, C. D. & Ishikawa, A. 2004. Petrology, geochemistry and paleogeographic reconstruction of the East Sulawesi Ophiolite, Indonesia. *Tectonophysics*, 392(1-4). 55-83. <https://doi.org/10.1016/j.tecto.2004.04.008>
- Kamenetsky, V. S., Crawford, A. J. & Meffre, S. 2001. Factors Controlling Chemistry of Magmatic Spinel: an Empirical Study of Associated Olivine, Cr-spinel and Melt Inclusions from Primitive Rocks. *Journal of Petrology*, 42(4). 655-671. <https://doi.org/10.1093/petrology/42.4.655>
- Koizumi, T., Tsunogae, T., Santosh, M., Tsutsumi, Y., Chetty, T. R. K. & Saitoh, Y. 2014. Petrology and zircon U-Pb geochronology of metagabbros from a mafic-ultramafic suite at Aniyapuram: Neoproterozoic to Early Paleoproterozoic convergent margin magmatism and Middle Neoproterozoic high-grade metamorphism in southern India. *Journal of Asian Earth Sciences*, 95. 51-64. <https://doi.org/10.1016/j.jseas.2014.04.013>
- Lindsley, D. H. 1983. Pyroxene thermometry. *American Mineralogist*, 68(5-6). 477-493.
- Lindsley, D. H. & Andersen, D. J. 2012. A two-pyroxene thermometer. *Journal of Geophysical Research: Solid Earth*, 117. 1-12. <https://doi.org/10.1029/2011JB008799>

Commented [S31]: apa jenis mineralnya..? agar lebih spesifik

Commented [S32]: perlu dijelaskan lebih detail, karena ada proses lain setelah pembentukan ultra basa

Commented [S33]: bagaimanakah paragenesa dari temperature 1200, 500-800 dan 800-1000 °C. Proses geologi seperti apa..tidak dijelaskan

Commented [S34]: Jelaskan dalam sari

Commented [S35]: menurut saya, hal ini bukan pembentukan jenis ultra basa, tetapi batuan ultra basa daerah penelitian mengalami beberapa kali proses geologi yaitu batuan beku, metamor dan mungkin batuan ubahan. Berdasarkan ciri-ciri kehadiran mineraloginya.

- Earth, 88(S02). A887-A906. <https://doi.org/10.1029/JB088iS02p0A887>
- Matysiak, A. K. & Trepmann, C. A. 2015. The deformation record of olivine in mylonitic peridotites from the Finero Complex, Ivrea Zone: Separate deformation cycles during exhumation. *Tectonics*, 34(12). 2514-2533. <https://doi.org/10.1002/2015tc003904>
- Maulana, A., Christy, A. G. & Ellis, D. J. 2015. Petrology, geochemistry and tectonic significance of serpentinitized ultramafic rocks from the South Arm of Sulawesi, Indonesia. *Geochemistry*, 75(1). 73-87. <https://doi.org/10.1016/j.chemer.2014.09.003>
- Monnier, C., Girardeau, J., Maury, R. C. & Cotten, J. 1995. Back-arc basin origin for the East Sulawesi ophiolite (eastern Indonesia). *Geology*, 23(9). 851-854. [https://doi.org/10.1130/0091-7613\(1995\)023<0851:Baboft>2.3.Co;2](https://doi.org/10.1130/0091-7613(1995)023<0851:Baboft>2.3.Co;2)
- Olfindo, V. S. V., Payot, B. D., Valera, G. T. V. & Arai, S. 2020. Petrogenesis of heterogeneous mantle peridotites with Ni-rich olivine from the Pujada Ophiolite, Philippines. *Journal of Asian Earth Sciences*: X, 4. <https://doi.org/10.1016/j.jaesx.2020.100039>
- Parkinson, C. 1998. Emplacement of the East Sulawesi Ophiolite: evidence from subophiolite metamorphic rocks. *Journal of Asian Earth Sciences*, 16(1). 13-28. [https://doi.org/10.1016/s0743-9547\(97\)00039-1](https://doi.org/10.1016/s0743-9547(97)00039-1)
- Payot, B., Arai, S., Yoshikawa, M., Tamura, A., Okuno, M. & Rivera, D. 2018. Mantle Evolution from Ocean to Arc: The Record in Spinel Peridotite Xenoliths in Mt. Pinatubo, Philippines. *Minerals*, 8(11). <https://doi.org/10.3390/min8110515>
- Pearce, J. A., Lippard, S. J. & Roberts, S. 1984. Characteristics and tectonic significance of supra-subduction zone ophiolites. Geological Society, London, Special Publications, 16(1). 77-94. <https://doi.org/10.1144/gsl.Sp.1984.016.01.06>
- Poli, S. & Schmidt, M. W. 2002. Petrology of Subducted Slabs. *Annual Review of Earth and Planetary Sciences*, 30(1). 207-235. <https://doi.org/10.1146/annurev.earth.30.091201.140550>
- Ray, D., Banerjee, R., Iyer, S. D. & Mukhopadhyay, S. 2010. A New Report of erpentinites from Northern Central Indian Ridge (at 6°S)-An Implication for Hydrothermal Activity. *Acta Geologica Sinica - English Edition*, 82(6). 1213-1222. <https://doi.org/10.1111/j.1755-6724.2008.tb00723.x>
- Simandjuntak, T. O., Surono & Sukido 1993. Peta Geologi Lembar Kolaka, Sulawesi, Bandung, Pusat Penelitian dan Pengembangan Geologi.p.
- Streckeisen, A. 1976. To each plutonic rock its proper name. *Earth-Science Reviews*, 12(1). 1-33. [https://doi.org/10.1016/0012-8252\(76\)90052-0](https://doi.org/10.1016/0012-8252(76)90052-0)
- Surono 2013. Geologi Lengan Tenggara Sulawesi, Bandung, Badan Geologi. 201 p.
- Surono & Hartono, U. 2013. Geologi Sulawesi, Jakarta, LIPI Press. 352 p.
- Syahrul. 2017. Studi Paragenesis dan Mineralisasi kompleks Batuan Ultramafik daerah Sopura, Kabupaten Kolaka, Provinsi Sulawesi Tenggara. Theses, Institut Teknologi Bandung.
- van Zuidam, R. A. 1986. Aerial photo-interpretation in terrain analysis and geomorphologic mapping, Netherland, Publisher The Hague.p.
- Whitney, D. L. & Evans, B. W. 2009. Abbreviations for names of rock-forming minerals *American Mineralogist*, 95(1). 185-187. <https://doi.org/10.2138/am.2010.3371>
- Yellappa, T., Koizumi, T. & Tsunogae, T. 2021. Geochemical constrains on pyroxenites from Aniyapuram Mafic-Ultramafic Complex, Cauvery Suture Zone, southern India: Suprasubduction zone origin. *Journal of Earth System Science*, 130(1). <https://doi.org/10.1007/s12040-020-01516-8>
- Zeng, Z., Wang, Q., Wang, X., Chen, S., Yin, X. & Li, Z. 2012. Geochemistry of abyssal peridotites from the super slow-spreading Southwest Indian Ridge near 65°E: Implications for magma source and seawater alteration. *Journal of Earth System Science*, 121(5). 1317-1336. <https://doi.org/10.1007/s12040-012-0229-z>



© 2016 Journal of Geoscience, Engineering, Environment and Technology. All rights reserved. This is an open access article distributed under the terms of the CC BY-SA License (<http://creativecommons.org/licenses/by-sa/4.0/>)

Commented [S36]: pustaka disesuaikan dengan kebutuhan JGEET. Diperbanyak pustaka 5 tahun terakhir



KEMENTERIAN PENDIDIKAN, KEBUDAYAAN, RISET, DAN TEKNOLOGI
UNIVERSITAS PEMBANGUNAN NASIONAL "VETERAN" YOGYAKARTA
FAKULTAS TEKNOLOGI MINERAL
Jl. Pajajaran Condongcatur, Yogyakarta 55283
Telp. (0274) 487813, 487814, Fax. (0274) 487813

SURAT TUGAS

Nomor : B/287d/UN62.11/ST/2023

Dekan Fakultas Teknologi Mineral Universitas Pembangunan Nasional "Veteran" Yogyakarta memberikan tugas kepada :

No.	Nama	NIP	Pangkat/Golongan
1.	Dr.Ir. Dwi Fitri Yudiantoro, MT	19630225 199003 1 002	Pembina Utama Muda /IVC

Keperluan: Untuk melaksanakan tugas dalam kegiatan sebagai **Reviewer Journal** pada **Journal of Geoscience, Engineering, Environment, and Technology (JGEET)** dengan judul artikel: "**Petrology and Mineral Chemistry of Ultramafic Rock in Baula - Pomalaa Ophiolite Complex, Southeast Sulawesi**"

Atas perhatian dan bantuannya kami ucapkan terima kasih.

18 Desember 2023



Dr. Ir. Sutarto, MT

NIP. 19650301199103 1 001