

REFERENCES

- Abad, M.J., Bedoya, L.M., Apaza, L., and Bermejo, P., 2012. *The Artemisia genus: a review of bioactive essential oils*. *Molecules* 17, 2542–2566.
- Agnello, A.M. 2002. Petroleum-derived Spray Oils: Chemistry, History, Refining and Formulation, p. 2-18.
- Alizadeh M., Aghaei M., Sharifian I., and Saadatian M. (2012). Chemical composition of essential oil of *Artemisia vulgaris* from West Azerbaijan, Iran. *Electron. J. Environ. Agric. Food Chem.* 11 493–496
- Appah, S., Jia, W., Ou, M., Wang, P., and Asante, E. 2020. Analysis of potential impaction and phytotoxicity of surfactant-plant surface interaction in pesticide application. *Crop Protection*, 127, 104961.
- Arshad, M., Ullah, M., Çağatay, N., Dikmen, F., Abdullah, A., and Afzal, M. (2019). *Cordia myxa* L.1, a New Host Plant Record for Asian Citrus Psyllid, *Diaphorina citri* Kuwayama2. *Southwestern Entomologist*, 44, 331 - 334.

Aslan, I., Kordali, S., and Calmasur, O., 2005. Toxicity of the vapours of *Artemisia absinthium* essential oil to *Tetranychus urticae* Koch and *Bemisia tabaci* Genn. Fresen. *Environ. Bull.* 14, 413–417.

Azam, M., Jiang, Q., Zhang, B., Xu, C., and Chen, K. 2013. Citrus Leaf Volatiles as Affected by Developmental Stage and Genetic Type. *Int J Mol Sci.* 14(9): 17744–17766.

Balabramani, S., Sabaphati, G. Moola, A, K., Solomon, R, V., Venuvanalingam, P., and Diana, R, K, B. 2018. *Evaluation of the Leaf Essential Oil from Artemisia vulgaris and Its Larvicidal and Repellent Activity against Dengue Fever Vector Aedes aegypti An Experimental and Molecular Docking Investigation.* Article. Journal ACS OMEGA. India.

Bano, A., Ahmad, M., Hadda, T.B., Saboor, A., Sultana, S., Zafar, M., Khan, M.P.Z., Arshad, M., and Ashraf, M.A., 2014. Quantitative ethnomedicinal study of plants used in the Skardu valley at high altitude of Karakoram-Himalayan range, Pakistan. *J. Ethnobiol. Ethnomed.* 10, 43–52.

Beattie, G.A.C. 2005. Using Petroleum-based Spray Oils in Citrus. *Agfact H2.AE.5.*

Benelli, G., 2015. Plant-borne ovicides in the fight against mosquito vectors of medical and veterinary importance: a systematic review. *Parasitol. Res.* 114, 3201–3212.

Borgoni PC, Vendramim JD, Lourenção AL, and Machado MA. 2014. Resistance of Citrus and related genera to *Diaphorina citri* Kuwayama (Hemiptera: Liviidae). *Neotrop Entomol* 43:465–46.

Borgoni PC, Vendramim JD, Lourenção AL, and Machado MA. 2014. Resistance of Citrus and related genera to *Diaphorina citri* Kuwayama (Hemiptera: Liviidae). *Neotrop Entomol* 43:465–469.

Bove JM. 2006. Huanglongbing: A Destructive, Newly-emerging, Century-old Disease of Citrus. *J of Plant Pathology*.

Bruce, T.J.sA., L.J. Wadhams and C.M. Woodcock. 2005. Insect Host Location: a Volatile Situation. *Trends in Plant Science*. 10: 269-274.

Caboni, P. 2002. Persistence of azadirachtin residues on olives after field treatment. *Journal of Agriculture and Food Chemistry*, 50:3491-3494, 2002.

Caboni, P. 2006. Residues and persistence of neem formulations on strawberry after field treatment., *Journal of Agriculture and Food Chemistry* 54:10026-10032.

Campbell, S.A. and J.H. Borden. 2006. Close-range, Inflight Integration of Visual and Olfactory Information by a Host-seeking Bark Beetle. *Entomologia Experimentalis et Applicata*. 120: 91-98.

Campos EV, de Oliveira JL, Pascoli M, de Lima R, and Fraceto LF. Neem Oil and Crop Protection: From Now to the Future. *Front Plant Sci*. 2016

Chiasson, H., Belanger, A., Bostanian, N., Vincent, C., and Poliquin, A., 2001. Acaricidal properties of *Artemisia absinthium* and *Tanacetum vulgare* (*Asteraceae*) essential oils obtained by three methods of extraction. *J. Econ. Entomol.* 94, 167e171.

Gonzalo O. D., Vioque. R.S. Berruga M.i., Penalver D. H., and Meridaz O.S. Antifeedant effects of common terpenes from Mediterranean aromatic plants on *Leptinotarsa decemlineata*. *J. Soil Sci. Plant Nutr.* vol.17 no.2

Gottwald TR, Graham JH, Irey M, McCollum G., and Wood B (2012) Inconsequential effect of nutritional treatments on

huanglongbing control, fruit quality, bacterial titer and disease progress. *Crop Protection* **36**: 72–82.

Govindarajan, M., Benelli, G., 2016. *Artemisia absinthium*-borne compounds as novel larvicides: effectiveness against six mosquito vectors and acute toxicity on non-target aquatic organisms. *Parasitol. Res.* 115, 4649–4661.

Greenberg, S.M., Showler, A.T., and Liu, T.X., 2005. Effects of neem-based insecticides on beet armyworm (Lepidoptera: Noctuidae). *Insect Sci* 12, 17 – 23.

Halbert, S. E., C. L. Niblett, K. L. Manjunath, R. F. Lee, and L. G. Brown. 2002. Establishment of two new vectors of citrus pathogens in Florida. *Proceeding of the international Society of Citriculture 9th Congress*. Alexandria, Va: ASHS Press.

Hall DG, Hentz MG, and Ciomperlik MA. A comparison of traps and stem tap sampling for monitoring adult Asian citrus psyllid (Hemiptera: Psyllidae) in citrus. *Fla. Entomol.* 2007; 90(2): 327–334.

Hall DG .2008b. Biological control of *Diaphorina citri*. North American Plant Protection Organization, Proceedings of the Taller Internacional Sobre

Huanglongbing y el Psilido Asiatico de los Citricos, del 7 al 9 de Mayo de 2008, Hermosillo, Sonora, p. 7.

Hall DG. 2010. *Biology, history and world status of Diaphorina citri*. North American Plant Protection Organization, *Proceedings of the Second Taller Internacional Sobre Huanglongbing y el Psilido Asiatico de los Citricos, 19–23 July, 2010*, pp. 8–36.

Hayat, M.Q., Ashraf, M., Khan, M.A., Yasmin, G., Shaheen, N., and Jabeen, S., 2009. Phylogenetic analysis of genus *Artemisia* (Asteraceae) based on micromorphological traits of pollen grains. *Afr. J. Biotec.* 8, 6561–6568.

Henry, E., Carlson, C., and Kuo, Y. 2023. Candidatus *Kirkpatrickella diaphorinae* gen. nov., sp. nov., an uncultured endosymbiont identified in a population of *Diaphorina citri* from Hawaii. *International journal of systematic and evolutionary microbiology*, 73 11.

Husain, M.A. and Nath, D. 1927. The citrus psylla (*Diaphorina citri*, Kuw.) [Psyllidae: Homoptera]. *Memoirs of the Department of Agriculture in India. Entomological Series*, 10(2): 5-27.

- Ikeda K, and Ashihara W. Preference of adult Asian citrus psyllid, *Diaphorina citri* (Homoptera: Psyllidae) for *Murraya paniculata* and *Citrus unshiu*. *Jpn J App Entomol Z.* 2008;52:27–30.
- Jiang Z., Guo X., Zhang K., Sekaran G., Cao B., and Zhao Q. 2019. The essential oils and eucalyptol from *Artemisia vulgaris* L. prevent acetaminophen-induced liver injury by activating Nrf2-Keap1 and enhancing APAP clearance through non-toxic metabolic pathway. *Front. Pharmacol.* 10:782.
- Khan, I., Zahid, M., and Khan, G. 2012. Toxicity of botanic and synthetic pesticide residues to citrus psyllid *Diaphorina citri* Kuwayama and *Chrysoperla carnea* (Stephens).. *Pakistan Journal of Zoology*, 44, 197-201.
- Knudsen JT, and Tollsten L. 1993. Trends in floral scent chemistry in pollination syndromes—floral scent composition in moth-pollinated taxa. *Bot J Linn Soc* 113: 263–284.
- Kordali, S., Aslan, I., Çalmaşur, O., and Cakir, A., 2006. *Toxicity of essential oils isolated from three Artemisia species and some of their major components to granary weevil, Sitophilus granarius (L.)(Coleoptera: curculionidae)*. *Ind. Crop. Prod.* 23, 162–170.

Mann R. S, Ali J. G, Hermann S. L, Tiwari S, Pelz-Stelinski KS, and Alborn H. T. Induced release of a plant-defense volatile ‘deceptively’ attracts insect vectors to plants infected with a bacterial pathogen. *PLoS Pathog.* 2012; 8(3).

McClellan APD., and Schwartz RE (1970) Greening of blotchy-mottle disease in citrus. *Phytophylactica* 2: 177–194.

Mihajilov-Krstev, T., Jovanovic, B., Jovic, J., Ilic, B., Miladinovic, D., Matejic, J., Rajkovic, J., Dordevic, L., Cvetkovic, V., and Zlatkovic, B., 2014. Antimicrobial, antioxidative, and insect repellent effects of *Artemisia absinthium* essential oil. *Planta Med.* 80, 1698e1705.

Mordue (Luntz) A. J., and Nisbet A. J. Azadirachtin from the neem tree *Azadirachta indica*: its action against insects. *Anais da Sociedade Entomológica do Brasil.* 2000;29(4):615–632.

Mulla M. S., Su T. (1999). Activity and biological effects of neem products against arthropods of medical and veterinary importance. *J. Am. Mosq. Control Assoc.* 15 133–152.

Nicoletti, M., Petitto, V., Gallo, F. R., Multari, G., Federici, E., and Palazzino, G. 2012. *The modern analytical determination of botanicals and similar*

novel natural products by the HPTLC fingerprint approach. Stud. Nat. Prod. Chem. 37, 217–258.

Nile, A., Kwon, Y., & Nile, S. 2019. Horticultural oils: possible alternatives to chemical pesticides and insecticides. *Environmental Science and Pollution Research*, 1-13.

Ninditya VI, Purwati E, Utami AT, Marwaningtyaz AS, Fairuz NK, Widayanti R, Hamid PH. *Artemisia vulgaris* efficacies against various stages of *Aedes aegypti*. *Vet World*. 2020 Jul;13(7):1423-1429

Ouyang G., Fang X., Lu H. Zhou X., 2013. Repellency of Five Mineral Oils Against *Diaphorina citri* (Hemiptera: Liviidae). *Florida Entomologist*. 96(3):976-982.

Patt JM, Sétamou M. Responses of the Asian citrus psyllid to volatiles emitted by the flushing shoots of its rutaceous host plants. *Environ Entomol*. 2010;39:618–624.

Patt JM, Meikle WG, Mafra-Neto A, Sétamou M, Mangan R, Yang C, *et al*. Multimodal cues drive host-plant assessment in Asian citrus psyllid (*Diaphorina citri*). *Environ. Entomol*. 2011. 40(6): 1494–1502.

- Poerwanto, M. E., Trisyono. A. Y., Subandiya. S. Martono. E., Holford .P., Andrew. G., 2008. Effects Of Mineral Oils On Host Selection Behavior Of *Diaphorina citri*. *Perlindungan Tanaman Indonesia*, Vol.14, No.1 : 23-28.
- Rodriguez, N.J.A., Walter, G., and Mensah. R. K. 2007. The efficacy of a petroleum spray oil against *Aphis gossypii* Glover on cotton. Part 1: Mortality rates and sources of variation. *Pest management Science*. 63, 586–595.
- Tsagkarakis AE, Rogers ME. Suitability of Cleopatra mandarin as a host plant for *Diaphorina citri* (Hemiptera: Psyllidae) *Fla Entomol*. 2010;93:451–453.
- Santos, M., Zanardi, O., Pauli, K., Forim, M., Yamamoto, P., and Vendramim, J. 2015. Toxicity of an azadirachtin-based biopesticide on *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) and its ectoparasitoid *Tamarixia radiata* (Waterston) (Hymenoptera: Eulophidae). *Crop Protection*, 74, 116-123.
- Sumbe RB, Dhalgade MS, Palve KN, Kale SS, Waghe VL, Autade KA, and Dr. Kothawade SN. 2022. A Pharmacotherapeutic Screening of *Artemisia vulgaris* Whole Plant: A Brief Review. *Journal of Pharmacognosy and Phytochemistry*. 12(1): 298-302.

Titouhi, F., Amri, M., Messaoud, C., Haouel, S., Youssfi, S., Cherif, A., Jemâa, J.M.B. 2017. Protective effects of three Artemisia essential oils against *Callosobruchus maculatus* and *Bruchus rufimanus* (Coleoptera: chrysomelidae) and the extended sideeffects on their natural enemies. *J. Stored. Prod. Res.* 72, 11–20.

TSAI, J. H., and Y. H. LIU. 2000. Biology of *Diaphorina citri* (Homoptera: Psyllidae) on four host plants. *J. Econ. Entomol.* 93: 1721-1725.

Twidle. A.M., Barker. D., Pilkington. L. I., Fedrizi B. Suckling D.M. 2022. Identification Of Herbivore-Induced Plant Volatiles From Selected *Rubus* Species Fed Upon By Raspberry Bud Moth (*Heterocrossa rubophaga*) larvae. [Phytochemistry](#). 22: 0031-9422.

Venkateswarlu B, Pirat M, Kishore N, Rasul A. Mycorrhizal inoculation in neem (*Azadirachta indica*) enhances azadirachtin content in seed kernels. *World Journal Microbiology Bio technology*. 2008; 24:1243- 1247.

Wang, J.; Zhu, F.; Zhou, X. M.; Iu, C. Y. and Lei, C. L. 2006, Repellent and fumigant activity of essential oil from *Artemisia vulgaris* to *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *J. Stored Prod. Res.*, 42, 339-347.

Weathersbee, A. A., and C. L. McKenzie. "Effect of a Neem Biopesticide on Repellency, Mortality, Oviposition, and Development of *Diaphorina citri* (Homoptera: Psyllidae)." *The Florida Entomologist*, vol. 88, no. 4, 2005, pp. 401–07.

Wirawan, I. G. P., S. Simanjuntak, M. Sritamin, and N. Wijaya. 2017. Detection Of Citrus Vein Phloem Degeneration (Cvpd) Disease And The Quality Of Healthy Fruits In Nutrient Deficiency Of Citrus. *Bali Medical Journal (Bali Med J)*, Volume 6.

Sakamaki Y .2005. Possible migration of the Asian citrus psyllid, *Diaphorina citri* Kuwayama (Homoptera: Psyllidae) between and within islands. *Occasional Papers Kagoshima University Research Center for the Pacific Islands* 42: 121–125.

Setamou, M., A. Sanchez, J. M. Patt, S. D. Nelson, J. Jifon, and E. S. Louzada. 2011. Diurnal patterns of flight activity and effects of light on host finding behavior of the Asian citrus psyllid. *J. Insect Behav.* 25: 264–276.

Singh, N., Devi, M., Biona, T., Sharma, N., Das, S., Chakravorty, J., Mukherjee, P., & Rajashekar, Y. 2023. Phytochemical Composition and Antimicrobial

Activity of Essential Oil from the Leaves of *Artemisia vulgaris* L.
Molecules, 28.

Sui, S. Leong, S., Beattie, G. 2021. Effect of Horticultural Mineral Oil on Huanglongbing Transmission by *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) Population in a Commercial Citrus Orchard in Sarawak, Malaysia, Northern Borneo. *J, Insect*. MDPI.

You, Chunxue, Guo, Shanshan, Zhang, Wenjuan, Yang, Kai, Geng, Zhufeng, Du, Shushan, Wang, Chengfang, Deng, Zhiwei. 2015. Identification of Repellent and Insecticidal Constituents from *Artemisia mongolica* Essential Oil against *Lasioderma serricorne*, *Journal of Chemistry*, 2015, 549057, 7 pages,

Zhang, J., Li, B., Lu, X., Zheng, Y., Wang, D., Zhang, Z., Zeng, D., & Du, S. 2022. Chemical Diversity and Anti-Insect Activity Evaluation of Essential Oils Extracted from Five *Artemisia* Species. *Plants*, 11.

Zhang J, Nielsen SE, Chen Y, Georges D, Qin Y, Wang SS, Thuiller W. 2017. Extinction risk of north American seed plants elevated by climate and land-use change. *J Appl Ecol* 54(1):303–3